


CLEVELAND CLINIC JOURNAL OF MEDICINE



INTERNAL MEDICINE BOARD REVIEW

■ DAVID L. LONGWORTH AND JAMES K. STOLLER, EDITORS

The elevated peripheral white blood count 267

B. J. BOLWELL

HIGHLIGHTS FROM MEDICAL GRAND ROUNDS

■ WILLIAM S. WILKE, EDITOR

Parkinson's disease: Where do we stand? 269

Although a cure has not been found, certain drugs can relieve symptoms and may actually slow disease progression.

P. J. SWEENEY

Irritable bowel syndrome: new perspectives on management 270

Long classified as psychosomatic, irritable bowel syndrome is now considered a motor disorder in which a number of factors are at play.

M. M. SCHUSTER

CURRENT DRUG THERAPY

■ DONALD G. VIDT AND ALAN BAKST, EDITORS

The fifth report of the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure: insights and highlights from the chairman 273

Highlights include the new classification schema and recommendations on the use of labetalol.

R. W. GIFFORD, JR

Assessing hypertension management: the role of 24-hour blood pressure monitoring 278

The best way to evaluate antihypertensive drug therapy may be to use, in a complementary fashion, the various methods of analyzing ambulatory blood pressure data.

B. ESAYAG-TENDLER AND W. B. WHITE

From bathtub ring to osteoporosis: a clinical review of the bisphosphonates 284

Clinical trials have aroused widespread interest in the use of these drugs to treat—and perhaps even to prevent—osteoporosis.

A. A. LICATA

Continued on p. 265

Continued from p. 263

The use of G-CSF and GM-CSF in bone marrow transplantation 291

Using hematopoietic growth factors appears to have made bone marrow transplantation safer. Treatment-related morbidity and mortality have decreased, and patients are discharged from the hospital sooner.

B. J. BOLWELL

ORIGINAL STUDIES

Lung transplantation: the Cleveland Clinic experience 303

The early perioperative and 1-month posttransplantation periods appear critical to long-term survival. Complications are usually manageable and do not lead to long-term morbidity.

S. R. PATEL, T. J. KIRBY, P. M. MCCARTHY, AND OTHERS

Stereotactically guided thrombolysis of deep cerebral hemorrhage: preliminary results 321

Results of this study support others' findings that the volume of the hematoma in intracerebral hemorrhage can be safely and dramatically reduced without resorting to open surgery.

D. W. MILLER, G. H. BARNETT, D. W. KORMOS, AND OTHERS

REVIEWS

The pulmonary effects of free-base cocaine 325

Respiratory symptoms, pulmonary hemorrhage, pulmonary edema, asthma, pulmonary barotrauma, and other pulmonary effects of free-base cocaine are reviewed.

I. S. MEISELS AND J. LOKE

Parathyroid carcinoma: 50-year experience at The Cleveland Clinic Foundation 331

Long-term survival is possible with repeated resection of locally recurrent tumors.

A. G. HAKAIM AND C. B. ESSELSTYN, JR

CASE REPORT

Hemopericardium and cardiac tamponade associated with warfarin therapy 336

K. S. LEE AND T. MARWICK

DEPARTMENTS

CME calendar 266

Information for authors 320