



New options for diagnosing and treating acromegaly

ARIEL L. BARKAN, MD*

Professor of Internal Medicine, Division of Endocrinology and Metabolism,
and Professor of Surgery, University of Michigan Medical Center, Ann Arbor

ABSTRACT

Insulin-like growth factor 1 (IGF-1) is the final pathway of the growth hormone cascade, and is a better clinical measure of acromegaly than growth hormone itself. In patients with acromegaly, medical treatment with a somatostatin analog normalizes growth hormone and IGF-1 levels and produces clinical improvement quickly.

A CROMEGALY, caused by an overproduction of growth hormone, was once diagnosed primarily by the physical changes it produces, followed by measurement of growth hormone. Now, a more complete understanding of the growth hormone cascade has shown that insulin-like growth factor 1 (IGF-1) is a sensitive clinical marker of the severity of acromegaly. In addition, two somatostatin analogs (eg, octreotide, lanreotide) are now available as medical treatment.

EPIDEMIOLOGY OF ACROMEGALY

Acromegaly is a classic endocrine disorder. Indeed, growth hormone was the first pituitary hormone identified.

Acromegaly occurs in 3 to 4 persons per million per year. It is equally frequent in both sexes. The mean age at diagnosis is 40 years in men and 45 years in women.

This disease confers a mortality rate two to

three times higher than in healthy persons. The mean age at death is approximately 60 years. Deaths are primarily due to vascular disorders (eg, cardiomyopathy) and various cancers.

PHYSIOLOGY OF ACROMEGALY

Two hypothalamic hormones regulate growth hormone:

Growth hormone-releasing hormone (GHRH) stimulates the synthesis and release of growth hormone from the pituitary gland.

Somatostatin inhibits release of growth hormone, but not its synthesis (FIGURE 1).

When growth hormone is released into the peripheral circulation, it stimulates the production of IGF-1—the true growth hormone—by the liver and other organs. Growth hormone also has independent metabolic effects on lipids and glucose.

Most cases of acromegaly are due to pituitary adenomas. However, a few are due to ectopic GHRH-secreting tumors.

In a study that provides insight into how pituitary adenomas develop, Thapar et al¹ found that they express the gene for GHRH. The amount of GHRH mRNA present was proportionate to the tumor's size, proliferating ability, and invasiveness, and to the amount of growth hormone it secreted. Further, the amount of GHRH mRNA was the single most important factor that predicted the likelihood of a cure. In effect, pituitary adenomas can stimulate themselves to grow and secrete growth hormone.

DIAGNOSING ACROMEGALY

Acromegaly is suspected on the basis of the classic clinical signs and symptoms (TABLE 1).

The characteristic appearance of persons with acromegaly identifies the condition at a glance: enlarged hands and feet, oily skin, thick lips, a protruding jaw. In uncertain cases, ask the patient to bring in his or her photo album. Often, one can pinpoint the year of onset by comparing pictures from different years.

**IGF-1 is the
true growth
hormone**

*Dr. Barkan's lecture was sponsored by Novartis Pharmaceuticals Corporation.



Examination may reveal some of the complications of acromegaly: thyroid hypertrophy, cardiomegaly, hepatomegaly, splenomegaly, nephromegaly, hypertension, and osteoarthritis. If the tumor impinges on the optic chiasma, the patient may have visual field defects.

MRI scanning usually reveals a macroadenoma, often quite large. There is a rough correlation between size of tumor and levels of growth hormone.

Growth hormone levels may be misleading. Normally, growth hormone is mostly secreted during sleep in pulses approximately 2 hours apart, and it has a short half-life. Thus, serum levels fluctuate widely through the day. Growth hormone levels may be above the normal limit during pulses of secretion in normal persons; conversely, they may be within the stated normal range between pulses in persons with acromegaly.

IGF-1 levels are the single best tool for diagnosing and monitoring acromegaly, because plasma levels remain steady through the day and night. The normal range for IGF-1 varies among laboratories and is sex-dependent and age-dependent. An elevated IGF-1 level nearly always indicates that the growth hormone level is elevated. However, IGF-1 increases with growth hormone only up to a growth hormone level of approximately 20 ng/mL, and reaches a plateau thereafter. Therefore, an elevated IGF-1 level does not distinguish between a growth hormone level of 500 or 50.

Of note: nearly everyone has high IGF-1 levels during puberty and pregnancy.

Dynamic tests are cumbersome and unreliable and not usually needed to diagnose acromegaly, but they may be useful after surgery to determine if the tumor is still active. Examples of dynamic tests:

- Oral glucose (75–100 g) suppresses growth hormone levels to less than 2 $\mu\text{g/L}$ in normal subjects, but not in persons with acromegaly.
- Thyrotropin-releasing hormone (TRH) injection has no effect on growth hormone in normal subjects, but increases growth hormone by more than 50% in 60% to 80% of persons with acromegaly.
- Bromocriptine suppresses growth hormone in 50% of persons with acromegaly.

■ Physiology of growth hormone secretion




FIGURE 1

GHRH assays are usually not needed. However, a few persons have acromegaly caused by oversecretion of GHRH by the hypothalamus or from ectopic tumors. In such patients, the histologic study of the excised tissue might show pituitary hyperplasia but not the tumor. Plasma GHRH should be measured in every acromegalic person who has evidence of another, extrapituitary tumor (eg, in a lung, the pancreas, or an adrenal gland).

TABLE 1

Signs and symptoms of acromegaly

General	Cardiopulmonary
Fatigue	Congestive heart failure
Heat intolerance	Arrhythmias, hypertension
Weight gain	Cardiomegaly
	Left ventricular hypertrophy
	Restrictive pulmonary disease
Skin structures	Genitourinary
Acral growth (hands and feet)	Decreased libido
Doughy and oily skin	Erectile dysfunction
Skin tags	Oligomenorrhea
Coarsening of facial features	Infertility
Hypertrichosis	Nephrolithiasis
Hyperhidrosis	
Acanthosis nigricans	Gastrointestinal
Lipomata	Cholelithiasis
Hyperpigmentation	Hepatosplenomegaly
	Colonic polyps, cancer
Head and neck	Neurologic
Headaches	Paresthesias
Goiter	Carpal tunnel syndrome
Frontal bossing	Seizures
Parotid enlargement	
Prognathism	
Eyes	Endocrinologic
Visual field defects	Glucose intolerance
Papilledema	Diabetes mellitus
	Hypolipidemia
Nose, throat, and sinuses	Pituitary apoplexy
Sinus congestion	
Enlarged tongue	Musculoskeletal
Malocclusion	Weakness
Voice change	Proximal myopathy
Sleep apnea	Arthralgias
Rhinorrhea	Osteoporosis
	Osteoarthritis

SOURCE: ADAPTED FROM MAUGANS TA, COATES ML. DIAGNOSIS AND TREATMENT OF ACROMEGALY. AMERICAN FAMILY PHYSICIAN 1995; 52:207-213.

TREATING ACROMEGALY

A preliminary, retrospective study showed that treatment of acromegaly normalizes the mortality rate—but only if the growth hormone level can be reduced to less than 2.5 ng/mL.² Even if this goal cannot be achieved, however, treatment can alleviate the symptoms of this painful and unpleasant disease.

There are three modalities of therapy: surgery, radiotherapy, and medical therapy.

Surgery

Ross and Wilson³ reviewed a series of 214 of their own surgical patients and 1,360 cases reported in the literature. Surgical complications occurred in 7% to 20% of patients, and hypopituitarism occurred in 5% to 18%. Growth hormone levels fell to less than 5 ng/mL in 56% of their own patients, and 60% in the other patients. However, the few studies to date that looked at IGF-1 levels, which we now know is the best parameter to follow, found success rates of only 20% to 30%.

Radiotherapy

Radiotherapy is often used when surgery fails to lower the growth hormone level to the normal range. This therapy takes time to work. In a series of 46 patients, Feek et al⁴ found that the growth hormone level fell by approximately 50% in the first few years after radiotherapy, and then by about 14% per year up to 20 years.

Approximately 50% of patients achieve growth hormone levels less than 5 ng/mL 10 years after starting radiotherapy. Fewer patients achieve a normal IGF-1 level: at our institution, only 2 of 38 patients achieved normal IGF-1 levels with radiotherapy.

Hypopituitarism is the most common complication, occurring in 40% to 50% of patients after 5 years. Other side effects are visual impairment, necrosis of brain tissue, and increased risk of brain malignancy.

Medical therapy

Two classes of drugs are used in acromegaly: dopamine agonists (eg, bromocriptine) and somatostatin analogs (eg, octreotide, lanreotide).

Dopamine agonists. In a review of 31 clinical trials, Jaffe and Barkan⁵ found that the dopamine agonist bromocriptine, in doses of 7.5 to 80 mg/day, lowered growth hormone levels to less than 10 ng/mL in 287 of 543 patients, and to less than 5 ng/mL in 115. Normal IGF-1 levels were achieved in approximately 10%.




Somatostatin analogs are highly effective, reducing growth hormone concentrations to less than 5 ng/mL in more than half of patients and normalizing IGF-1 levels in approximately 70%.⁶ They also work rapidly, often causing soft-tissue regression within 48 hours of starting therapy. Another advantage: they are the only medications that address acromegaly due to ectopic GHRH production.

A disadvantage of currently available preparations is that they must be given by injection three times per day. Long-acting formulations of octreotide and lanreotide are being developed.

A mutant form of growth hormone, which occupies the growth hormone receptor but is physiologically inactive, is also under development.

REFERENCES

1. Thapar K, Kovacs K, Stefaneau L, et al. Overexpression of the growth-hormone-releasing hormone gene in acromegaly-associated pituitary tumors. *Am J Pathol* 1997; 151:769-784.
2. Bates AS, Van't Hoff W, Jones JM, Clayton RN. Does treatment of acromegaly affect life expectancy? *Metab Clin Exper* 1995; 44(suppl 1):1-5.
3. Ross DA, Wilson CB. Results of transsphenoidal microsurgery for growth hormone-secreting pituitary adenoma in a series of 214 patients. *J Neurosurg* 1988; 68:854-867.
4. Feek CM, McLelland J, Seth J, et al. How effective is external pituitary irradiation for growth hormone-secreting pituitary tumors? *Clin Endocrinol* 1984; 20:401-408.
5. Jaffe CA, Barkan AL. Treatment of acromegaly with dopamine agonists. *Endocrinol Metab Clin North Am* 1992; 21:713-735.
6. Lambers SWJ, van der Lely A-J, de Herder WW, Hofland LJ. Octreotide. *N Engl J Med* 1996; 334:246-254.



**Earn Valuable
CME Credit!**

Prepare and Review with Distinguished Cleveland Clinic Faculty Members

THE CLEVELAND CLINIC INTENSIVE REVIEW OF INTERNAL MEDICINE

James F. Stoller, MD, Muzaffar Ahmad, MD
and David L. Longworth, MD

Due June 1998/904 pages/570 illustrations/30087-3/\$89.00

Phone orders accepted 24 hours a day, 7 days a week (US only).
Prices subject to change without notice.

From the US:

Call: 1-800-638-0672

Fax: 1-800-447-8438

From Canada:

Call: 1-800-665-1148

Fax: 1-800-665-0103

Outside the US & Canada:

Call: 410-528-4223

Fax: 410-528-8550

PUT YOUR FINGER ON IT
W&Wonline
www.wilkins.com


Williams & Wilkins
A WAVERLY COMPANY

Printed in US 98 STOLLEAD ► S8B049 A