



FIRAS ZAHR, MD

Department of Pulmonary, Allergy,
and Critical Care Medicine,
Cleveland Clinic

ATUL C. MEHTA, MD

Department of Pulmonary, Allergy, and
Critical Care Medicine, Cleveland Clinic

OMAR A. MINAI, MD

Department of Pulmonary, Allergy, and
Critical Care Medicine, Cleveland Clinic

The Clinical Picture

A young man with unexplained dyspnea




FIGURE 1. Flow-volume loop in a patient with persistent dyspnea.

A 22-YEAR-OLD MAN presents to the pulmonary clinic with persistent dyspnea with exertion, nasal congestion, and epistaxis.

About a year ago, the patient was evaluated for the same symptoms at a community hospital. At the time, computed tomography (CT) of the chest revealed multiple pulmonary nodules with cavitation. Serum was positive for antineutrophil cytoplasmic antibodies with a cytoplasmic staining pattern (c-ANCA). Pulmonary function tests showed a forced vital capacity of 4.1 L (70% of predicted value) and a forced expiratory volume in 1 second (FEV₁) of 2.62 L (54% of predicted).

He underwent sinus surgery with difficult intubation. He was diagnosed with Wegener granulomatosis based on histopathology and was prescribed daily prednisone and cyclophosphamide. The pulmonary nodules resolved but the shortness of breath persisted.

Physical examination. Now, the patient has bilateral sinus tenderness, saddle-nose deformity, and malodorous nasal crusting. His lung fields are clear and his heart sounds are normal. His breathing pattern is normal with no use of accessory muscles or paradoxical breathing.

CT of the chest shows new nonspecific bilateral parenchymal nodules (2–3 mm), but marked improvement compared with the previous CT.


Pulmonary function tests reveal normal flows, volumes, and diffusion capacity. His FEV₁ has improved (3.4 L, 69% of predicted). His flow-volume loop is seen in **FIGURE 1**.

■ DYSPNEA IN WEGENER GRANULOMATOSIS

Q: What is the most likely cause of his persistent shortness of breath?

- ☐ Pulmonary vascular disease complicating Wegener granulomatosis
- ☐ Parenchymal lung disease due to Wegener granulomatosis
- ☐ Fixed upper airway obstruction
- ☐ Neuromuscular weakness

A: Pulmonary vascular disease complicating Wegener's granulomatosis is extremely rare¹ and does not explain the shape of the flow-volume loop or the bronchoscopic findings.



ADAPTED FROM KAVURU MS, WIEDEMANN HP. ASTHMA. IN: GEORGE RB, LIGHT RW, MATTHAY MA, MATTHAY RA, EDITORS. CHEST MEDICINE. ESSENTIALS OF PULMONARY AND CRITICAL CARE MEDICINE. 5TH ED. PHILADELPHIA: LIPPINCOTT WILLIAMS AND WILKINS. 2005:131-162.

Fixed upper airway obstruction is correct. Although the patient's pulmonary function tests are normal, the flow-volume loop (FIGURE 1) shows flattening of the inspiratory and expiratory limbs, indicating that fixed upper airway obstruction is likely.

Upper airway obstruction can be categorized into three types,³ each having characteristic flow-volume loops (**FIGURE 2**).

Fixed upper airway obstruction occurs

CLEVELAND CLINIC JOURNAL OF MEDICINE VOLUME 73 • NUMBER 6 JUNE 2006 555




FIGURE 3. Bronchoscopic image showing the vocal cords (arrows A and C) and severe subglottic stenosis (arrow B).

with tracheal stenosis or goiter. Both limbs of the flow-volume loop are flattened.

■ TRACHEAL STENOSIS

Our patient undergoes flexible bronchoscopy, which demonstrates severe subglottic stenosis (**FIGURE 3**). In this case, the cause is likely to be either Wegener granulomatosis or the previous traumatic intubation.

■ REFERENCES

1. Sanchez O, Garcia G, Hamid A, et al. Pulmonary hypertension complicating systemic diseases other than connective tissue diseases [in French]. *Presse Med* 2003; 32:800–803.
2. Schwarz MI, Brown KK. Small vessel vasculitis of the lung. *Thorax* 2000; 55:502–510.
3. Aboussouan LS, Stoller JK. Diagnosis and management of upper airway obstruction. *Clin Chest Med* 1994; 15:35–53.
4. Aozasa K, Ohsawa M, Tajima K, et al. Nation-wide study of lethal mid-line granuloma in Japan: frequencies of Wegener's granulomatosis, polymorphic reticulosis, malignant lymphoma and other related conditions. *Int J Cancer* 1989; 44:63–66.
5. Langford CA, Sneller MC, Hallahan CW, et al. Clinical features and therapeutic management of subglottic stenosis in patients with Wegener's granulomatosis. *Arthritis Rheum* 1996; 39:1754–1760.
6. Alaani A, Hogg RP, Drake Lee AB. Wegener's granulomatosis and subglottic stenosis: management of the airway. *J Laryngol Otol* 2004; 118:786–790.
7. Kastanos N, Estopa Miro R, Marin Perez A, Xaubet Mir A, Agusti-Vidal A. Laryngotracheal injury due to endotracheal intubation: incidence, evolution, and predisposing factors. A prospective long-term study. *Crit Care Med* 1983; 11:362–367.

ADDRESS: Omar A. Minai, MD, Department of Pulmonary, Allergy, and Critical Care Medicine, G62, Cleveland Clinic, 9500 Euclid Ave, Cleveland OH 44195; e-mail minai@ccf.org.

Wegener granulomatosis causes otolaryngologic problems at some stage of the disease in more than 90% of cases: subglottic stenosis occurs in 10% to 20% due to circumferential scarring and narrowing of the airway.⁴ It is frequently resistant to systemic immunosuppressive therapy, and fibrosis often develops despite aggressive treatment.⁵ Tracheostomy is required in up to 60% of cases. Decannulation is not always successful: restenosis often occurs due to failure of the intervention failure or to continued disease activity.⁶ Intralesional steroids can be safe and effective and may reduce the need for systemic therapy when used with tracheal dilation.⁴ Surgical resection by laser or reconstruction should only be offered to patients who are in remission and have required no immunosuppressive therapy for at least 1 year.

Intubation trauma or excessive cuff pressure during prolonged mechanical ventilation can also cause tracheal stenosis.⁷ These complications are rarely seen now with better intubation techniques and routine monitoring of cuff pressure.

Case revisited. The patient was treated with a steroid injection of the subglottis, dilation of the subglottic stenosis, and administration of mitomycin-C. His dyspnea substantially improved following the procedure. ■