

THE CLINICAL PICTURE

ANNA CAMILLE MORENO, DO, NCMP

Duke Women's Health Associates, Department of OB/GYN, Duke University Medical Center; Assistant Professor, Duke University, Raleigh, NC

OLUWATOSIN GOJE, MD, MSCR, FACOG

Center for Gynecologic Infectious Disease; Director, Ob/Gyn & Women's Health Institute, Cleveland Clinic

MELISSA PECK PILIANG, MD, FAAD

Dermatology and Anatomic Pathology; Vice Chair, Education; Associate Program Director, Dermatology Residency; Associate Program Director, Dermatopathology Fellowship, Cleveland Clinic

PELIN BATUR, MD, NCMP, CCD

Ob/Gyn & Women's Health Institute, Cleveland Clinic; Associate Professor of Medicine, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University, Cleveland, OH; Steering Committee, Women's Preventive Services Initiative, American College of Obstetricians and Gynecologists and US Department of Health and Human Services, Health Resources & Services Administration; Clinical Guideline Committee of the American College of Physicians; Deputy Editor, *Cleveland Clinic Journal of Medicine*

Vulvar and gluteal manifestations of Crohn disease



Figure 1. Symmetric edema and erythema of the vulva.



Figure 2. A linear ulcer with sharply demarcated borders in the gluteal cleft in another patient.

Crohn disease is associated with extraintestinal manifestations in up to 45% of patients

A 37-YEAR-OLD WOMAN presented with recurring painful swelling and erythema of the vulva over the last year. Despite a series of negative vaginal cultures, she was prescribed multiple courses of antifungal and antibacterial treatments, while her symptoms continued to worsen. She had no other relevant medical history except for occasional diarrhea and abdominal cramping, which were attributed to irritable bowel syndrome.

On examination, she had symmetric edema and erythema of the vulva (**Figure 1**). Closer inspection revealed a nonulcerated, slightly friable nodule of approximately 4 mm on her right labium minus. A biopsy of this region demonstrated multiple noncaseating granulomas and mixed inflammatory infiltrates. An

acid-fast stain for mycobacteria was negative. Vulvar skin ultrasonography demonstrated fistulas and increased dermal thickness with altered subcutaneous tissue. She was encouraged to undergo colonoscopy, which showed findings suggestive of Crohn disease.

■ CROHN DISEASE OUTSIDE THE GASTROINTESTINAL TRACT

Crohn disease primarily affects the gastrointestinal tract but is associated with extraintestinal manifestations (in the oral cavity, eyes, skin, and joints) in up to 45% of patients.¹

The most common mucocutaneous manifestations are granulomatous lesions that extend directly from the gastrointestinal tract, including perianal and peristomal skin tags, fistulas, and perineal ulcerations. In most cas-

doi:10.3949/ccjm.86a.19062

es, the onset of cutaneous manifestations follows intestinal disease, but vulvar Crohn disease may precede gastrointestinal symptoms in approximately 25% of patients, with the average age at onset in the mid-30s.¹

The pathogenesis of vulvar Crohn disease remains unclear. One theory involves production of immune complexes from the gastrointestinal tract and a possible T-lymphocyte-mediated type IV hypersensitivity reaction.²

The most commonly reported symptoms include pain, dyspareunia, pruritus, and discharge.¹ The classic findings, found in 50% of cases, include perianal and peristomal skin tags, fistulae, perineal ulcers, linear ulcers (resembling knife cuts), abscesses, and fissures.³ **Figure 2** shows a linear ulcer in the gluteal cleft, with sharply demarcated borders and resembling a knife cut, in another patient treated at our institution. Associated perianal fissures are also seen, in addition to vulvar edema.⁴

The diagnosis of vulvar Crohn disease

should be considered in a patient who has vulvar pain, edema, and ulcerations not otherwise explained, whether or not gastrointestinal Crohn disease is present. The diagnosis is established with clinical history and characteristic histopathology on biopsy. Multiple biopsies may be needed, and early endoscopy is recommended to establish the diagnosis. The histologic features include noncaseating and nonnecrotizing granulomatous dermatitis or vulvitis with occasional reports of eosinophilic infiltrates and necrobiosis.^{5,6} An imaging study such as ultrasonography is sometimes used to differentiate between a specific cutaneous manifestation of Crohn disease and its complications such as perianal fistula or abscess.

Clinical vulvar lesions are nonspecific, and those of Crohn disease are frequently mistaken for infectious, inflammatory, or traumatic vulvitis. Diagnostic biopsy for histologic analysis is warranted.

REFERENCES

1. Andreani SM, Ratnasingham K, Dang HH, Gravante G, Giordano P. Crohn's disease of the vulva. *Int J Surg* 2010; 8(1):2–5. doi:10.1016/j.ijssu.2009.09.012
2. Siroy A, Wasman J. Metastatic Crohn disease: a rare cutaneous entity. *Arch Pathol Lab Med* 2012; 136(3):329–332. doi:10.5858/arpa.2010-0666-RS
3. Foo WC, Papalas JA, Robboy SJ, Selim MA. Vulvar manifestations of Crohn's disease. *Am J Dermatopathol* 2011; 33(6):588–593. doi:10.1097/DAD.0b013e31820a2635
4. Amankwah Y, Haefner H. Vulvar edema. *Dermatol Clin* 2010; 28(4):765–777. doi:10.1016/j.det.2010.08.001
5. Emanuel PO, Phelps RG. Metastatic Crohn's disease: a histopathologic study of 12 cases. *J Cutan Pathol* 2008; 35(5):457–461. doi:10.1111/j.1600-0560.2007.00849.x
6. Hackzell-Bradley M, Hedblad MA, Stephansson EA. Metastatic Crohn's disease: report of 3 cases with special reference to histopathologic findings. *Arch Dermatol* 1996; 132(8):928–932. doi:10.1001/archderm.132.8.928

ADDRESS: Pelin Batur, MD, Women's Health Institute, A8-406, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195; baturp@ccf.org