

Hemodynamic effects of narcotics

Theodore H. Stanley, M.D.

Salt Lake City, Utah

In intact animals and nonaddicted supine patients, morphine and fentanyl and some new, as yet unreleased narcotics, e.g., sulfentanil, produce minimal changes in cardiovascular dynamics.¹⁻³ Indeed, Lowenstein et al¹ observed that the introduction of large (anesthetic) doses of morphine (as much as 1 mg/kg) intravenously as the sole or principal anesthetic did not cause significant circulatory changes in patients without cardiac disease. In patients with aortic valvular disease, stroke volume and cardiac output increased. Yet several investigators observed depression of myocardial contractility by narcotics when isolated myocardial muscle and heart-lung preparations were perfused with saline solution.⁴ When blood was substituted as the perfusing solution, no depressant effect was observed with as much as 30 mg/kg morphine in dog and cat heart-lung preparations.⁵ Apparently, fentanyl also does not change myocardial mechanics in blood-perfused preparations, whereas meperidine does cause considerable myocardial depression.⁶

Myocardial depression does not occur after morphine administration in intact animals or man. Schmidt and Livingston⁷ found no evidence of cardiac depression in awake dogs given as much as 100 mg/kg morphine. Vasco et al⁸ found that morphine had a positive inotropic effect in dogs, which

was dependent on endogenous catecholamine release and blocked by beta-adrenergic blocking agents or previous surgical adrenalectomy. Morphine also increases the concentrations of catecholamines in both blood and urine of human subjects.⁹⁻¹¹ Endogenous catecholamine secretion appears to be dependent on the functional state of the cardiovascular system and plasma concentrations of morphine.^{9, 10} Liu et al¹² have shown that elevated plasma and urinary concentrations of epinephrine and norepinephrine also occur after administration of large doses (0.5 to 30 mg/kg) of fentanyl to dogs. However, recently we found that anesthetic doses of fentanyl do not alter plasma catecholamine concentrations in man.¹³

The potential for cardiovascular stability and anesthesia with large doses of morphine has made it a popular drug in modern anesthesia, especially in critically ill patients with little cardiovascular reserve.^{14, 15} However, morphine may not always be benign with respect to circulatory dynamics. Bradycardia, tachycardia, hypotension, and hypertension have been observed in both man and experimental animals after administration of morphine.¹⁵⁻¹⁹ Therefore, certain precautions must be taken during the induction and maintenance of anesthesia with morphine.

Hypotension can occur during and after administration of even relatively small doses of morphine.¹⁷ Due to arterial and venous dilation and increased vagal activity with cardiac slowing, hypotension most often occurs in hypovolemic, hypertensive patients and in those with high vagal tone or during rapid administration of morphine. Hypotension is less common (a) when morphine is given slowly (5 mg/min or slower), (b) when a rapid infusion of intravenous fluids is given concurrently,

(c) in patients who are in a slight Trendelenburg (head down) position, and (d) in patients with congestive heart failure. The latter findings suggest that a large component of hypotension observed after morphine is given is secondary to venodilation. Experiments in animals⁷ and man^{14, 17, 20} have confirmed the arterial and venodilating ability of morphine, with the latter property being more dominant. Ward et al²¹ have shown that the differential arterial and venous actions of morphine may be due to a selective venous alpha adrenergic blocking effect. However, Zelis et al^{22, 23} and Flaim et al²⁴ found that morphine selectively impairs certain sympathetic reflexes involving peripheral veins; they concluded that this was caused by the action of the drug on the central nervous system. Apparently venoconstriction to tilting or inhalation of carbon dioxide is impaired by morphine, whereas venoconstriction to a single deep breath is not. Whatever the mechanism, we found that venodilation after morphine is dose-related and results in occasional hypotension during induction and increases in blood requirements during and after surgery when compared to patients anesthetized with halothane.²⁵ There is evidence that venodilation and increased blood requirements do not occur with lower doses of morphine (<0.5 mg/kg plus N₂O). (Unpublished data.)

Arterial vasodilation after morphine, in contrast to venodilation, lasts for only a short period of time (15 to 30 minutes)²⁶ and is related to histamine release²⁷ and the direct effects of morphine on vascular smooth muscle.⁷ Histamine release following morphine varies, although dose and rate of morphine administration undoubtedly have some influence. Hypotension secondary to histamine release can be corrected by vasopressors; however, administration

of antihistamines either before or after morphine-induced hypotension is of no benefit.

Hypotension also occurs after the intravenous administration of meperidine (Demerol).⁷ Hypotension occurs more frequently after meperidine and is more profound than after comparable doses of morphine. This is due to a significant negative inotropic effect of meperidine^{7, 28, 29} as well as a marked reduction in systemic vascular resistance. As a result, meperidine has had little value as a "complete" anesthetic, although it is still popular as a supplement in nitrous oxide-narcotic "balanced" anesthesia.³⁰ Meperidine, in contrast to morphine, rarely results in bradycardia but can cause tachycardia.³¹ This may be related to the structural similarity of meperidine to atropine.

Alphaprodine (Nisentil) is structurally similar to meperidine and has had limited use as an analgesic supplement in patients in the operating room and delivery suite. Unfortunately, although alphaprodine is somewhat shorter acting than meperidine, it possesses all of the cardiovascular depressant qualities of the latter.³² Large anesthetic doses (50 to 100 $\mu\text{g}/\text{kg}$) in man and small analgesic doses (5 to 10 $\mu\text{g}/\text{kg}$) of fentanyl also reduce arterial blood pressure but rarely below 90 torr systolic.^{2, 33} Hypotension, when it occurs after fentanyl administration, is primarily due to bradycardia, and can be reversed with atropine, ephedrine, or even large doses of pancuronium.³⁴

Hypertension, occasionally severe enough to necessitate the use of vasodilators, has been found with narcotic-oxygen anesthesia or narcotic-nitrous oxide balanced anesthesia.^{15, 16, 35, 36} This complication is a frequent and especially serious problem in patients with coronary artery disease.¹⁸ However, it

most often occurs in healthy patients or those with only minimal impairment of cardiovascular reserve and is much less common in patients with valvular heart disease or in those who are severely ill irrespective of the primary pathology.³⁷ Hypertension during and after coronary artery surgery may be related to operative manipulation of the coronary arteries or to the type of patient who acquires coronary artery disease rather than to morphine or the other opioids.¹⁸ Perhaps the primary cause of hypertension during opioid anesthesia may be inadequate analgesia or narcosis or both.^{16, 38} This belief is supported by occasional awareness when opioids are the major component of the anesthetic technique.¹⁶ Also hypertension rarely occurs before endotracheal tube intubation or surgical stimulation.³⁶ Both sympathetic nervous stimulation and renin-angiotensin responses are probably not blocked by morphine in doses of 1 to 3 mg/kg .^{10, 39} However, larger doses of morphine, 4 mg/kg or higher, block increases in circulating catecholamine levels in response to surgical stimulus.⁹ Hypertension produced during narcotic anesthesia is often difficult to treat. Higher doses of morphine, other narcotics, and even potent inhalation agents may not be effective in restoring blood pressure when noxious stimulation has elevated it. In these situations, vasodilators or adrenergic blockers may be required to treat hypertension. In addition, increased doses of morphine, a variety of intravenous hypnotics, or low concentrations of inhalation anesthetics may also be needed to suppress awareness.¹⁶

As mentioned, bradycardia often occurs following the intravenous administration of morphine, fentanyl, and some of the other narcotics.^{2, 40, 41} Although bradycardia almost always accompanies

induction of anesthesia with narcotics during breathing of oxygen, it occurs less frequently when nitrous oxide is also used for induction.^{2, 40, 42} This may be due to a possible stimulating effect of nitrous oxide on the sympathetic nervous system. Although premedication with the belladonna drugs or glycopyrrolate minimizes bradycardia after morphine and fentanyl have been given intravenously, these drugs do not always eliminate bradycardia.⁴² Also, treatment of narcotic-induced bradycardia with atropine (0.4 to 0.8 mg, intravenously) is usually, but not always effective. On occasion even large doses of atropine (1 to 2 mg, intravenously) are ineffective in treating narcotic-induced bradycardia. In these situations ephedrine (15 to 25 mg, intravenously) is almost always effective.

Anesthesiologists experienced with narcotic anesthesia are usually not concerned with bradycardia accompanying induction of anesthesia as long as arterial blood pressure or cardiac output or both remain unchanged or are only moderately reduced. This is especially true when patients are brought to the operating room with low heart rates secondary to chronic use of propranolol. Thus, narcotic-induced bradycardia is not usually detrimental to myocardial function and indeed may be advantageous in patients with ischemic cardiac disease. Furthermore, bradycardia from narcotics is relatively transient, usually lasting only 10 to 20 minutes. Most experimental evidence indicates that bradycardia following the use of morphine is caused by stimulation of the vagal nucleus in the medulla of the brain.^{41, 43} However, morphine may also have a direct effect on the sinoatrial node⁴⁴ and depress atrioventricular conduction.⁴⁵ Although narcotics usually do not cause arrhythmias, large doses of

morphine and probably other narcotics used as anesthetics can cause increasing sinoatrial and atrioventricular block and decrease the duration of the refractory phase of atrial depolarization.⁴⁵ Theoretically, this could lead to the development of reentry type arrhythmias, but this rarely occurs. In a recent study it was found that the doses of epinephrine necessary to induce arrhythmias in dogs are not any greater during narcotic-nitrous oxide anesthesia than during halothane, enflurane, or nitrous oxide anesthesia. However, these authors reported that the incidence of malignant arrhythmias (ventricular tachycardia and ventricular fibrillation) was much less with narcotics plus nitrous oxide than with most potent inhalation anesthetics.⁴⁶ Despite these data many clinicians are of the opinion that patients anesthetized with narcotics are less susceptible to catecholamine-induced arrhythmias than patients anesthetized with halogenated compounds, and that the absence of dangerous arrhythmias is the strongest reason for considering narcotic anesthesia in patients with cardiac problems.

The most common arrhythmia noted during narcotic anesthetic administration other than bradycardia, is supraventricular tachycardia.³⁶ This occurs more often during or immediately after endotracheal intubation or surgical stimulation and, therefore, is probably evidence of inadequate anesthesia rather than a direct effect of the narcotic. However, there are certain patients who experience supraventricular tachycardias during narcotic anesthetic induction without concurrent noxious stimulation. This is especially true following the use of large doses of meperidine and alpha-prodine but also occurs with the use of morphine. Although tachycardia following the administra-

tion of meperidine and alpha-prodine may be attributed to the structural similarity of these compounds to atropine, tachycardia following morphine cannot. Tachycardia, when it occurs during induction with morphine, is usually accompanied by facial and upper torso flushing. This suggests that increased circulation concentrations of histamine or catecholamines or both may be the explanation for this infrequent but certainly not uncommon finding.

The pulmonary vascular effects of narcotic administration have not been nearly as well studied as have the systemic vascular actions. However, there are data that demonstrate that morphine significantly increases pulmonary artery blood pressure and decreases pulmonary blood volume.^{40, 47} Hug (unpublished data) has found that large doses of fentanyl intravenously can also occasionally dramatically increase pulmonary artery pressure in man. The mechanism(s) involved in producing these changes have not been elucidated.

As noted, narcotics are usually used along with other drugs (supplements) during anesthesia. The most common supplement used with intravenous narcotics is nitrous oxide. Although nitrous oxide alone has minimal effects on myocardial mechanics⁴⁸ and cardiovascular dynamics,⁴⁹ when given to patients receiving intravenous narcotics, myocardial depression frequently occurs.^{14, 50, 51} McDermott and Stanley⁵⁰ found that after administration of morphine (2 mg/kg), nitrous oxide produces concentration-dependent decreases in stroke volume, cardiac output, and arterial blood pressure and increases in systemic vascular resistances. Impairment of cardiac output was always greater than the reduction in blood pressure because of an increase in peripheral vascular resistance. Heart rate was usually little

changed by addition of nitrous oxide. This could lead to a reduction in total blood flow before the variables that are usually monitored (arterial blood pressure and heart rate) are changed enough to alert the anesthesiologist to the impending danger. Similar decreases in stroke volume and cardiac output and increases in peripheral vascular resistances have resulted from lower doses of morphine,³⁶ other narcotics plus nitrous oxide,^{30, 33, 52} and also with Innovar-nitrous oxide anesthesia. Surprisingly, cardiovascular depression during nitrous oxide-narcotic anesthesia appears not to be related to the plasma concentration of the narcotic.⁵²

A variety of other intravenous supplements are often combined with narcotics during anesthesia to increase analgesia, ensure adequate hypnosis and amnesia, and correct undesirable cardiovascular alterations (usually hypertension). This therapy is effective, but it is usually achieved at the expense of significant cardiovascular depression. Stoelting⁵³ has demonstrated, for example, that when a barbiturate has been given for induction of anesthesia, thiamylal (4 mg/kg intravenously), morphine (1 mg/kg intravenously), reduced cardiac output and stroke volume 30% to 40% and decreased blood pressure 16%; systemic vascular resistance was increased 35%. Similarly, diazepam (5 to 10 mg intravenously), a compound with little cardiovascular effects when employed alone, results in marked cardiovascular depression when added after morphine, fentanyl, and probably most narcotics.^{2, 54} Of the intravenous compounds that have been studied it appears that only scopolamine and droperidol do *not* produce significant cardiovascular depression when combined with intravenously administered narcotics.^{35, 54}

Morphine, meperidine, and more re-

cently fentanyl have also been combined with potent anesthetics.^{55, 56} These techniques have varied from the addition of narcotics to patients who were primarily anesthetized with potent inhalation anesthetics⁵⁵ and vice versa.⁵⁶ These procedures have frequently included numerous other intravenous supplements as well. The rationale for this practice is based on the minimal effects of narcotics on cardiovascular dynamics, their presumed ability to decrease the MAC of the inhalation anesthetics and the belief that use of these combinations produces deep anesthesia without marked cardiovascular depression.^{1, 55, 57} Although all of the above seem reasonable, there are, unfortunately, little data that these combinations have any more benign effect on cardiovascular dynamics than the use of any of the compounds alone or that there are not more potential problems from drug interactions than advantages. Stoelting et al⁵⁶ have shown that addition of low to moderate concentrations of halothane after large doses of morphine produces severe cardiovascular depression. Bennett and Stanley⁵⁵ have found that the same changes occur when fentanyl (200 µg intravenously) is added during enflurane-nitrous oxide anesthesia, but that lower doses of fentanyl (50 to 100 µg) produced little change in cardiovascular dynamics during enflurane-nitrous oxide anesthesia.

Narcotic compounds decrease cerebral blood flow and intracranial pressure.^{58, 59} For this reason nitrous oxide-narcotic anesthesia with or without other supplements is a popular technique for patients with increases in intracranial pressure. Some narcotics can decrease the consumption of cerebral oxygen, but the reduction is transient and usually less than the corresponding reduction in cerebral blood flow, at least

in dogs.^{58, 60, 61} Although there is no evidence that narcotic-induced decreases in cerebral flow are dangerous in man, some anesthesiologists are reluctant to use narcotic techniques in patients with marginal cerebral circulation.

References

1. Lowenstein E, Hallowell P, Levine FH, Daggett WM, Austen WG, Laver MB. Cardiovascular response to large doses of intravenous morphine in man. *N Engl J Med* 1969; **281**: 1389-93.
2. Stanley TH, Webster LR. Anesthetic requirements and cardiovascular effects of fentanyl-oxygen and fentanyl-diazepam-oxygen anesthesia in man. *Anesth Analg* 1978; **57**: 411-6.
3. Van de Walle J, Lauwers P, Adriaensen H. Double blind comparison of fentanyl and sulfentanil in anesthesia. *Acta Anaesth Belg* 1976; **27**: 129-38.
4. Rothschild F, Roth L. *Lipoidwirkungen am isolierten Forschherz und isolierten Dünndarm. Einfluss von lipoiden auf Giftwirkungen am isolierten Organ.* *Arch Int Pharmacodyn* 1937; **57**: 1-6.
5. Falcke W. Some actions of morphine on the circulation. *Fed Proc* 1965; **24**: 163.
6. Ostheimer GW, Shanahan EA, Guyton RA, Daggett WM, Lowenstein E. Effects of fentanyl and droperidol on canine left ventricular performance. *Anesthesiology* 1975; **42**: 288-91.
7. Schmidt CF, Livingston AE. The action of morphine on the mammalian circulation. *J Pharmacol Exp Ther* 1933; **47**: 411-41.
8. Vasco JS, Henney RP, Brawley RK, Oldham HN, Morrow AG. Effects of morphine on ventricular function and myocardial contractile force. *Am J Physiol* 1966; **210**: 329-34.
9. Stanley TH, Isern-Amaral J, Lathrop GD. The effects of morphine and halothane anesthesia on urine norepinephrine during and after coronary artery surgery. *Can Anaesth Soc J* 1975; **22**: 478-85.
10. Stanley TH, Isern-Amaral J, Lathrop GD. Urine norepinephrine excretion in patients undergoing mitral or aortic valve replacement with morphine anesthesia. *Anesth Analg* 1975; **54**: 509-17.
11. Balasoroswathi K, Glisson SN, El-Etr A, et al. Serum epinephrine and norepinephrine during valve replacement and aorta-coronary bypass. *Can Anaesth Soc J* 1978; **25**: 198-203.

12. Liu WS, Bidwai AV, Lunn JK, Stanley TH. Urine catecholamine excretion after large doses of fentanyl, fentanyl and diazepam and fentanyl, diazepam and pancuronium. *Can Anaesth Soc J* 1977; **24**: 371-9.
13. Stanley TH, Berman L, Green O, et al. Fentanyl-oxygen anesthesia for coronary artery surgery; plasma catecholamine and cortisol responses. *Anesthesiology Abstracts*. In press.
14. Stoelting RK, Gibbs PS. Hemodynamic effects of morphine and morphine-nitrous oxide in valvular heart disease and coronary-artery disease. *Anesthesiology* 1973; **38**: 45-52.
15. Arens JF, Benbow BP, Ochsner JL, et al. Morphine anesthesia for aortocoronary bypass procedures. *Anesth Analg* 1972; **51**: 901-9.
16. Lowenstein E. Morphine "anesthesia"; a perspective. *Anesthesiology* 1971; **35**: 563-5.
17. Drew JH, Dripps RD, Comroe JH Jr. Clinical studies on morphine. II. The effect of morphine upon the circulation of man and upon the circulatory and respiratory responses to tilting. *Anesthesiology* 1946; **7**: 44-61.
18. Estafanous FG, Tarazi RC, Viljoen JF, El Tawil MY. Systemic hypertension following myocardial revascularization. *Am Heart J* 1973; **85**: 732-8.
19. Henney RP, Vasco JS, Brawley RK, Oldham HN, Morrow AG. The effects of morphine on the resistance and capacitance vessels of the peripheral circulation. *Am Heart J* 1966; **72**: 242-50.
20. Stanley TH, Gray NH, Stanford W, Armstrong A. The effects of high-dose morphine on fluid and blood requirements in open-heart operations. *Anesthesiology* 1973; **38**: 536-41.
21. Ward JM, McGrath RL, Weil JV. Effects of morphine on the peripheral vascular response to sympathetic stimulation. *Am J Cardiol* 1972; **29**: 659-66.
22. Zelis R, Mansour EJ, Capone RJ, Mason DT. The cardiovascular effects of morphine; the peripheral capacitance and resistance vessels in human subjects. *J Clin Invest* 1974; **54**: 1247-58.
23. Zelis R, Flaim SF, Eisele JH. Effects of morphine on reflex arteriolar constriction induced in man by hypercapnia. *Clin Pharmacol Ther* 1977; **22**: 172-8.
24. Flaim SF, Vismara LA, Zelis R. The effects of morphine on isolated cutaneous canine vascular smooth muscle. *Res Commun Chem Pathol Pharmacol* 1977; **16**: 191-4.
25. Stanley TH, Gray NH, Isern-Amaral JH, Patton C. Comparison of blood requirements during morphine and halothane anesthesia for open-heart surgery. *Anesthesiology* 1974; **41**: 34-8.
26. Wong KC, Martin WE, Hornbein TF, Freund FG, Everett J. The cardiovascular effects of morphine sulfate with oxygen and with nitrous oxide in man. *Anesthesiology* 1973; **38**: 542-9.
27. Thompson WL, Walton RP. Elevation of plasma histamine levels in the dog following administration of muscle relaxants, opiates and macromolecular polymers. *J Pharmacol Exp Ther* 1964; **143**: 131-6.
28. Strauer BE. Contractile responses to morphine piritramide, meperidine, and fentanyl; a comparative study of effects on the isolated ventricular myocardium. *Anesthesiology* 1972; **37**: 304-10.
29. Rees HA, Muir AL, McDonald HR, Lawrie DM, Burton JL, Donald KW. Circulatory effects of pethidine in patients with acute myocardial infarction. *Lancet* 1967; **2**: 863-6.
30. Stanley TH, Liu WS. Cardiovascular effects of meperidine-N₂O anesthesia before and after pancuronium. *Anesth Analg* 1977; **56**: 669-73.
31. King BD, Elder JD Jr, Dripps RD. The effect of the intravenous administration of meperidine upon the circulation of man and upon the circulatory response to tilt. *Surg Gynecol Obstet* 1952; **94**: 591-7.
32. Reddy P, Liu WS, Stanley TH, Johansen R. Hemodynamic effects of anesthetic doses of alpha-prodine and sufentanil in dogs. (Abstr) *Anesthesiology* 1979; **51**: S102.
33. Stoelting RK, Gibbs PS, Creasser CW, Peterson C. Hemodynamic and ventilatory responses to fentanyl, fentanyl-droperidol, and nitrous oxide in patients with acquired valvular heart disease. *Anesthesiology* 1975; **42**: 319-24.
34. Liu WS, Bidwai AV, Stanley TH, et al. The cardiovascular effects of diazepam and of diazepam and pancuronium during fentanyl and oxygen anesthesia. *Can Anaesth Soc J* 1976; **23**: 395-403.
35. Bennett GM, Loeser EA, Stanley TH. Cardiovascular effects of scopolamine during morphine-oxygen and morphine-nitrous oxide-oxygen anesthesia in man. *Anesthesiology* 1977; **46**: 225-7.
36. Bennett GM, Stanley TH. Comparison of the cardiovascular effects of morphine-N₂O and fentanyl-N₂O balanced anesthesia before and after pancuronium. (Abstr) *Anesthesiology* 1979; **51**: S138.

37. Estafanous FG, Tarazi RC, Buckley S, Taylor PC. Arterial hypertension in immediate post-operative period after valve replacement. *Br Heart J* 1970; **40**: 718-24.
38. Hug CC. *Pharmacology-Anesthetic Drugs*. Kaplan JA, ed. New York: Grune & Stratton, 1979: 19-24.
39. Bailey DR, Miller ED Jr, Kaplan JA, Rogers PW. The renin-angiotensin-aldosterone system during cardiac surgery with morphine-nitrous oxide anesthesia. *Anesthesiology* 1975; **42**: 538-44.
40. Lappas DG, Geha D, Fischer JE, Laver MB, Lowenstein E. Filling pressures of the heart and pulmonary circulation of the patient with coronary-artery disease after large intravenous doses of morphine. *Anesthesiology* 1975; **42**: 153-9.
41. Cohn AE. The effect of morphine on the mechanism of the dog's heart after removal of one vagus nerve. *Proc Soc Exp Biol Med* 1913; **10**: 93-7.
42. Liu WS, Bidwai AV, Stanley TH, et al. Cardiovascular dynamics after large doses of fentanyl and fentanyl plus N₂O in the dog. *Anesth Analg* 1972; **55**: 168-72.
43. Robbins BH, Fitzhugh OG, Baxter JH Jr. The action of morphine in slowing the pulse. *J Pharmacol Exp Ther* 1939; **66**: 216-23.
44. Kennedy BL, West TC. Effect of morphine on electrically-induced release of autonomic mediators in the rabbit sinoatrial node. *J Pharmacol Exp Ther* 1967; **157**: 149-58.
45. Eyster JAE, Meek WJ. Cardiac irregularities in morphine poisoning in the dog. *Heart* 1952; **4**: 542.
46. Puerto BA, Wong KC, Puerto AX, Blatnick RA. Epinephrine-induced dysrhythmias; comparison during anaesthesia with narcotics and with halogenated inhalation agents in dogs. *Can Anaesth Soc J* 1979; **26**: 263-8.
47. Roy SB, Singh I, Bhatia MH, Khanna PK. Effect of morphine on pulmonary blood volume in convalescents from high altitude pulmonary oedema. *Br Heart J* 1965; **27**: 876-8.
48. Goldberg AH, Sohn YZ, Phear WPC. Direct myocardial effects of nitrous oxide. *Anesthesiology* 1975; **43**: 61-9.
49. Craythorne NWB, Darby TD. The cardiovascular effects of nitrous oxide in the dog. *Br J Anaesth* 1965; **37**: 560-5.
50. McDermott RW, Stanley TH. The cardiovascular effects of low concentrations of nitrous oxide during anesthesia. *Anesthesiology* 1974; **41**: 89-91.
51. Lappas DG, Buckley MJ, Laver MB, Daggett WM, Lowenstein E. Left ventricular performance and pulmonary circulation following addition of nitrous oxide to morphine during coronary-artery surgery. *Anesthesiology* 1975; **43**: 61-9.
52. Lunn JK, Stanley TH, Eisele J, et al. High dose fentanyl anesthesia for coronary artery surgery; plasma fentanyl concentrations and influence of N₂O on cardiovascular responses. *Anesth Analg*. In press.
53. Stoelting RK. Influence of barbiturate anesthetic induction on circulatory responses to morphine. *Anesth Analg* 1977; **56**: 615-7.
54. Stanley TH, Bennett GM, Loeser EA, Kawamura R, Sentker CR. Cardiovascular effects of diazepam and droperidol during morphine anesthesia. *Anesthesiology* 1976; **44**: 255-8.
55. Bennett GM, Stanley TH. Cardiovascular effects of fentanyl during enflurane anesthesia in man. *Anesth Analg* 1979; **58**: 179-82.
56. Stoelting RK, Creasser CW, Gibbs PS, et al. Circulatory effects of halothane added to morphine anesthesia in patients with coronary artery disease. *Anesth Analg* 1974; **53**: 449-55.
57. Eger EI II. *Anesthetic Uptake and Action*. Baltimore: Williams and Wilkins, 1974: 14.
58. Michenfelder JD, Theye RA. Effects of fentanyl, droperidol, and Innovar on canine cerebral metabolism and blood flow. *Br J Anaesth* 1971; **43**: 630-6.
59. Larson CP Jr, Mazze RI, Cooperman LH, Wollman H. Effects of anesthetics on cerebral, renal, and splanchnic circulations; recent developments. *Anesthesiology* 1974; **41**: 169-81.
60. Takeshita H, Michenfelder JD, Theye RA. The effects of morphine and N-allylnormorphine on canine cerebral metabolism and circulation. *Anesthesiology* 1972; **37**: 605-12.
61. Miller JD, Barker J. The effect of neuroleptanalgesic drugs on cerebral blood flow and metabolism. (Abstr) *Br J Anaesth* 1969; **41**: 554-5.