



Early exercise testing following percutaneous transluminal coronary angioplasty

DONNA H. KORZICK, MS; DONALD A. UNDERWOOD, MD; CONRAD C. SIMPFENDORFER, MD

■ The value of early symptom-limited stress electrocardiography following percutaneous transluminal coronary angioplasty in assessing late outcome was evaluated in 218 patients. All subjects were tested using the Bruce or Sheffield Protocols, 2.5 ± 1.3 days after percutaneous transluminal coronary angioplasty. Repeat coronary angiography was performed after percutaneous transluminal coronary angioplasty because of symptoms (58%) or as routine follow-up (42%). Stress electrocardiography results were compared to coronary angiography. The sensitivity and specificity were 35.3% and 52.6%, respectively. The positive and negative predictive values were 39.6% and 48.0%. Two acute myocardial infarctions and one coronary angiographic-proven restenosis occurred within hours of the stress electrocardiogram in three patients (1.4%). It is concluded that symptom-limited stress testing immediately following percutaneous transluminal coronary angioplasty has no prognostic value and may carry increased risk for immediate negative coronary events.

□ INDEX TERMS: ANGIOPLASTY; EXERCISE TEST □ CLEVE CLIN J MED 1990; 57:53-56

GRADED exercise testing is performed soon after percutaneous transluminal coronary angioplasty for a variety of reasons. These include confirmation of resolution of symptoms, establishment of a new functional baseline, and reassurance of the patient or spouse.¹⁻¹⁰ While numerous exercise studies have demonstrated the positive effects of percutaneous transluminal coronary angioplasty on work capacity, few have looked at the diagnostic accuracy of early stress-electrocardiography for assessing late success or failure. Failure of investigators to define such terms as "soon after" or "early," or to include follow-up catheterization data, have made interpretation

of early results and selection of an optimal testing protocol difficult. In addition, reports of complications¹¹⁻¹⁴ accompanying early stress testing after percutaneous transluminal coronary angioplasty raise questions about the safety of such a procedure during this time period. The purpose of this investigation was to assess the diagnostic value of early symptom-limited stress electrocardiography following percutaneous transluminal coronary angioplasty and to document any associated negative events.

MATERIALS AND METHODS

Patient population

Follow-up data were available in 437 patients who underwent symptom-limited stress testing at The Cleveland Clinic Foundation following percutaneous transluminal coronary angioplasty between January 1984 and January 1985. Of these, 219 patients were eliminated

From the Department of Cardiology, The Cleveland Clinic Foundation.

Address reprint requests to D.A.U., Department of Cardiology, The Cleveland Clinic Foundation, One Clinic Center, 9500 Euclid Avenue, Cleveland, Ohio 44195.

from the analysis because of one of the following reasons: lack of follow-up catheterization data (51%), submaximal heart rate response (26%), or an abnormal resting electrocardiogram (23%) making stress response impossible to assess. The study population represented 100% of the eligible patients who underwent early symptom-limited stress electrocardiography following percutaneous transluminal coronary angioplasty at our institution during the analysis period. All stress tests were performed prior to hospital discharge.

Stress testing protocol

All patients were tested utilizing Bruce or Sheffield protocols.¹⁵ The exercise test was terminated because of fatigue, hypotension, severe ST depression (≥ 3 mm), chest pain (grade 3+), claudication, dyspnea, or arrhythmia. Modified 12-lead electrocardiograms were recorded in all subjects at rest, every three minutes during exercise, during peak exercise, and every two minutes during recovery using a Mason Likar lead system. An abnormal stress-electrocardiographic result was defined as ≥ 1 mm of horizontal ST depression 0.08 seconds after the J point. The stress test was considered to be "early" if performed within one week of percutaneous transluminal coronary angioplasty. All tests were interpreted by the same trained observer to avoid interobserver variance.

Follow-up catheterization

Follow-up catheterization data were available for all patients included in the study population. For the purposes of this investigation, recurrence was defined as a $\geq 50\%$ luminal diameter stenosis in an artery of previously successful percutaneous transluminal coronary angioplasty. A successful result was defined as a $\leq 50\%$ diameter stenosis.

RESULTS

The study sample consisted of 172 men and 46 women. The mean age of the patients was 55.2 ± 8.6 years (range, 36–76 years). Prior to percutaneous transluminal coronary angioplasty, 101 patients were classified as having single-vessel coronary artery disease, with the remaining 117 patients possessing two or more stenotic vessels. A total of 315 vessels were successfully dilated: 58% because of left anterior descending disease, 26% because of disease in the right coronary artery, and 16% because of circumflex disease. Following percutaneous transluminal coronary angioplasty, stenoses were reduced to a mean value of $17.8\% \pm 14.3\%$.

The exercise test was performed 2.5 ± 1.3 days after

percutaneous transluminal coronary angioplasty (range, 1–6 days). The Bruce protocol was selected for 213 patients and the Sheffield protocol was chosen for the remaining five patients. All patients exercised to at least 80% of their maximum-predicted heart rate, with a mean value of $92.9\% \pm 11.0\%$. Ninety-one tests were considered positive and 127 were considered normal.

Within hours of the symptom-limited stress tests, acute myocardial infarction occurred in two patients and catheterization-proven restenosis in another. These patients had been considered stable and had not undergone percutaneous transluminal coronary angioplasty for acute myocardial infarction intervention prior to the early stress test. Acute occlusion had occurred in one of the patients early in the percutaneous transluminal coronary angioplasty procedure while still in the catheterization laboratory.

Repeat catheterization was performed in all 218 patients one week to 12 months after PTCA. Reasons for follow-up angiography included symptoms ($n=127$) and routine follow-up ($n=91$). Recurrence was documented in 46% of the patients as defined by previously mentioned criteria.

Comparing follow-up angiography with stress-electrocardiographic data, a false-negative result was observed in 66 patients, a false-positive result in 55 patients, a true-negative result in 61 patients, and a true-positive result in 36 patients. Sensitivity and specificity were 35.3% and 52.6%, respectively, with a positive and negative predictive accuracy of 39.6% and 48.0%, respectively. The degree of coronary artery disease (single *v* multi-vessel) did not affect the outcome nor did the time interval between initial PTCA and follow-up catheterization.

DISCUSSION

In the presence of established coronary artery disease, graded exercise testing has been identified as a valuable noninvasive tool for assessing prognosis and stratifying risk.^{16–19} Early after myocardial infarction, the presence of ≥ 1 mm exercise-induced ST depression has been associated with a three-to-20-fold increase in risk for subsequent cardiac events in comparison to those without ST abnormality.^{19–21}

Our data show that an abnormal stress electrocardiogram within one week of percutaneous transluminal coronary angioplasty was not useful in predicting future re-stenosis. Of the total number of negative exercise tests, 52% were considered false-negative when compared to follow-up angiography. Moreover, a false-posi-

tive result occurred in 60% of the stress tests interpreted as normal.

In their initial report on percutaneous transluminal coronary angioplasty, Gruntzig et al²² recommended the use of *submaximal* bicycle ergometric evaluation two days after dilation to evaluate the immediate success of the angioplasty. Some, however, have adopted the use of *symptom-limited* exercise testing during this period.^{6,8} It is clear from the literature (Table 1) that testing protocols vary considerably and lack standardization. The time frame for "early" testing ranges from two days to eight months, with no clear consensus definition. A recent American College of Cardiology/American Heart Association task force¹⁶ concluded that exercise testing following percutaneous transluminal coronary angioplasty may be performed safely prior to hospital discharge but did not specify a maximal or submaximal protocol.

There have been four reports of complications accompanying early symptom-limited exercise testing following percutaneous transluminal coronary angioplasty: exercise-induced vasospasm,¹³ acute thrombus formation,¹² and two reocclusions.^{11,14} In our series, acute myocardial infarction was documented in two patients and one acute occlusion in another patient. The overall incidence of acute occlusion following successful percutaneous transluminal coronary angioplasty at our institution has been reported at 2%.²³ Of these, 84% occurred within six hours of the percutaneous transluminal coronary angioplasty. The only late occlusion (>24 hours) occurred following a symptom-limited stress test. The complication rate of 1.4% (three events per 218 patients) exceeds the generally accepted risk in stress testing of three cardiac events per 10,000 patients.¹⁵

While the mechanism of re-stenosis remains to be established,²⁴ it is clear that the process generally occurs during the first six months following angioplasty.²⁵ If exercise testing is to be used effectively as a predictor of future cardiac events, the timing of the exercise test is crucial. Wijns et al³ reported that myocardial thallium perfusion scintigraphy four weeks after angioplasty was highly predictive of re-stenosis, with a positive predictive value of 82% and no reported complications. However, because the stress electrocardiogram and thallium scintigraphic responses may revert from positive to negative in a delayed fashion during the recovery period, Scholl et al¹ have suggested that follow-up exercise testing be conducted at three months postangioplasty. In support of this, Manyari and colleagues²⁶ compared thallium stress scintigraphy at nine days and at three, six, and nine months postangioplasty in patients without evidence of recurrence on follow-up catheterization. Al-

TABLE 1
EXERCISE TESTING PROTOCOLS FOLLOWING PERCUTANEOUS TRANSLUMINAL CORONARY ANGIOPLASTY

Principal investigators	No. of patients	Protocol	Time to early graded exercise testing	Follow-up angiography
Scholl et al ¹	54	85% maximal predicted heart rate	1 mo	20 mo
Rosing et al ²	45	Symptom-limited. Treadmill	8 mo	—
Wijns et al ³	89	Symptom-limited. Bicycle	4.7 mo	12 mo
Miri et al ⁸	71	Symptom-limited. Bicycle	2 d	28 mo
Rod et al ⁵	14	Symptom-limited. Bicycle	7.8 d	—
Meier et al ⁶	160	Symptom-limited. Bicycle	2 d	—
Ernst et al ⁷	25	Symptom-limited. Bicycle	14 d	—

though thallium scans improved significantly immediately after angioplasty, evidence of "ischemia" was still present in over one-third of their patients. By three months, the perfusion scans normalized in all patients, with no further changes at six or nine months.

CONCLUSIONS

The results of this study suggest that early symptom-limited stress electrocardiography following percutaneous transluminal coronary angioplasty is of no diagnostic value for the prediction of future coronary events. In addition, maximal exercise testing may carry some added risk for acute complications.

The optimal timing of the test requires further investigation and will ultimately depend on the individual clinical circumstances of each patient. Delaying stress testing for four weeks or limiting the test to a moderate workload are alternative solutions. The use of testing at three or six months may also maximize early detection of recurrence. If predischarge testing is to be used routinely, test endpoints and timing need to be standardized and the potential for complications related to the exercise test in this population needs to be considered.

REFERENCES

- Scholl JM, Chatman BR, David PR, et al. Exercise electrocardiography and myocardial scintigraphy in the serial evaluation of the results of percutaneous transluminal coronary angioplasty. *Circulation* 1982; 66:380-390.

2. Rosing DR, VanRaden MJ, Mincemoyer RM, et al. Exercise, electrocardiographic and functional responses after percutaneous transluminal coronary angioplasty. *Am J Cardiol* 1984; 53:36C-41C.
3. Wijns W, Serruys PW, Reiber JH, et al. Early detection of restenosis after successful percutaneous transluminal coronary angioplasty by exercise-redistribution thallium scintigraphy. *Am J Cardiol* 1985;55:357-361.
4. Wijns W, Serruys PW, Simoons ML, et al. Predictive value of early maximal exercise test and thallium scintigraphy after successful percutaneous transluminal coronary angioplasty. *Br Heart J* 1985; 53:194-200.
5. Rod JL, Foster C, Schmidt DH. Evaluation of percutaneous transluminal coronary angioplasty by symptom-limited graded exercise testing. *J Cardiac Rehabil* 1984; 4:70-73.
6. Meier B, Gruentzig AR, Siegenthaler WE, Schlumpf M. Long-term exercise performance after percutaneous transluminal coronary angioplasty and coronary artery bypass grafting. *Circulation* 1983; 68:796-802.
7. Ernst SM, Hillebrand FA, Klein B, Ascoop CA, van Tellingen C, Plokker HW. The value of exercise tests in the follow-up of patients who underwent transluminal coronary angioplasty. *Int J Cardiol* 1985; 7:267-279.
8. Miri M, Meyer J, Erbel R. Early and long-term exercise performance after percutaneous transluminal coronary angioplasty of the left anterior descending coronary artery. *Acta Med Iugosi* 1986; 40:107-113.
9. Rosing DR, Cannon RO, Watson RM, et al. Three year anatomic, functional and clinical follow-up after successful percutaneous transluminal coronary angioplasty. *J Am Coll Cardiol* 1987; 9:1-7.
10. Okada RD, Lim YL, Boucher CA, Pohost GM, Chesler DA, Block PC. Clinical, angiographic, hemodynamic, perfusional and functional changes after one-vessel left anterior descending coronary angioplasty. *Am J Cardiol* 1985; 55:347-356.
11. Dash H. Delayed coronary occlusion after successful percutaneous transluminal coronary angioplasty: association with exercise testing. *Am J Cardiol* 1983; 52:1143-1144.
12. Przbojewski JZ, Weich HF. Acute coronary thrombus formation after stress testing following percutaneous transluminal coronary angioplasty. *S Afr Med J* 1985; 67:378-382.
13. Przbojewski JZ, Weich HF. Late symptomatic exercise-induced coronary vasospasm after percutaneous transluminal coronary angioplasty. *S Afr Med J* 1986; 69:314-320.
14. Nygaard TW, Beller GA, Mentzer RM, Gibson RS, Moeller CM, Burwell LR. Acute coronary occlusion with exercise testing after initially successful coronary angioplasty for acute myocardial infarction. *Am J Cardiol* 1986; 57:687-688.
15. Ellestad M. Stress Testing, 2nd ed. Philadelphia, FA Davis Company, 1980, pp 113-116.
16. Schlant RC, Blomquist CG, Brandenburg RO, et al.-A report of the American College of Cardiology/American Heart Association task force on assessment of cardiovascular procedures. *J Am Coll Cardiol* 1986; 8:725-738.
17. Chaitman BR. The changing role of the exercise electrocardiogram as a diagnostic and prognostic test for chronic ischemic heart disease. *J Am Coll Cardiol* 1986; 8:1195-1210.
18. DeBusk RF, Blomquist CG, Kouchoukos NT, et al. Identification and treatment of low-risk patients after acute myocardial infarction and coronary-artery bypass graft surgery. *N Engl J Med* 1986; 314:161-166.
19. Stone PH, Turi ZG, Muller JE, et al. Prognostic significance of the exercise treadmill test performance 6 months after myocardial infarction. *J Am Coll Cardiol* 1986; 8:1007-1017.
20. Waters DD. Exercise testing after myocardial infarction: a perspective. *J Am Coll Cardiol* 1986; 8:1018-1019.
21. Weiner DA. Role of exercise testing after myocardial infarction. *J Am Coll Cardiol* 1986; 8:1020-1021.
22. Gruntzig AR, Senning A, Siegenthaler WE. Nonoperative dilatation of coronary-artery stenosis. *N Engl J Med* 1979; 301:61-68.
23. Simpfendorfer C, Belardi J, Bellamy G, Galan K, Franco I, Hollman J. Frequency, management, and follow-up of patients with acute coronary occlusions after percutaneous transluminal coronary angioplasty. *Am J Cardiol* 1987; 59:267-269.
24. Shaw RE, Meler RK, Stertz SH, Clark DA. Restenosis after coronary angioplasty. *Cardiology* 1987; 74:42-84.
25. Kemp H. Coronary angioplasty and the persistent problem of restenosis. *J Am Coll Cardiol* 1986; 8:1277-1278.
26. Manyari DE, Knudtson M, Kloiber R, Roth D. Sequential thallium-201 myocardial perfusion studies after successful percutaneous transluminal coronary artery angioplasty: delayed resolution of exercise-induced scintigraphic abnormalities. *Circulation* 1988; 77:86-95.