

Evaluation and treatment of generalized pruritus

GARY R. KANTOR, MD

■ Generalized pruritus may be symptomatic of dermatologic disease or it can be a manifestation of a systemic disorder. When the cause of generalized pruritus is not immediately evident, careful clinical and laboratory evaluation and close follow-up are needed to ensure appropriate diagnosis and treatment.

□ INDEX TERM: PRURITUS □ CLEVE CLIN J MED 1990; 57:521-526

PRURITUS is the most common dermatologic symptom that patients present. It accompanies a wide range of skin diseases, most of them characterized by primary lesions such as papules, vesicles, and plaques (*Table 1*).

Generalized pruritus of undetermined origin (GPUO) is continuous itching of at least 3 weeks duration that is unresponsive to 2 weeks of conservative management and unassociated with primary skin lesions. Systemic diseases may be associated with generalized pruritus (*Table 2*); but the prevalence of the association ranges from 14% to 50%, depending on the source of the report.¹⁻⁴

This article discusses the systemic diseases associated with generalized pruritus, and provides recommendations for evaluating the patient with generalized pruritus.

SYSTEMIC DISEASES ASSOCIATED WITH PRURITUS

Uremia

Chronic renal failure is the systemic disease most commonly associated with generalized pruritus.⁵ Up to

90% of patients on hemodialysis have either localized or generalized pruritus.⁶ These patients complain of severe, paroxysmal itching that tends to be most severe in the summer. In one study, 25% of patients had discomfort soon after hemodialysis and 42% had their most severe pruritus during hemodialysis.⁶

The etiology of uremic pruritus is unknown, but several mechanisms have been proposed. Increased numbers of mast cells have been found in the spleen and bone marrow of uremic patients who have no evidence of systemic mast cell disease.⁷ The mast cell proliferation may be related to elevated parathyroid hormone levels. Histamine has been implicated, given the increased sensitivity of uremic patients to intradermal histamine.⁸ Calcium hemostasis is defective in uremia, and medical or surgical parathyroidectomy improves the associated pruritus.^{9,10} Other possibilities include peripheral neuropathy, xerosis, hypermagnesemia, or accumulation of an unknown pruritogenic substance.

Ultraviolet B phototherapy is the most effective treatment of pruritus associated with uremia. Most patients improve within six treatments; if a second course of therapy is necessary, the response is more rapid.¹¹ Ultraviolet A has some advantages because of its wider availability and higher safety profile.¹² Emollients are useful for associated xerosis. Other effective treatments include topical capsaicin,¹³ lidocaine infusion, intravenous heparin, and oral charcoal. It also helps to

From the Section of Dermatopathology, Division of Dermatology, Department of Medicine, Hahnemann University School of Medicine, Philadelphia.

Address reprint requests to G.R.K., Division of Dermatology, Hahnemann University School of Medicine, Broad and Vine Streets, Mail Stop 401, Philadelphia, Pennsylvania 19102-1192.

TABLE 1
SKIN DISEASES ASSOCIATED WITH PRURITUS

Xerosis
Urticaria
Sunburn
Dermatitis herpetiformis
Contact dermatitis
Fungal infections
Fiberglass dermatitis
Pruritic urticarial papules and plaques of pregnancy
Pediculosis
Miliaria
Folliculitis
Polymorphous light eruption
Pityriasis rosea
Lichen simplex chronicus
Psoriasis
Scabies
Insect bites
Atopic dermatitis
Drug reactions
Mycosis fungoides
Bullous pemphigoid
Pemphigus foliaceus
Lichen planus

lower the dialysate magnesium concentration for patients undergoing hemodialysis.

Hepatobiliary diseases

Pruritus may be associated with primary biliary cirrhosis, drug-induced cholestasis (as occurs with chlorpropamide, oral contraceptives, and anabolic steroids), biliary obstruction from intrahepatic or extrahepatic sources, and cholestasis of pregnancy. Cyclic premenstrual pruritus due to recurrent cholestasis has also been reported.¹⁴

Pruritus may develop weeks or months before other signs and symptoms of hepatic disease. Itching often begins on the hands, feet, and other pressure areas of the skin and is especially severe at night. The presence of the “butterfly sign” suggests chronic obstructive hepatobiliary disease.¹⁵ The butterfly sign is characterized by the relative hypopigmentation of the skin on the mid- to upper back, surrounded by postinflammatory hyperpigmentation.

Although accumulated bile salts have been implicated as the cause of pruritus in hepatobiliary disease, this notion is controversial.¹⁶ Factors that support the role of bile salts include relief of pruritus with bile salt-binding resins such as cholestyramine¹⁷ and the production of pruritus when bile salts are applied to the skin.¹⁸ Nevertheless, other studies on the correlation of itching with serum and skin bile salt concentrations have had conflicting results.^{19,20} Furthermore, in some patients phenobarbital relieves pruritus without a decrease in serum bile acids, while in others, androgen therapy is

TABLE 2
SYSTEMIC DISEASES ASSOCIATED WITH GENERALIZED PRURITUS

System involved	Diseases that cause pruritus
Renal	Uremia
Hepatic (obstructive hepatobiliary disease)	Primary biliary cirrhosis Drugs (chlorpropamide, oral contraceptives, anabolic steroids) Extrahepatic biliary obstruction Intrahepatic cholestasis of pregnancy
Hematologic	Polycythemia vera Iron deficiency anemia Multiple myeloma and paraproteinemia Mastocytosis Thyroxinosis
Endocrine	Hypothyroidism Diabetes
Neurologic	Infarcts Brain abscess Multiple sclerosis Tumor
Lymphoreticular neoplasms	Hodgkin's disease Lymphoma
Internal malignancy	Systemic carcinoma Carcinoid syndrome
Miscellaneous	AIDS Sjögren's syndrome Dumping syndrome Bullous pemphigoid

beneficial despite the associated increase in serum bile acid concentration.²¹

The general consensus is that pruritus with hepatobiliary disease is an effect of bile salts on the liver, not the skin. The pruritus is related to the release of an unknown pruritogenic substance that binds to cholestyramine and is photolabile.²² Supporting this theory is the beneficial effect of rifampin on pruritus associated with primary biliary cirrhosis; this drug inhibits bile acid uptake in the liver and stimulates hepatic enzymatic degradation of bile salts.²³

Pruritus associated with hepatobiliary disease can be severe and can interfere with quality of life. Cholestyramine is the first treatment of choice; in resistant cases, the drug may be combined with phototherapy.²⁴ UVA or UVB phototherapy alone is less useful than in the treatment of pruritus associated with uremia. Naloxone, a central opioid antagonist, is beneficial for intractable itching related to primary biliary cirrhosis.²⁵ Besides rifampin,²³ other treatments include surgical external diversion of bile²⁶ and plasma perfusion through charcoal coated glass beads.²⁷

Hematologic diseases

Patients with polycythemia vera may present with generalized pruritus. The itching is described as “prickling” in quality and lasts from minutes to hours. It starts

after the patient emerges from a shower or bath as the skin begins to cool—hence the term “bath pruritus.” The cause of the pruritus is unknown, although elevated plasma and urine histamine levels were detected in one study and correlated with an increase in circulating basophils.²⁸ Earlier reports claimed that cimetidine was beneficial for these patients,^{29,30} but subsequent controlled studies did not confirm efficacy.³¹ Aspirin and cyproheptadine (Periactin) have also been used for treatment.

Generalized pruritus has been reported in association with iron deficiency anemia.³² Although small studies have described relief of pruritus with iron therapy,^{33,34} the relationship between generalized pruritus and iron deficiency is questionable. A large series of patients needs to be investigated before any conclusions can be made.

Rarely, benign gammopathy³⁵ and multiple myeloma³⁶ are associated with pruritus. Systemic mastocytosis has also been reported in a patient presenting with generalized pruritus and no observable skin lesions.³⁷




FIGURE 1. Flow chart for evaluating patients with generalized pruritus of unknown origin.

Endocrine disorders

Pruritus may occur in up to 5% of patients with thyrotoxicosis and subside with treatment of the thyroid disorder.³⁸ Some patients with hyperthyroidism may present with pruritus.³⁹ Various mechanisms have been suggested for the pruritus associated with hyperthyroidism; these include cutaneous vasodilation, lowered pain threshold, enhanced sen-

sitivity to catecholamines, activation of kalli-krein-kinin system, and prostaglandin mediation.⁴⁰

Hypothyroidism has not been linked to generalized pruritus in controlled studies. Dry skin and associated itching often accompany the hypothyroid state; but GPUO as defined above rarely is solely attributable to

TABLE 3
LABORATORY WORKUP FOR GENERALIZED PRURITUS

Indications	Suggested tests
Screening	CBC with differential BUN, creatinine Alkaline phosphatase, bilirubin, SGOT T ₄ , TSH Glucose Chest radiograph
Follow-up	Stool for occult blood (for patients >40 years) Serum iron, ferritin Serum protein electrophoresis, serum immunoelectrophoresis Urine 5-hydroxyindoleacetic acid (5-HIAA) Skin biopsy for direct immunofluorescence Skin biopsy for special stain (to exclude mastocytosis) Stool for ova and parasites Radiologic studies as indicated

hypothyroidism.

The association of diabetes mellitus with pruritus has been largely anecdotal.⁴¹ A report that compared 300 diabetic patients with controls showed an increased prevalence of pruritus vulvae, but not generalized pruritus, in the diabetic population.⁴² The presence of diabetes does not satisfactorily explain GPUO, although persistent scalp pruritus may be related to diabetes.⁴³

Neurologic disorders

Unilateral pruritus has been reported secondary to central nervous system infarct⁴⁴ or brain abscess.⁴⁵ Paroxysmal itching has also been described in multiple sclerosis.^{46,47} Intractable nasal pruritus may be a sign of brain tumor⁴⁸ or atypical angina,⁴⁹ suggesting that itching may originate in the cerebral cortex. Bombesin, an endogenous opioid peptide, may be a central mediator of pruritus⁵⁰; when administered intrathecally to animals, it produces intense and long-lasting pruritus. This mediator and its antagonist require further investigation.

Lymphoreticular malignancies

Generalized pruritus may be associated with or the presenting symptom of Hodgkin's disease. Opinions conflict regarding the prognostic significance of pruritus in this disease. The Ann Arbor Conference on Hodgkin's disease in 1971 dropped pruritus as a criterion for B categorization, but two later reports disputed this action.^{51,52} Further studies are needed to resolve this question.

Non-Hodgkin's lymphoma and mycosis fungoides may also be associated with generalized pruritus.

Internal malignancy

Systemic malignancy is a rare cause for GPUO and does not warrant extensive, cost-consuming, laboratory evaluation.⁵³ The association of pruritus and internal malignancy was reported in 1942⁵⁴; however, in most instances pruritus was accompanied by some other sign such as rash, and was only rarely the sole presenting symptom. Cormia stated in 1965 that pruritus was an uncommon, although important, symptom of systemic carcinoma—especially if it was severe, prolonged, and localized to the pretibial legs, inner thighs, upper trunk, and extensor surfaces of the upper arms.⁵⁵

Subsequent studies reported that 3% to 26% of patients with generalized pruritus have an underlying malignancy.¹⁻⁴ However, when a population of 125 patients with generalized pruritus were followed over a 6-year period, there was no significant increase in malignant neoplasms, except for a slight increase in malignant lymphomas.⁵⁶

Carcinoid tumors⁵⁷ have been associated with pruritus—presumably because of a direct effect of mediators on cutaneous vasculature.

Other systemic disorders

Patients with acquired immunodeficiency syndrome may present with generalized pruritus.⁵⁸ Pruritus has been reported in 30% of HIV-infected prostitutes.⁵⁹ The pruritus associated with HIV infection is generally refractory to treatment, although psoralen and UVA phototherapy (PUVA) may benefit some patients.⁶⁰ In many AIDS patients, pruritus may be associated with bacterial folliculitis, with or without atopic eczema or scabies.

Pruritus can be the presenting symptom for Sjögren's syndrome,⁶¹ dumping syndrome,⁶² and bullous pemphigoid.⁶³

Psychogenic disorders

Pruritus is aggravated by stress; one study correlated high levels of psychological stress with increased experimentally induced itching.⁶⁴ Nevertheless, a diagnosis of psychogenic pruritus should be made with great caution and after careful observation. Itching may reasonably be ascribed to psychogenic factors if there is a relevant history of major psychiatric disease. Fluoxetine (Prozac), 10 mg to 20 mg daily, may be useful for some of these patients.

EVALUATING THE PATIENT

The patient with GPUO requires careful evaluation and long-term follow-up (Figure 1). In the absence of primary skin disease, scabies, and lymphadenopathy or organomegaly at the initial visit, a 2-week trial of

symptomatic therapy is appropriate before starting laboratory evaluation.

Conservative symptomatic measures include educating the patient about bathing procedures, especially in fall and winter in cold climates—for example, the use of lukewarm bath water, mild soaps such as Dove or Basis, pat drying instead of rubbing with the bath towel, limited frequency, and immediate application of a moisturizing lotion such as Moisturel or Nivea. A sedating H₁-antagonist such as hydroxyzine, 10 mg to 25 mg bid, as initial therapy is helpful for some patients. Fluorinated topical corticosteroids should be avoided if there is no primary cutaneous disease, but sparing use of nonfluorinated corticosteroid creams and lotions has an antipruritic effect. Menthol-containing lotions such as Sarna appeal to some patients. Also useful is topical pramoxine alone (Prax), with menthol 0.5% (Prame-Gel), or with hydrocortisone (Pramasone). Topical antihistamines should not be used because of their potential for sensitization.

If 2 weeks of symptomatic therapy does not bring relief or significant improvement of symptoms, then initial screening laboratory tests are indicated to rule out underlying systemic disease (Table 3), including complete blood count, blood urea nitrogen, creatinine, liver enzymes, thyroid studies, blood glucose, and chest

radiograph. Screening for fecal occult blood is appropriate for patients older than 40 years.

If these screening tests are normal, then additional studies may be performed as indicated (Table 3). For example, a perilesional skin biopsy may reveal clinically masked primary skin disease such as dermatitis herpetiformis and bullous pemphigoid. Extensive radiographic examinations are unwarranted unless the history or physical examination suggests a need for them.

If this secondary evaluation is unrevealing, then continued symptomatic therapy is warranted. At this point, alternative antihistamines may be indicated, such as cyproheptadine and doxepin (Sinequan). Nonsedating antihistamines have not been helpful in these patients.

Systemic antibiotics may be needed if there are extensive excoriations and secondary infection. A trial of UVB or PUVA phototherapy may be indicated. Localized, disabling pruritus may respond to a soft cast of zinc oxide-impregnated gauze (Domepaste) applied over a soothing topical corticosteroid ointment or tar-based paste.

Reassessment and reevaluation of these patients is crucial; for example, it is not unreasonable to repeat laboratory studies at intervals of 3 to 6 months. Some patients will manifest features of systemic disease after their initial visit,⁴ and only careful follow-up will ensure timely diagnosis of occult medical problems.

REFERENCES

1. Rajka G. Investigation of patients suffering from generalized pruritus, with special references to systemic disease. *Acta Derm Venereol (Stockh)* 1966; **46**:190-194.
2. Lyell A. The itching patient. A review of the causes of pruritus. *Scott Med J* 1972; **17**:334-338.
3. Beare JM. Generalized pruritus. A study of 43 cases. *Clin Exp Dermatol* 1976; **1**:343-352.
4. Kantor GR, Lookingbill DP. Generalized pruritus and systemic disease. *J Am Acad Dermatol* 1983; **9**:375-382.
5. Denman ST. A review of pruritus. *J Am Acad Dermatol* 1986; **14**:375-392.
6. Gilchrist BA, Stern RS, Steinman TI, Brown RS, Arndt KA, Anderson WW. Clinical features of pruritus among patients undergoing maintenance hemodialysis. *Arch Dermatol* 1982; **118**:154-156.
7. Neiman RS, Bischel MD, Lukes RJ. Uraemia and mast-cell proliferation. *Lancet* 1972; **1**:959.
8. Stahle-Backdahl M, Hagermark O, Lins LE. The sensitivity of uremic and normal human skin to histamine. *Arch Derm Venerol (Stockh)* 1988; **68**:230-235.
9. Massry SG, Popovtzer MM, Coburn JW, Makoff DL, Maxwell MH, Kleeman CR. Intractable pruritus as a manifestation of secondary hyperparathyroidism in uremia. *N Engl J Med* 1968; **279**:697-700.
10. Hampers CL, Katz AI, Wilson RE, Merrill JP. Disappearance of "uremic" itching after subtotal para-thyroidectomy. *N Engl J Med* 1968; **279**:695-697.
11. Gilchrist BA, Rowe JW, Brown RS, Steinman TI, Arndt KA. Ultraviolet phototherapy of uremic pruritus. *Ann Intern Med* 1979; **91**:17-21.
12. Hindson C, Taylor A, Martin A, Downey A. UVA light for relief of uraemic pruritus. *Lancet* 1981; **1**:215.
13. Breneman DL, Cardone JS, Kauffman RF, Lather RM. Topical capsaicin for treatment of pruritus related to hemodialysis. *Clin Pharmacol Ther* 1989; **45**:188A.
14. Dahl MGC. Premenstrual pruritus due to recurrent cholestasis. *Trans St Johns Hosp Dermatol Soc* 1970; **56**:11-13.
15. Goldman RD, Rea TH, Cinque J. The "Butterfly" sign—a clue to generalized pruritus in a patient with chronic obstructive hepatobiliary disease. *Arch Dermatol* 1983; **119**:183-184.
16. Ghent CN, Bloomer JR. Itch in liver disease: facts and speculations. *Yale J Biol Med* 1979; **52**:77-82.
17. Van Itallie TB, Hashim SA, Crampton RS, Tennent DM. The treatment of pruritus and hypercholesterolemia of primary biliary cirrhosis with cholestyramine. *N Engl J Med* 1961; **265**:469-474.
18. Varadi DP. Pruritus induced by crude bile and purified bile acids—experimental production of pruritus in human skin. *Arch Dermatol* 1974; **109**:678-681.
19. Bloomer JR, Allen RM, Klatskin G. Serum bile acids in primary biliary cirrhosis. *Arch Intern Med* 1976; **136**:57-61.
20. Ghent CN, Bloomer JR, Klatskin G. Elevations in skin tissue levels of bile acids in human cholestasis: relation to serum levels and to pruritus. *Gastroenterology* 1978; **73**:1125-1130.
21. Alva J, Iber FL. Relief of the pruritus of jaundice with methandrolone and speculations of the nature of pruritus in liver disease. *Am J Med Sci* 1965; **250**:60-65.
22. Ghent CN. Pruritus of cholestasis is related to effects of bile salts on the liver, not the skin. *Am J Gastroenterol* 1987; **82**:117-118.
23. Ghent CN, Carruthers SG. Treatment of pruritus in primary biliary cirrhosis with rifampin—results of a double-blind, crossover, ran-

- domized trial. *Gastroenterology* 1988; **94**:488–493.
24. Cerio R, Murphy GM, Sladen GE, MacDonald DM. A combination of phototherapy and cholestyramine for the relief of pruritus in primary biliary cirrhosis. *Br J Dermatol* 1987; **116**:265–267.
25. Bernstein JE, Swift R. Relief of intractable pruritus with naloxone. *Arch Dermatol* 1979; **115**:1366–1367.
26. Whittington PF, Whittington GL. Partial external diversion of bile for the treatment of intractable pruritus associated with intrahepatic cholestasis. *Gastroenterology* 1988; **95**:130–136.
27. Carey WD, Smith J, Asanuma Y, Malchesky P, Ferguson DR, Greenstreet R, Nose Y. Pruritus of cholestasis treated with plasma perfusion. *Am J Gastroenterol* 1981; **76**:330–337.
28. Gilbert HS, Warner RRP, Wasserman LR. A study of histamine in myeloproliferative disease. *Blood* 1966; **28**:795–806.
29. Easton P, Galbraith PR. Cimetidine treatment of pruritus in polycythemia vera. *N Engl J Med* 1978; **299**:1134.
30. Harrison AR, Littenberg G, Goldstein L, Kaplowitz N. Pruritus, cimetidine, and polycythemia. *N Engl J Med* 1979; **300**:433–434.
31. Bauer E, Densen P. Cimetidine for the treatment of pruritus. *N Engl J Med* 1979; **300**:370.
32. Lewiecki EM. Pruritus, a manifestation of iron deficiency. *JAMA* 1976; **236**:2319–2320.
33. Vickers CF. Iron deficiency and the skin. *Br J Dermatol* 1973; **89**:10.
34. Salem HH, Van der Weyden MB, Young IF, Wiley JS. Pruritus and severe iron deficiency in polycythemia vera. *Br Med J* 1982; **285**:91–92.
35. Zelicovici Z, Lahav M, Cahane P, Bianu G. Pruritus as a possible early sign of paraproteinemia. *Isr J Med Sci* 1969; **5**:1079–1081.
36. Erskine JG, Rowan RM, Alexander JO, et al. Pruritus as a presentation of myelomatosis. *Br Med J* 1977; **1**:687–688.
37. Kendall ME, Fields JP, King LE. Cutaneous mastocytosis without clinically obvious skin lesions. *J Am Acad Dermatol* 1984; **10**:903–905.
38. Caravati CM, Richardson DR, Wood BT, Cawley EP. Cutaneous manifestations of hyperthyroidism. *South Med J* 1969; **62**:1127–1130.
39. Barrow MV, Bird ED. Pruritus in hyperthyroidism. *Arch Derm* 1966; **93**:237–238.
40. Barnes HM, Sarkany I, Calnan CD. Pruritus and thyrotoxicosis. *Trans St Johns Hosp Dermatol Soc* 1974; **60**:59–62.
41. Jelinek JE. The skin in diabetes mellitus: cutaneous manifestations, complications, and associations [In] Kopf AW, Andrade R (eds): *Yearbook of Dermatology*, 1970. Chicago, Year Book Medical Publishers, Inc. 1970, pp 5–35.
42. Neilly JB, Martin A, Simpson N, MacCuish AC. Pruritus in diabetes mellitus: investigation of prevalence and correlation with diabetes control. *Diabetes Care* 1986; **9**:273–275.
43. Scribner M. Diabetes and pruritus of the scalp. *JAMA* 1977; **237**:1559.
44. King CA, Huff FJ, Jorizzo JL. Unilateral neurogenic pruritus: paroxysmal itching associated with central nervous system lesions. *Ann Intern Med* 1982; **97**:222–223.
45. Sullivan MJ, Drake ME. Unilateral pruritus and Nocardia brain abscess. *Neurology* 1984; **34**:828–829.
46. Osterman PO. Paroxysmal itching in multiple sclerosis. *Br J Dermatol* 1976; **95**:555–558.
47. Yamamoto M, Yabuki S, Hayabara T, Otsuki S. Paroxysmal itching in multiple sclerosis: a report of three cases. *J Neurol Neurosurg Psychiatry* 1981; **44**:19–22.
48. Andreev VC, Petkov I. Skin manifestations associated with tumours of the brain. *Br J Dermatol* 1975; **92**:675–678.
49. Reichstein RP, Stein WG. Nasal pruritus as atypical angina? *N Engl J Med* 1983; **309**:667.
50. Gmerek DE, Cowan A. Bombesin—a central mediator of pruritus? *Br J Dermatol* 1983; **109**:239.
51. Feiner AS, Mahmood T, Wallner SF. Prognostic importance of pruritus in Hodgkin's disease. *JAMA* 1978; **240**:2738–2740.
52. Gobbi PG, Attardo-Parrinello G, Lattanzio G, Rizzo SC, Ascari E. Severe pruritus should be a B-symptom in Hodgkin's disease. *Cancer* 1983; **51**:1934–1936.
53. Lober CW. Should the patient with generalized pruritus be evaluated for malignancy. *J Am Acad Dermatol* 1988; **19**:350–352.
54. Becker SW, Kahn D, Rothman S. Cutaneous manifestations of internal malignant tumors. *Arch Dermat Syph* 1942; **45**:1069–1080.
55. Cormia FE. Pruritus, an uncommon but important symptom of systemic carcinoma. *Arch Derm* 1965; **92**:36–39.
56. Paul R, Paul R, Jansen CT. Itch and malignancy prognosis in generalized pruritus: a 6-year follow-up of 125 patients. *J Am Acad Dermatol* 1987; **16**:1179–1182.
57. Menzel CE. Cutaneous manifestations of the malignant carcinoid syndrome. *Ann Intern Med* 1963; **58**:989–993.
58. Shapiro RS, Samorodin C, Hood AF. Pruritus as a presenting sign of acquired immunodeficiency syndrome. *J Am Acad Dermatol* 1987; **16**:1115–1117.
59. Van De Perre P, Carael M, Robert-Guroff M, et al. Female prostitutes: a risk group for infection with human T-cell lymphotropic virus type III. *Lancet* 1985; **2**:524–526.
60. Gorin I, Lessana-Leibowitch M, Fortier P, Leibowitch J, Escande JP. Successful treatment of the pruritus of human immunodeficiency virus infection and acquired immunodeficiency syndrome with psoralens plus ultra-violet A therapy. *J Am Acad Dermatol* 1989; **20**:511–513.
61. Feuerman EJ. Sjogren's syndrome presenting as recalcitrant generalized pruritus. *Dermatologica* 1968; **137**:74–86.
62. Harries AD, Tredree R, Heatley RV, Rhodes J, Holt PJA. Pruritus as a manifestation of the dumping syndrome. *Br J Dermatol* 1982; **107**:707–709.
63. Barker DJ. Generalized pruritus as a presenting feature of bullous pemphigoid. *Br J Dermatol* 1983; **109**:237–239.
64. Edwards AE, Shellow WVR, Wright ET, Dignam TF. Pruritic skin disease, psychological stress, and the itch sensation. *Arch Dermatol* 1976; **112**:339–343.