



ALAN J. TAEGE, MD

Department of Infectious Disease, Cleveland Clinic

Listeriosis: Recognizing it, treating it, preventing it

ABSTRACT

Listeria monocytogenes has become a major pathogen in foodborne illness. It most often affects patients who are pregnant, at the extremes of life, or immunocompromised in some way. A variety of clinical manifestations are possible, but bacteremia and meningitis are most common. This article reviews the epidemiology, microbiology, populations at risk, clinical manifestations, treatment, and prevention of listeriosis.

KEY POINTS

In suspected or proven cases of listerial infection, prompt treatment is crucial. The first-line treatment is the combination of either ampicillin or penicillin plus gentamicin. Patients allergic to penicillin can receive trimethoprim-sulfamethoxazole instead.

L. monocytogenes can survive and even slowly proliferate at 4° to 10°C. Therefore, refrigerating food does not completely prevent listeriosis.

Groups at risk are pregnant women, neonates, immunocompromised people, and the very elderly.

Meningitis and bacteremia are the most common manifestations of listeriosis, and more than 20% of people affected die.

THE RECENT WIDESPREAD OUTBREAK of listeriosis associated with tainted processed meats from a food manufacturing plant in Michigan is responsible for 100 illnesses and 15 deaths thus far, refocusing attention on the potential menace of *Listeria monocytogenes* to public health. Although this organism has been recognized as a human pathogen for 70 years,¹ until recent foodborne outbreaks occurred it was an uncommon infection associated with sporadic disease and fostered only a passing interest.^{2,3}

SOURCES OF PAST OUTBREAKS

Epidemics of listeriosis have been linked to many foods. Some of the more publicized and larger outbreaks have been ascribed to soft cheese,⁴ cole slaw,⁵ deli meats,⁶ hot dogs,⁷ and pasteurized milk.^{8,9} Other foods involved in outbreaks have included raw vegetables, raw milk, fish, pâté,¹⁰ raw and undercooked poultry,⁷ ice cream,¹⁰ sausage,¹¹ raw eggs, and prepackaged mixed salads.¹¹ Any meat can harbor *L. monocytogenes*, but chicken appears to support its growth best.¹⁰

LISTERIA'S MICROBIOLOGIC PROFILE

Although there are at least seven species of *Listeria*, only *L. monocytogenes* appears to cause disease in humans.

L. monocytogenes is a gram-positive bacillus that displays tumbling motility and beta-hemolysis. In mixed cultures it is difficult to isolate and is often mistakenly identified as a contaminant diphtheroid or corynebacterium. In partially treated patients, culture isolates of *Listeria* may resemble *Hemophilus influenzae*, leading to further confusion.




FIGURE 1. *Listeria monocytogenes* (arrows) inside macrophages in a cerebrospinal fluid sample from a patient with listerial meningitis. Oil, $\times 1,000$.

SOURCE: COURTESY OF GERALDINE S. HALL, PHD

L. monocytogenes is ubiquitous.¹² It prefers moist environments¹³ and can be isolated from soil, decaying vegetation, sewage, manure, and silage.⁹ It is also part of the fecal flora of birds, fish, and mammals. Five to ten percent of humans may carry it in their intestinal tracts.^{9,12} It is responsible for epizootics in herd animals and causes abortions, stillbirths, and the “circling disease” (a form of basilar meningitis) in sheep.¹⁴

This organism has several capabilities that make it particularly dangerous:

It can survive at low temperatures. In fact, because *L. monocytogenes* can proliferate at 4° to 10°C, cold is used in the laboratory to help isolate it.¹² Refrigeration therefore does not completely protect against listeriosis.

It survives inside human phagocytic cells, shielding itself from many of the host’s humoral defenses (FIGURE 1).^{10,15} Therefore, cell-mediated immunity is critical in fighting this pathogen.

It expresses listerolysin O, a hemolysin similar to streptolysin O.^{10,12,15} This is the organism’s major virulence factor. When the organism is engulfed by a macrophage, the macrophage encloses it in a vacuole called a phagosome. Listerolysin O dissolves the phagosome and allows the organism to escape into the macrophage’s cytoplasm, evading destruction. (The presence of iron, particularly in iron-overload states, further adds to its virulence.^{12,16})

■ POPULATIONS AT RISK

Although *L. monocytogenes* can infect healthy people,^{11,17} the vast majority afflicted are either at the extremes of life (< 1 month or > 60 years old), pregnant, or immunocompromised in some way.

Pregnant women account for more than 25% of cases,¹² most commonly in the third trimester, although disease has been recorded throughout all trimesters.^{2,14}

Patients with malignancy or some alteration of cell-mediated immunity account for approximately 70% of nonperinatal infections.¹² Chemotherapy, particularly with fludarabine plus prednisone to treat chronic lymphocytic leukemia, enhances the risk of infection with *L. monocytogenes*.^{18,19} Organ transplant recipients,²⁰ particularly renal transplant recipients, are at risk because of their immunosuppressed status.² HIV-positive patients have perhaps a 150 times higher risk of *L. monocytogenes* infection than healthy persons.²¹

Other groups at risk are people with diabetes, renal failure, alcohol abuse,¹⁴ intravenous drug abuse,¹¹ hypochlorhydria,^{7,12} and iron overload states.¹⁶

■ INCIDENCE AND MORTALITY

The estimated annual incidence of listeriosis is approximately 7.4 cases/million population—about 2,000 cases in the United States each year.^{7,12,22} However, these figures may be low, due to under-recognition and under-reporting.

Of persons infected, 20% to 30% die—about 400 deaths per year.⁷ This percentage is higher than in any other foodborne illness and reflects the host population¹¹: the mortality rate is even higher in patients that are older and more seriously immunocompromised¹⁴ and who have more serious underlying diseases.²³

■ ROUTES OF INFECTION

The usual route of infection is by eating contaminated food. Vertical transmission (ie, from mother to infant) also occurs, but other human-to-human transfer is very rare. Cutaneous infection by contact with infected

Any meat can harbor *L. monocytogenes*, but chicken supports its growth best



TABLE 1

**Characteristics of central nervous system infections:
Listeria monocytogenes vs other bacteria**

	LISTERIA MONOCYTOGENES	OTHER BACTERIA
Cerebrospinal fluid analysis		
Glucose level	Normal in > 60% of cases	Typically depressed
White blood cell count	High (50%–70% polymorphonuclear)	High (≥ 80% polymorphonuclear)
Protein level	High	High
Gram stain	Negative in ≥ 60% of cases	Frequently positive
Blood cultures	Positive in about 75% of cases	Less commonly positive
Clinical abnormalities		
Cranial nerve deficits	May occur	Rare
Ataxia and tremor	May occur	Rare
Altered mental status	Common	Less common
Nuchal rigidity	Absent in 15% to 20% of cases	Usually present
Cortical invasion	May occur	Uncommon
Seizures	May occur	Uncommon

animals has been documented in poultry handlers, ranchers, and veterinarians. The amount of inoculum necessary to produce infection is not known. Experts have postulated the presence of coinfecting microorganisms not yet identified.²⁴

CLINICAL MANIFESTATIONS

One to 2 days after exposure, some patients experience a self-limited gastrointestinal syndrome, which probably occurs more often than recognized.⁹ Clinical signs and symptoms may include fever, abdominal cramps, diarrhea, fatigue, headache, and myalgia. A listerial infection should be considered in suspected foodborne gastroenteritis when other pathogens are not isolated.¹² However, most laboratories do not specifically culture stool specimens for *Listeria*.

The more serious syndromes of listerial meningitis and bacteremia do not arise until days to weeks later, often making it difficult to trace the source of infection. The incubation period for these syndromes ranges from 11 to 70 days (mean: approximately 21 days).^{12,24} The amount of inoculum and host factors likely influence the duration between exposure and disease.

Central nervous system infection

Central nervous system (CNS) infection is the most common manifestation of listerial infection.² The mortality rate is greater than 30%, the highest of all causes of meningitis.¹²

Blood cultures are positive in 60% to 75% of patients with listerial CNS infections.¹⁴

CNS infection may manifest as meningitis, meningoencephalitis (rhombencephalitis), or abscesses.

Listerial meningitis. *Listeria* is the most common cause of meningitis in patients with hematologic malignancies, in transplant recipients, and in those receiving corticosteroids.¹² It is the third most common cause of meningitis in newborns (after *Escherichia coli* and group B streptococci),²⁶ and the third most common cause of meningitis in adults (after *Streptococcus pneumoniae* and *Neisseria meningitidis*).²⁷ Overall, it is the fifth-leading cause of bacterial meningitis in all age groups.¹² Interestingly, listerial meningitis in pregnant women, one of the groups at high risk, is exceedingly rare.^{2,12,28}

Listerial meningitis is similar in many ways to other forms of meningitis, except that meningismus occurs less often and mental status changes, movement disorders, and seizures are more common (TABLE 1).¹² The cere-

Serologic testing is not reliable for diagnosing listeriosis

TABLE 2

Differential diagnosis of listerial meningoencephalitis

Infectious diseases

Aspergillosis	Toxoplasmosis
Brucellosis	Tuberculosis
Cryptococcosis	Viral encephalitides
Coccidioidomycosis	Adenovirus
Cysticercosis	Echovirus
Herpes zoster	Epstein-Barr virus
Legionellosis	Herpes simplex virus
Lyme disease	Japanese virus
Melioidosis	Poliovirus
Mycoplasma	Rabies
Syphilis	St. Louis virus

Noninfectious conditions

Behçet syndrome	Multiple sclerosis
Carcinomatosis	Sarcoidosis
Central pontine myelinolysis	Vasculitis
Lymphomatous meningitis	Wegener granulomatosis
Miller-Fisher syndrome	

brospinal fluid usually has an elevated protein level and an elevated white blood cell count with a neutrophilic predominance, but the glucose level may be normal in up to 60% of cases.¹⁴ The Gram stain is negative in more than 60% of cases, possibly owing to the organism's intracellular life cycle.¹⁵ In the uncommon scenario of an adult without underlying disease, listerial meningitis may present with lymphocytic pleocytosis (ie, an increased number of lymphocytes in the cerebrospinal fluid),¹⁷ therefore increasing the possibility of incorrectly attributing the infection to a virus.

Meningoencephalitis and rhombencephalitis are very uncommon in adults^{29,30} and are almost never seen in neonates. Frequently, the patient is not immunocompromised. Rhombencephalitis (ie, inflammation of the medulla, pons, and cerebellum) manifests as a biphasic process, beginning with fever, headache, nausea, and vomiting and followed by neurologic findings. Because *L monocytogenes* has an affinity for the brain stem, common neurologic findings include asymmetric cranial nerve deficits, cerebellar abnormalities, tremor,¹⁴ hemiparesis, seizures, and respiratory impairment. Personality changes have also been noted.

The differential diagnosis for this type of syndrome is broad and should include many infectious and noninfectious diseases (TABLE 2). The mortality rate in patients with rhombencephalitis may reach 50%. Neurologic sequelae are common, occurring in approximately 60%. The cerebrospinal fluid findings are not as impressive as in listerial meningitis and often more closely resemble those found in brain abscess.²⁹

Isolated bacteremia

Isolated *L monocytogenes* bacteremia presents as a nonspecific flulike illness with fever, arthralgia, myalgia, and headache. It most commonly affects pregnant women, usually in the third trimester, but cases may occur throughout pregnancy. Because the organism has an affinity for the placenta, listerial bacteremia may result in premature labor, stillbirth, or miscarriage. Maternal death is rare, but fetal mortality may approach 40%.³¹ If the disease is recognized, it is possible to treat the mother successfully and still allow the birth of a healthy infant.

Spontaneous resolution of primary bacteremia in pregnancy has been reported.² When bacteremia results in amnionitis, the baby must be delivered immediately, and maternal recovery usually follows. Infection during pregnancy does not predispose to a carrier state.¹⁰

Neonatal listerial bacteremia

In neonates, listerial bacteremia usually occurs during the first 2 days of life. It may manifest as granulomatosis infantisepticum, a severe disseminated form of bacteremia with widespread microabscesses and granulomas and hepatosplenomegaly. This is believed to represent in utero infection and is associated with a poor prognosis and a mortality rate of up to 50%.^{10,14} Although late-onset neonatal disease (10–14 days of age or older) most commonly presents as meningitis, bacteremia does occur in this age group as well.³¹

Listerial endocarditis

Listerial endocarditis is uncommon and usually has a subacute presentation.^{2,32} Often, the host is not immunocompromised but has underlying cardiac valvular abnormalities.



There appears to be a left-sided predominance of the affected valves.

Localized listerial infection

Localized listerial infections are rare.¹⁴ Lymph node abscesses, osteomyelitis, septic arthritis, pneumonia, empyema, and peritonitis have been described.³³ Cutaneous infections occur as occupational hazards for ranchers, veterinarians, and poultry workers. Myocarditis has been described in heart transplant recipients.³⁴

■ DIAGNOSIS AND TREATMENT

In suspected or proven cases of listeriosis, prompt initiation of antibiotic therapy is crucial. Diagnosis is established by culture, growth, isolation, and characterization of *L monocytogenes* from the blood and cerebrospinal fluid. Serologic testing is not reliable for diagnosis.

Antibiotic regimens

Ampicillin in maximal doses (2 g every 4 hours for adults, or 200–400 mg/kg/day for children) is the preferred agent to treat listeriosis, often in synergistic combination with gentamicin.^{12,14} Penicillin G may be as effective as ampicillin. However, ampicillin and penicillin are only bacteriostatic for *L monocytogenes*.

Trimethoprim and sulfamethoxazole are synergistic,³⁵ and the combination is accepted as a second-line therapy for patients allergic to penicillin. Trimethoprim-sulfamethoxazole is bactericidal and penetrates the cerebrospinal fluid well.¹⁴

Erythromycin as monotherapy demonstrates good antilisterial activity³⁶ but is considered a third-line therapy at present. It may be a suitable alternative for penicillin-allergic pregnant patients, in whom trimethoprim-sulfamethoxazole would be contraindicated. Erythromycin penetrates the meninges very poorly, even when the meninges are inflamed, and therefore would be a poor choice in treating meningitis or rhombencephalitis.

Vancomycin has some activity against *L monocytogenes*. Most authorities do not recommend using it for this purpose however, because of limited efficacy and because of the desire to limit the use of vancomycin in general.

TABLE 3

Avoiding foodborne listeriosis: Advice for patients

Advice for all persons

Cook all meat thoroughly

Wash all raw vegetables carefully

Keep raw meats separate from vegetables, cooked foods, and ready-to-eat foods during preparation to avoid cross-contamination

Wash hands, knives, and cutting boards after handling each uncooked food

Avoid unpasteurized milk products

Advice for pregnant or immunocompromised persons

Avoid soft cheeses such as Mexican-style, feta, Brie, Camembert, and blue cheese (hard cheeses, cream cheese, cottage cheese, and yogurt are OK)

Reheat leftovers and ready-to-eat foods such as hot dogs until steaming hot before eating

Avoid deli foods unless they are thoroughly heated before eating

SOURCE: ADAPTED FROM SCHUCHAT A, BROOME CV. INFECTIONS CAUSED BY *LISTERIA MONOCYTOGENES*. IN ISSELBACHER KJ, EDITOR. HARRISON'S PRINCIPLES OF INTERNAL MEDICINE 13TH ED. NEW YORK: MCGRAW-HILL, 1994; 631–632.

Bacterial resistance

An isolate of *L monocytogenes* that is resistant to trimethoprim-sulfamethoxazole has been reported,³⁷ but overall resistance to first-line, second-line, or third-line therapy is rare. Cephalosporins, chloramphenicol, tetracycline, and quinolones are probably ineffective.

Treatment duration

Bacteremia should be treated for 2 weeks, meningitis for 3 weeks, endocarditis for 4 to 6 weeks, and rhombencephalitis or brain abscess for 6 weeks^{2,12} or until the patient has improved and the symptoms have resolved.

■ PREVENTION

Government agencies have launched programs to improve and ensure the safety of our foods.^{38,39} In addition, new methods of sterilizing foods are under investigation. Irradiation was recently approved as one such method. Unfortunately, this technology has a bad pub-



lic image, and it changes the appearance and taste of food. If these obstacles can be overcome, irradiation may hold promise.

Persons at high risk for listeriosis must exercise caution in their dietary choices, and common sense must prevail for the rest of the population. TABLE 3 provides some useful guidelines. Persons at risk should avoid raw vegetables unless they are known to be carefully washed. All dairy products should be pasteur-

ized, and soft cheeses should be avoided. Meats should be cooked until the internal temperature is 160 to 180°F, ie, medium-well to well done. Refrigerated leftover foods need to be thoroughly reheated to destroy *L. monocytogenes*, which can proliferate at low temperatures. Although deli foods usually carry a low risk for listerial infection, they either should be well heated before eating or should be avoided.



REFERENCES

- Nyfeldt A. Etiologie de la mononucleose infectieuse. CR Soc Biol 1929; 101:590-592.
- Nieman R, Lorber B. Listeriosis in adults: a changing pattern. Report of eight cases and review of the literature. 1968-1978. Rev Infect Dis 1980; 2:207-227.
- Ciesielski C, Hightower A, Parsons S, Broome C. Listeriosis in the United States: 1980-1982. Arch Intern Med 1988; 148:1416-1419.
- Linnan M, Mascola L, Lou X, et al. Epidemic listeriosis associated with Mexican-style cheese. N Engl J Med 1988; 319:823-828.
- Schlech W, Lavigne P, Bortolussi R, et al. Epidemic listeriosis—evidence for transmission by food. N Engl J Med 1983; 308:203-206.
- Pinner R, Schuchat A, Swaminathan B, et al. Role of foods in sporadic listeriosis. II. Microbiologic and epidemiologic investigation. JAMA 1992; 267:2046-2050.
- Schuchat A, Deaver K, Wenger J, et al. Role of foods in sporadic listeriosis. I. Case-control study of dietary risk factors. JAMA 1992; 267:2041-2045.
- Fleming D, Cochi S, MacDonald L, et al. Pasteurized milk as a vehicle of infection in an outbreak of listeriosis. N Engl J Med 1985; 312:404-407.
- Dalton C, Austin C, Sobel J, et al. An outbreak of gastroenteritis and fever due to *Listeria monocytogenes* in milk. N Engl J Med 1997; 336:100-105.
- Farber J, Peterkin P. *Listeria monocytogenes*, a foodborne pathogen. Microbiol Rev 1991; 55:476-511.
- WHO Working Group. Foodborne listeriosis. Bull World Health Org 1988; 66:421-428.
- Lorber B. Listeriosis. Clin Infect Dis 1997; 24:1-11.
- Altekruse S, Swerdlow D. The changing epidemiology of foodborne diseases. Am J Med Sci 1996; 311:23-29.
- Gellin B, Broome C. Listeriosis. JAMA 1989; 261:1313-1320.
- Southwick F, Purich D. Intracellular pathogenesis of listeriosis. N Engl J Med 1996; 334:770-776.
- Sword C. Mechanisms of pathogenesis in *Listeria monocytogenes* infection. I. Influence of iron. J Bacteriol 1966; 92:536-542.
- Hearmon C, Ghosh S. *Listeria monocytogenes* meningitis in previously healthy adults. Postgrad Med J 1989; 65:74-78.
- Anaissie E, Kontoyiannis P, O'Brien S, et al. Infections in patients with chronic lymphocytic leukemia treated with fludarabine. Ann Intern Med 1998; 129:559-566.
- Anaissie E, Kontoyiannis P, Kantarjian H, et al. Listeriosis in patients with chronic lymphocytic leukemia who were treated with fludarabine and prednisone. Ann Intern Med 1992; 117:466-469.
- Schroter G, Weil R. *Listeria monocytogenes* infection after renal transplantation. Arch Intern Med 1977; 137:1395-1399.
- Jurado R, Farley M, Pereira E, et al. Increased risk of meningitis and bacteremia due to *Listeria monocytogenes* in patients with human immunodeficiency virus infection. Clin Infect Dis 1993; 17:224-227.
- Altekruse S, Cohen M, Swerdlow D. Emerging foodborne diseases. Emerg Infect Dis 1997; 3:285-293.
- Skogberg K, Syrjanen J, Jähkola M, et al. Clinical presentation and outcome of listeriosis in patients with and without immunosuppressive therapy. Clin Infect Dis 1992; 14:815-821.
- Schwartz B, Hexter D, Broome C, et al. Investigation of an outbreak of listeriosis: new hypotheses for the etiology of epidemic *Listeria monocytogenes* infections. J Infect Dis 1989; 159:680-685.
- Riedo F, Pinner R, de Lourdes Tosca M, et al. A point-source foodborne listeriosis outbreak: documented incubation period and possible mild illness. J Infect Dis 1994; 170:693-696.
- Visintine K, Oleske J, Nahmias A. *Listeria monocytogenes* infection in infants and children. Am J Dis Child 1977; 131:393-397.
- Durand M, Calderwood S, Weber D, et al. Acute bacterial meningitis in adults. N Engl J Med 1993; 328:21-28.
- Boucher M, Yonekura L. *Listeria* meningitis during pregnancy. Am J Perinatol 1984; 1:312-318.
- Armstrong R, Fung P. Brainstem encephalitis (rhombencephalitis) due to *Listeria monocytogenes*: Case report and review. Clin Infect Dis 1993; 16:689-702.
- Uldry P, Kuntzer T, Bogousslavsky J, et al. Early symptoms and outcome of *Listeria monocytogenes* rhombencephalitis: 14 adult cases. Neurol 1993; 240:235-242.
- Gellin B, Broome C, Bibb W, et al. The epidemiology of listeriosis in the United States—1986. Am J Epidemiol 1991; 133:392-401.
- Bassan R. Bacterial endocarditis produced by *Listeria monocytogenes*. Am J Clin Pathol 1975; 63:522-527.
- Sivalingam J, Martin P, Fraimow H, et al. *Listeria monocytogenes* peritonitis: case report and literature review. Am J Gastroenterol 1992; 87:1839-1845.
- Stamm A, Smith S, Kirklin J, McGiffin D. Listerial myocarditis in cardiac transplantation. Rev Infect Dis 1990; 12:820-823.
- Spitzer P, Hammer S, Karchmer A. Treatment of *Listeria monocytogenes* infection with trimethoprim-sulfamethoxazole: case report and review of the literature. Rev Infect Dis 1986; 8:427-430.
- Jones E, MacGowan A. Antimicrobial chemotherapy of human infection due to *Listeria monocytogenes*. Eur J Clin Microbiol Infect Dis 1995; 14:165-175.
- Charpentier E, Gerbaud G, Jacquet C, Rocourt J, Corvalin P. Incidence of antibiotic resistance in *Listeria* species. J Infect Dis 1995; 172:277-281.
- Tauxe RV. New approaches to surveillance and control of emerging foodborne infectious diseases. Emerg Infect Dis 1998; 4:455-456.
- Yang S. FoodNet and Enter-net: emerging surveillance programs for foodborne diseases. Emerg Infect Dis 1998; 4:457-458.

ADDRESS: Alan J. Taega, MD, Department of Infectious Disease, S32, The Cleveland Clinic Foundation, 9500 Euclid Avenue, Cleveland, OH 44195.