



Q: Can calcium and vitamin D supplementation adequately treat most patients with osteoporosis?

CHAD DEAL, MD

Department of Rheumatic and Immunologic Disease,
Cleveland Clinic

A: ALTHOUGH CALCIUM AND VITAMIN D play an important role, they are not sufficient when used alone. Rather, they should be used in combination with an antiresorptive agent such as estrogen, raloxifene, alendronate, risedronate, or calcitonin.

■ CATEGORIES OF BONE LOSS

The World Health Organization¹ has defined three levels of low bone mass; the definitions are based on T scores, which are standard deviation units below peak bone mass:

- **Osteopenia**—a T score between -1 and -2.5
- **Osteoporosis**—a T score less than -2.5
- **Severe osteoporosis**—a T score less than -2.5 with a fracture.

The National Osteoporosis Foundation² recommends treatment with an antiresorptive agent in patients with a T score less than -2.0 ; patients with T scores less than -1.5 to -2.0 should also be treated if they have any of the following risk factors:

- Family history of osteoporosis
- Previous fracture
- Current tobacco use
- Body weight less than 127 pounds.

Many other risk factors such as steroid use are also important and should be considered.

■ STUDIES OF CALCIUM AND VITAMIN D

In almost all randomized controlled trials of antiresorptive agents, the control groups took calcium and vitamin D supplements. In these trials, patients who took alendronate, risedronate, or raloxifene³⁻⁵ had significantly fewer fractures than those who took calcium and vitamin D alone. Thus, we conclude that supplements alone are not adequate.

Still, calcium and vitamin D are important in treating age-related bone loss, as they reduce the rate of bone loss and possibly reduce fracture risk.

Heaney^{6,7} reviewed 43 studies of calcium published between 1988 and 1993. Although 16 studies showed that calcium had no effect on bone loss, 16 of the 19 placebo-controlled studies in which calcium intake was controlled did show that the mineral prevented or slowed bone loss. In the 12 studies that excluded women who were within 5 years of menopause—a period when estrogen deficiency overwhelms the effect of calcium supplementation⁸—all showed that calcium had a significantly beneficial effect.

Several well-controlled studies showed that calcium and vitamin D can reduce hip and nonvertebral fractures in elderly patients. Chapuy et al⁹ gave healthy elderly women either 800 IU of vitamin D₃ and 1.2 g of elemental calcium or a double placebo every day for 18 months. The number of hip fractures was 43% lower and the number of nonvertebral fractures was 32% lower in the supplement group compared to the placebo group. In addition, the hip bone density in the supplement group increased by 2.7%.

Dawson-Hughes et al¹⁰ gave supplements to men and women age 65 and older who had low calcium intakes (400–650 mg/day); the supplements contained calcium 500 mg and vitamin D₃ 700 IU. At 36 months, bone density was higher and the nonvertebral fracture rate was 50% lower in the supplement group.

In a 2-year study of women who were already calcium-replete (ie, who had an average daily calcium intake of 750 mg) and who had reached menopause at least 3 years before enrollment, Reid et al¹¹ found that total body bone loss was 43% lower in those given a calcium supplement of 1,000 mg/day than in those given a placebo. In a follow-up study, 86

By itself,
supplementation
is not enough



of the original 122 patients were followed for another 2 years (4 years total); the amount of bone loss and the number of fractures was lower in those who continued receiving calcium supplements than in those who continued receiving a placebo.¹²

In a 3-year study by Tilyard et al,¹³ women with vertebral fractures at enrollment who were treated with calcitriol (1,25 dihydroxy-vitamin D₃ 0.25 µg twice a day) had a lower rate of new vertebral fractures during the second and third year than did women treated with calcium 1,000 mg/day.

■ MORE RESEARCH IS NEEDED

In the treatment of age-related bone loss, numerous questions remain unanswered. Are the effects of calcium supplements more (or only) significant in patients with calcium-deficient diets? Research suggests that patients with calcium intakes lower than 500 mg/day are more likely to respond to calcium supplementation. In addition, is vitamin D therapy more effective in those with low calcium intake or low vitamin D levels? And is calcitriol more effective than vitamin D₃?

Following are a few points that can be stated with some certainty:

- Most postmenopausal women have a deficient intake of dietary calcium. In addition, serum vitamin D levels are low in at least 20% of women who live in North America and Northern Europe.¹⁴
- Laboratories consider the lower limit of normal for blood levels of 25-hydroxyvitamin D to be 9 ng/mL. Physiologically, however, values at the low end of the "normal" range are too low because parathyroid hormone levels rise as vitamin D levels fall below 20 to 25 ng/mL, and elevated parathyroid hormone increases bone turnover. Therefore, the RDA for vitamin D (400 IU/day) is probably too low and should be increased to 600 to 800 IU/day.¹⁴
- A postmenopausal woman who is not on estrogen therapy should take 1,500 mg of calcium a day; those on estrogen therapy should take 1,000 mg a day.¹⁵
- Calcium and vitamin D have a substantial effect on age-related bone loss in elderly women. It is likely that supplementation could prevent a proportion of bone fractures.

■ REFERENCES

1. Lindsay R, Meunier PJ. Osteoporosis: review of the evidence for prevention, diagnosis, and treatment and cost-effectiveness analysis. *Osteoporosis Int* 1998; 8(Suppl 14):S1-S588.
2. National Osteoporosis Foundation. Pocket guide to the prevention and treatment of osteoporosis. Washington DC: National Osteoporosis Foundation, 1998:8.
3. Black DM, Cummings SR, Karpf DB, et al. Randomized trial of effect of alendronate on the risk of fracture in women with existing vertebral fractures. *Lancet* 1996; 348:1535-1541.
4. Harris ST, Watts NB, Genant HR, et al. Effects of risedronate treatment on vertebral and nonvertebral fractures in women with postmenopausal osteoporosis: a randomized controlled trial. *JAMA* 1999; 282:1344-1352.
5. Ettinger B, Black DM, Mitlak BH, et al. Reduction of vertebral fracture risk in postmenopausal women with osteoporosis treated with raloxifene: results from a 3-year randomized clinical trial. *JAMA* 1999; 282:637-645.
6. Heaney RP. Thinking straight about calcium. *N Engl J Med* 1993; 328:503-505.
7. Heaney RP. Nutritional factors in osteoporosis. *Annu Rev Nutr* 1993; 13:287-316.
8. Riis B, Thomsen K, Christiansen C. Does calcium supplementation prevent post menopausal bone loss? A double-blind, controlled clinical study. *N Engl J Med* 1987; 316:173-177.
9. Chapuy MC, Arlot ME, Duboeuf F, et al. Vitamin D3 and calcium to prevent hip fractures in elderly women. *N Engl J Med* 1992; 327:1637-1642.
10. Dawson-Hughes B, Harris SS, Krall EA, et al. Effect of calcium and vitamin D supplementation on bone density in men and women 65 years of age or older. *N Engl J Med* 1997; 337:670-676.
11. Reid IR, Ames RW, Evans MC, et al. Effect of calcium supplementation on bone loss in postmenopausal women. *N Engl J Med* 1993; 328:460-464.
12. Reid IR, Ames RW, Evans MC, et al. Long-term effects of calcium supplementation on bone loss and fractures in postmenopausal women: a randomized controlled trial. *Am J Med* 1995; 98:331-335.
13. Tilyard MW, Spears GFS, Thomson J, et al. Treatment of postmenopausal osteoporosis with calcitriol or calcium. *N Engl J Med* 1992; 326:357-362.
14. McKenna MJ, Freaney R. Secondary hyperparathyroidism in the elderly: means to defining hypovitaminosis D. *Osteoporosis Int* 1998; 8(Suppl 2):S3-S6.
15. Consensus conference: osteoporosis. *JAMA* 1984; 252:799-802.

ADDRESS: Chad Deal, MD, Department of Rheumatic and Immunologic Disease, A50, The Cleveland Clinic Foundation, 9500 Euclid Avenue, Cleveland, OH 44195; e-mail dealc@ccf.org.

CME ANSWERS



Answers to the CREDIT TEST on page 775 of this issue

1 E 2 D 3 D 4 E 5 D 6 E 7 C 8 B 9 A 10 C
11 B 12 A