

**KEVIN R. FLAHERTY, MD**Division of Pulmonary and Critical Care Medicine,
University of Michigan Health System, Ann Arbor**FERNANDO J. MARTINEZ, MD**Division of Pulmonary and Critical Care Medicine,
University of Michigan Health System, Ann Arbor

Diagnosing interstitial lung disease: A practical approach to a difficult problem

■ ABSTRACT

Interstitial lung disease has a variety of causes: environmental, infectious, autoimmune, and drug-related. Accurate diagnosis is essential because the prognosis and treatment of the disease varies widely depending on the cause. However, the respiratory symptoms and pulmonary radiographic picture of these various causes of interstitial lung disease are often similar, making the diagnosis of its cause confusing and frustrating. The practical, algorithmic approach to diagnosis outlined here identifies key diagnostic clues in the patient's history, physical exam, and radiographic findings.

■ KEY POINTS

Interstitial lung disease is characterized by cough, progressive dyspnea, restrictive pulmonary physiology, and abnormalities on chest radiography.

A standardized, logical evaluation yields a diagnosis in the majority of patients with interstitial lung disease.

Although laboratory studies, radiography, and bronchoscopic procedures may provide useful information, they often do not provide the diagnosis, and surgical lung biopsy remains the standard for definitive diagnosis.

RECOGNIZING INTERSTITIAL lung disease (ILD) and identifying its cause can be difficult for several reasons:

- Many diseases can cause the cough, progressive dyspnea, and pulmonary fibrosis that are the chief features of ILD
- Symptoms are often mild and slowly progressive
- Patients wait long before reporting symptoms
- No underlying cause may be found (idiopathic pulmonary fibrosis).¹

The prognosis and treatments for interstitial lung disease vary widely depending on the cause, so accurate diagnosis is essential. Whenever ILD is suspected, a disciplined evaluation using an algorithmic approach is the key to diagnosis of the causative disease, if there is one. In this article we outline such a practical approach and review key historical, physical, and radiographic features that help to narrow the differential diagnosis. Treatment options are discussed.

■ THE CAUSES

The pathogenesis of ILD is thought to center around an injury to the lung—environmental, infectious, autoimmune, or drug-induced—followed by an attempt to heal the injury.² Whether this injury represents an ongoing insult, a series of multiple events, or an abnormal response to a single event that is no longer present is unknown. Whatever the scenario, it is believed that the attempt to control the injury eventually leads to inflammation and fibrosis, with subsequent destruction of lung

TABLE 1

Causes and categories of interstitial lung disease

Inhaled agents

Inorganic: asbestos, beryllium, silica

Organic: animal and bird antigens, farm antigens

Drug-induced

Antiarrhythmics

Antibiotics

Antidepressants

Anti-inflammatory agents

Chemotherapeutic agents

Oxygen

Radiation

Connective tissue disease

Ankylosing spondylitis

Behçet syndrome

Mixed connective tissue disease

Polymyositis/dermatomyositis

Rheumatoid arthritis

Scleroderma

Sjögren syndrome (primary)

Systemic lupus erythematosus

Infectious

Atypical pneumonias

Pneumocystis carinii pneumonia

Tuberculosis

Other

Acute interstitial pneumonia

Bronchiolitis obliterans organizing pneumonia

Desquamative interstitial pneumonia

Eosinophilic granuloma

Idiopathic

Lymphangioleiomyomatosis

Lymphocytic interstitial pneumonia

Nonspecific interstitial pneumonia

Respiratory bronchiolitis with interstitial lung disease

Sarcoidosis

Usual interstitial pneumonia

Malignant

Bronchoalveolar cell carcinoma

Lymphangitic carcinomatosis

ADAPTED FROM KING TE JR. APPROACH TO THE PATIENT WITH INTERSTITIAL LUNG DISEASE. IN: B. ROSE, EDITOR. UP TO DATE, [ON CD-ROM] WELLESLEY, MA, 2000

Patients often realize the true duration of cough and dyspnea only in retrospect

architecture and disruption of pulmonary function. An abbreviated list of causes of ILD is presented in TABLE 1.

FEATURES OF INTERSTITIAL LUNG DISEASE

Patients with ILD typically present with cough and progressive dyspnea, but these symptoms are often subtle, nonspecific, and slowly progressive; it is common for patients to realize the true duration of symptoms only in retrospect. Therefore, the physician needs to maintain a suspicion of ILD to facilitate diagnostic testing.

Patients can also present with abnormalities on chest radiography, with a systemic illness that includes pulmonary symptoms such as cough and dyspnea, or with abnormalities on pulmonary function testing, such as a restrictive ventilatory defect or a gas exchange abnormality.

Other chest symptoms are unusual but may provide clues to the cause of ILD. For example, hemoptysis can be seen in patients with alveolar hemorrhage syndromes, pulmonary vascular disease, lymphangioleiomyomatosis, tuberous sclerosis, and chronic mitral valve disease. Acute chest pain may represent pleurisy in patients with collagen vascular illness or pneumothorax in patients with lymphangioleiomyomatosis, tuberous sclerosis, or eosinophilic granuloma.

Symptom onset

The time over which symptoms develop can be important to the differential diagnosis. Acute onset is often seen in atypical infections, eosinophilic pneumonia, pulmonary hemorrhage, Wegener granulomatosis, acute interstitial pneumonia, and initial hypersensitivity reactions. An insidious onset is seen in patients with idiopathic pulmonary fibrosis, silicosis, asbestosis, long-standing hypersensitivity pneumonitis, and drug-induced lung diseases. Because dyspnea and cough are often subtle, nonspecific, and slowly progressive, patients often realize the true duration of symptoms only in retrospect.

Frequently, patients with ILD first present with abnormalities on a chest radiograph or pulmonary function testing that indicates a restrictive ventilatory defect or abnormal gas exchange. Keeping ILD in mind facilitates appropriate diagnostic testing.

Clues from the medical history

A careful medical history is an essential part of the evaluation of patients with suspected ILD. Important features include the patient's age and sex: ILD caused by sarcoidosis, eosinophilic granuloma, familial idiopathic pulmonary fibrosis, and Gaucher disease is more common in younger patients, while other conditions are more likely to cause ILD in older patients. ILD caused by lymphangiomyomatosis and pulmonary involvement in tuberous sclerosis is seen exclusively in premenopausal women.

The medical history can also uncover a previous diagnosis of collagen vascular disease, a common cause of ILD. Evaluation of risk for human immunodeficiency virus (HIV) infection is also important because patients with HIV can have ILD due to infection, neoplasm, or other causes related to immune deficiency.³

History of occupational and environmental exposures

A thorough history of occupational and environmental exposures aids the diagnosis and may also help to direct therapy. The range of exposures associated with the development of ILD is vast and includes avian, animal, and fish proteins, fungal spores, asbestos, silica, cobalt, beryllium, aluminum, isocyanates, and copper sulfate.³ Because the latency period between disease onset and the development of symptoms can be years, a detailed assessment of all previous occupations and potential environmental exposures is also necessary.

Information about the patient's home environment is also important. A recent move or home remodeling can expose people to new antigens, and exposure-related conditions such as hypersensitivity pneumonitis can cause ILD. Similarly, information about pets, especially birds, narrows the differential diagnosis and directs therapy, such as removal or avoidance of the offending antigen.

The occupations of family members should also be ascertained, as contaminated clothing can also be a source of exposure.⁴⁻⁶

Current and previous medications

A detailed account of current and previous medications should be obtained: many prescription and over-the-counter drugs have

been associated with the development of ILD (TABLE 1). The history should also include the use of recreational drugs such as cocaine.^{7,8}

Smoking history

Several forms of ILD are seen almost exclusively in smokers: eg, respiratory bronchiolitis/ILD, desquamative interstitial pneumonia, and eosinophilic granuloma.⁹ Cigarette smoking has also been implicated in the development of idiopathic pulmonary fibrosis.¹⁰⁻¹³

Paradoxically, smoking may reduce the incidence of sarcoidosis and hypersensitivity pneumonitis. However, when hypersensitivity pneumonitis is present in smokers, it follows a more chronic course with a worse clinical outcome.¹⁴

Family medical history

Familial types of pulmonary fibrosis have been reported,^{15,16} so obtaining a family medical history is advised. This may be helpful not only in narrowing the differential diagnosis but also in possibly identifying other family members with an earlier stage of disease.

FINDINGS IN THE PHYSICAL EXAMINATION

Physical findings related to ILD are nonspecific. The characteristic finding is dry bibasilar crackles, although inspiratory high-pitched rhonchi ("squeaks") can be heard with bronchiolitis. Clubbing (most common in idiopathic pulmonary fibrosis) and signs of right heart failure can also be seen in patients with advanced disease.

The physical examination is particularly helpful when it uncovers signs of an underlying connective tissue disorder. The presence of a rash (malar, heliotropic, vasculitic, or due to erythema nodosum), Raynaud phenomenon, joint deformity, synovial swelling, or muscle weakness should prompt a more complete evaluation for an underlying rheumatologic disorder.

LABORATORY TESTING

Laboratory testing can be useful in the diagnosis and management of patients with ILD. A list of various laboratory findings and disease associations is shown in TABLE 2.

Ask about occupational, environmental exposures—past and present

TABLE 2

Laboratory findings in interstitial lung disease

ABNORMALITY	ASSOCIATED CONDITION
Elevated angiotensin-converting enzyme (serum)	Sarcoidosis, hypersensitivity silicosis, Gaucher disease
Elevated antibasement membrane antibody	Goodpasture syndrome
Elevated antineutrophil cytoplasmic antibody	Wegener granulomatosis, Churg-Strauss syndrome, microscopic polyangiitis
Eosinophilia	Eosinophilic pneumonia, sarcoidosis, systemic vasculitis, drug-induced (sulfa, methotrexate)
Hemolytic anemia	Connective tissue disease, sarcoidosis, lymphoma, drug-induced
Hypergammaglobulinemia	Connective tissue disease, sarcoidosis, systemic vasculitis, lymphocytic interstitial pneumonia, lymphoma
Hypogammaglobulinemia	Lymphocytic interstitial pneumonitis
Immune complexes	Idiopathic pulmonary fibrosis, lymphocytic interstitial pneumonitis, systemic vasculitis, connective tissue disease, eosinophilic granuloma
Lactate dehydrogenase elevation	Alveolar proteinosis, idiopathic pulmonary fibrosis
Leukopenia	Sarcoidosis, connective tissue disease, lymphoma, drug-induced
Lymphocyte transformation test	Chronic beryllium disease, aluminum potroom positive worker's disease, gold-induced pneumonitis
Normocytic anemia	Diffuse alveolar hemorrhage syndromes, connective tissue disease, lymphangitic carcinomatosis
Serum precipitating antibodies	Hypersensitivity pneumonitis
Thrombocytopenia	Sarcoidosis, connective tissue disease, drug-induced, Gaucher disease
Urinary sediment abnormalities	Connective tissue disease, systemic vasculitis, drug-induced

ADAPTED FROM SCHWARZ MI, KING JR TE, CHERNIAK RM. GENERAL PRINCIPLES AND DIAGNOSTIC APPROACH TO THE INTERSTITIAL LUNGS DISEASES. IN: J. MURRAY AND J. NADEL, EDITORS. TEXTBOOK OF RESPIRATORY MEDICINE, 2ND EDITION. PHILADELPHIA: WB SAUNDERS, 1994: 1803-1826.

The time over which symptoms develop is an important diagnostic clue

A minimum panel of initial laboratory tests includes a complete blood count with differential, electrolytes, renal function studies, liver function studies, antinuclear antibodies, rheumatoid factor, and urinalysis.³ If chronic beryllium disease is suspected, a lymphocyte transformation test should be performed.

■ TESTING FOR ABNORMAL PULMONARY PHYSIOLOGY

Physiologically, a restrictive ventilatory defect that reflects reduced lung volumes is common

in patients with ILD.¹⁷ The vital capacity is typically reduced to a greater extent than the total lung capacity and functional residual capacity.¹⁷

However, these abnormalities are not unique to ILD or idiopathic pulmonary fibrosis and may not occur in all patients with pulmonary fibrosis. In two studies,^{18,19} smokers and ex-smokers with interstitial pulmonary fibrosis had preserved lung volumes. Doherty et al¹⁹ compared two groups of patients with cryptogenic fibrosing alveolitis—those with relatively preserved lung volumes and those with

TABLE 3

Physiologic features that may suggest specific diagnoses

PHYSIOLOGIC FEATURE	POSSIBLE DIAGNOSES
Airflow obstruction	Sarcoidosis Lymphangioleiomyomatosis Tuberous sclerosis Interstitial lung disease with superimposed chronic obstructive pulmonary disease
Isolated decrease in diffusing capacity of lung for carbon monoxide	Interstitial lung disease with superimposed emphysema Pulmonary vascular disease Eosinophilic granuloma Lymphangioleiomyomatosis

the typical pulmonary restriction—and found that both groups had the same degree of pulmonary fibrosis, but that 86% of patients with preserved volumes had concomitant emphysema vs 19% of those with pulmonary restriction.

In ILD, the diffusing capacity of carbon monoxide is typically reduced to a greater extent than the lung volume at which it is measured,¹⁷ and it appears to be more decreased in idiopathic pulmonary fibrosis than it is in ILD due, for example, to sarcoidosis.^{20,21}

Arterial blood gas abnormalities characteristic of idiopathic pulmonary fibrosis include resting hypoxemia and an increase in the alveolar-arterial oxygen pressure difference.¹⁷ Gas exchange abnormalities are more evident during exercise, with hypoxemia being highly prevalent.^{22–25} TABLE 3 lists physiologic features and the diagnoses they suggest.

Pulmonary function testing

Pulmonary function testing is a helpful early diagnostic tool in patients with appropriate symptoms, perhaps more so than chest radiography and high-resolution computed tomography (CT).

In one study,²⁶ 44 dyspneic patients with biopsy-proven ILD had normal chest radiographs but several pulmonary function test abnormalities: the diffusing capacity of lung for carbon monoxide was decreased in 73% of patients, the vital capacity was low in 57%, and the total lung capacity was low in 16%.²⁶ In another study,²⁷ three of 25 dyspneic patients with biopsy-confirmed ILD had nor-

mal high-resolution CT scans, despite abnormal pulmonary function tests.²⁷ These data suggest that abnormal pulmonary function tests in patients with symptoms such as breathlessness and cough should prompt further evaluation for ILD.

Unfortunately, pulmonary function tests may be normal despite histologic and radiographic evidence of ILD, even in patients with idiopathic pulmonary fibrosis.^{28,29} For example, in one study³⁰ two patients with biopsy-proven idiopathic pulmonary fibrosis had a diffusing capacity of lung for carbon monoxide greater than 70% of predicted, whereas the alveolar-arterial oxygen pressure difference was increased during both rest and exercise.³⁰ Therefore, normal pulmonary function test results cannot be assumed to exclude ILD when a patient has clinical or radiographic abnormalities that suggest it, although this is unusual.

Chest radiography and high-resolution computed tomography

Radiographic studies are usually abnormal in patients with ILD, although chest radiographs and high-resolution CT scans are normal in 10%.^{26,27} In general, chest radiographs and high-resolution CT scans show a mixture of interstitial and alveolar infiltrates. The techniques are most helpful, however, when they show characteristics that are “diagnostic” of a specific form of ILD (TABLES 4 AND 5).

The chief drawback with chest radiography in patients with ILD is a specificity of only 50%.³

Surgical lung biopsy is still the gold standard in ILD diagnosis

High-resolution CT, on the other hand, has dramatically improved the diagnostic evaluation of patients with ILD. The technique allows a detailed evaluation of the lung parenchyma by using 1-mm to 2-mm slices reconstructed with an algorithm that maximizes spatial resolution.^{31,32} Several studies³³ have confirmed that abnormalities not visible with chest radiography can be seen with high-resolution CT.³³ Furthermore, observer variability is less of a factor with high-resolution CT than with chest radiography, and a confident diagnosis is more likely with high-resolution CT.

High-resolution CT is particularly likely to be diagnostic in patients with idiopathic pulmonary fibrosis, lymphangitic carcinoma, sarcoidosis, silicosis, subacute hypersensitivity pneumonitis, and pulmonary alveolar proteinosis.³ **FIGURE 1** shows high-resolution CT scans of a patient with idiopathic pulmonary fibrosis and the characteristic findings of bibasilar interstitial and intralobular reticular opacities, irregular interlobular septal thickening, and subpleural honeycombing in the lower lobes in the absence of ground-glass densities and pleural abnormalities.³⁴

The diagnostic sensitivity of high-resolution CT in patients with idiopathic pulmonary fibrosis is estimated at 84%, compared with a sensitivity of 73% for chest radiography.³³ When experienced radiologists interpreting high-resolution CT are confident of the diagnosis of idiopathic pulmonary fibrosis using these features, they are usually correct.³³ The diagnostic sensitivity and specificity for many other forms of ILD are less well defined and depend on the experience of the thoracic radiologist interpreting the study.³³

Bronchoscopic procedures

In the diagnosis of ILD, the role of bronchoscopy, including bronchoalveolar lavage and transbronchial biopsy, is unclear.

Bronchoalveolar lavage is easy to perform, is associated with little risk, and can be diagnostic in certain occupational exposures to inorganic dusts, malignancy, hematological disease, drug-induced lung disease, and pulmonary alveolar proteinosis.³ A recent discriminant diagnostic model generated from bronchoalveolar lavage counts in a population

TABLE 4

Parenchymal abnormalities on chest radiography: Associated diagnoses

ABNORMALITY	POSSIBLE ASSOCIATED DIAGNOSES
Micronodules (< 1.5 mm)	Alveolar microlithiasis Talc granulomatosis Early pneumoconiosis Thesauriosis (hair spray inhalation) Early pneumocystis Pulmonary alveolar proteinosis
Nodules (up to 1 cm)	Infection Granulomatous inflammation Pneumoconiosis Cancer Pulmonary hemosiderosis
Linear opacities	Acute Congestive heart failure Atypical pneumonia Collagen vascular disease Chronic Idiopathic pulmonary fibrosis Lymphocytic interstitial pneumonitis Collagen vascular disease Asbestosis Radiation pneumonia Drug-induced disease Radiation pneumonitis Lymphangitic tumor Lymphagioleiomyomatosis Sarcoidosis
Honeycombing	Idiopathic pulmonary fibrosis Eosinophilic granuloma Collagen vascular disease Pneumoconiosis End-stage hypersensitivity pneumonitis
Airspace opacity	Pulmonary hemorrhage Acute or chronic eosinophilic pneumonia Bronchiolitis obliterans with organizing pneumonia Pulmonary alveolar proteinosis

ADAPTED FROM ZIESCHE R, HOFBAUER E, WITTMANN K, PETKOV V, BLOCK L. A PRELIMINARY STUDY OF LONG-TERM TREATMENT WITH INTERFERON GAMMA-1B AND LOW-DOSE PREDNISONE IN PATIENTS WITH IDIOPATHIC PULMONARY FIBROSIS. *N ENGL J MED* 1999; 341:1264-1269, AND KING JR TE. APPROACH TO THE PATIENT WITH INTERSTITIAL LUNG DISEASE. IN: B. ROSE, EDITOR. UP TO DATE, WELLESLEY, MA. 2000

of patients with sarcoidosis, hypersensitivity pneumonitis, and idiopathic pulmonary fibrosis was applied to a second group of patients with a similar distribution of ILD: importantly, 94.5% of the patients were correctly diag-

nosed.³⁵ Still, the diagnostic role of lavage remains controversial, and definitive diagnosis usually requires biopsy.

Transbronchial biopsy can be performed at the same time as bronchoalveolar lavage but carries an additional risk of bleeding and pneumothorax. It can be very useful in the diagnosis of some patients with ILD, as enumerated in **TABLE 6**.³⁶ For example, the combination of bronchoalveolar lavage, transbronchial biopsy, and transbronchial mediastinal lymph node aspiration has proven highly sensitive in the diagnosis of sarcoidosis. These procedures were performed by Leonard and colleagues in 13 patients with suspected sarcoidosis and, in combination, provided a sensitivity of 100%.³⁷

Unfortunately, transbronchial biopsy is of limited value in the diagnosis of idiopathic pulmonary fibrosis due to the small amount of tissue obtained.

Surgical lung biopsy

Surgical lung biopsy remains the gold standard for the diagnosis of ILD.³⁸ In some cases, such as an elderly patient with advanced lung disease and high-resolution CT findings typical of idiopathic pulmonary fibrosis, the risks of surgery may outweigh the benefits in terms of the information it may provide regarding prognosis and treatment.

■ ALGORITHMIC APPROACH TO DIAGNOSIS

While the diagnosis of ILD remains a challenging problem, a systematic approach as outlined in this article is helpful. **FIGURE 2** presents our approach to evaluating patients with suspected ILD. This algorithm reflects published recommendations.^{3,34,39}

In brief, a careful history and physical examination are followed by selected laboratory testing, chest radiography, and pulmonary function testing. A look at the criteria in **TABLE 6** will help determine if a bronchoscopic procedure such as bronchoalveolar lavage or transbronchial biopsy will help the diagnosis. If not, or if the results of bronchoscopy are not diagnostic, then high-resolution CT assumes a pivotal role in further diagnostic efforts.

Typical historical features, physical findings, and typical features seen on high-resolu-

TABLE 5

Other abnormalities on chest radiography: Associated diagnoses*

ABNORMALITY	POSSIBLE ASSOCIATED DIAGNOSES
Normal or large lung volumes	Eosinophilic granuloma Sarcoidosis Lymphangioleiomyomatosis Cystic fibrosis
Upper zone distribution	Granulomatous inflammation Pneumoconiosis (except asbestosis) Ankylosing spondylitis Cystic fibrosis Infections (tuberculosis, fungal) Drug-induced disease
Lower zone distribution	Idiopathic pulmonary fibrosis Desquamative interstitial pneumonitis Nonspecific interstitial pneumonitis Drug-induced disease Asbestosis Scleroderma Rheumatoid arthritis
Pleural disease	Asbestosis Collagen vascular disease Lymphangitic tumor Lymphangioleiomyomatosis Drug-induced disease Sarcoidosis
Pneumothorax	Eosinophilic granuloma Lymphangioleiomyomatosis Tuberous sclerosis
Mediastinal adenopathy	Sarcoidosis Lymphoma Infection (tuberculosis, fungal) Berylliosis Amyloidosis Malignancy

*The chest radiograph may be normal in 10% of patients with interstitial lung disease with no known cause (ie, idiopathic pulmonary fibrosis) or due to collagen vascular disease, desquamative interstitial pneumonitis, hypersensitivity pneumonitis, or sarcoidosis.

tion CT (**FIGURE 1**) offer a high degree of certainty that the patient has idiopathic pulmonary fibrosis.

Evidence supporting this algorithmic approach

A prospective study³⁴ in a referral center with expertise in the evaluation of ILD recently val-




FIGURE 1. High-resolution computed tomography scans of the upper lung zones (left) and lower lung zones (right) in a 58-year-old man with progressive breathlessness over the previous 3 years. Both images demonstrate subpleural honeycombing, which is worse in the lower lung zones. Open lung biopsy confirmed the diagnosis of idiopathic pulmonary fibrosis (usual interstitial pneumonitis).

**In 10%
of cases,
the x-ray
and CT
are normal**

idated a similar diagnostic algorithm. Fifty-nine patients with new-onset ILD (29 with idiopathic pulmonary fibrosis, 13 with granulomatous inflammation, 17 with miscellaneous diseases) were evaluated according to specific clinical and radiographic criteria. A clinical diagnosis was made by an expert clinician after initial evaluation including high-resolution CT and bronchoscopic biopsy. Using surgical lung biopsy as the gold standard, the sensitivity of the algorithm for diagnosing ILD other than idiopathic pulmonary fibrosis as 88.8%, the specificity was 40%, the positive predictive value was 94%, and the negative predictive value was 25%. For idiopathic pulmonary fibrosis, the sensitivity was 62%, specificity 97%, positive predictive value 95%, and negative predictive value 73%.

Importantly, one third of patients with a final diagnosis of idiopathic pulmonary fibrosis would not have been diagnosed without surgical lung biopsy, despite evaluation by a highly experienced group of physicians. Clearly, a strong suspicion of idiopathic pulmonary fibrosis on clinical and radiographic grounds is quite valuable, although the reverse may not be true.

■ PROGNOSIS AND SURVIVAL IN PATIENTS WITH INTERSTITIAL LUNG DISEASE

The overall prognosis for patients with ILD varies widely according to the specific diagno-

sis. In general, patients with idiopathic pulmonary fibrosis have the worst prognosis, with median survival times ranging from 2 to 5 years after the onset of symptoms.³¹ The most common causes of death in these patients are respiratory failure, heart failure, bronchogenic carcinoma, ischemic heart disease, infection, and pulmonary embolism.⁴⁰

Patients with nonspecific interstitial pneumonia⁴¹ have a better prognosis, with an estimated median survival greater than 10 years.^{42–44} Patients with bronchiolitis obliterans organizing pneumonia (BOOP), desquamate interstitial pneumonia, and respiratory bronchiolitis-associated ILD have an excellent prognosis.⁴⁴

The prognosis for patients with granulomatous lung diseases such as sarcoidosis or hypersensitivity pneumonitis is generally good. A study of 479 patients with sarcoidosis identified only 13 (3%) deaths due to respiratory failure.⁴⁵ Similarly, a study from Finland⁴⁶ identified only 13 deaths attributable to hypersensitivity pneumonitis (farmer's lung) over a 10-year period. In both of these studies mortality was associated with the fibrotic changes on chest radiography.

Lung cancer risk


Progressive pulmonary fibrosis, however, is not the only cause of death in patients with ILD. Patients with silicosis, asbestosis, and idio-



Diagnostic algorithm for patients with suspected interstitial lung disease

Ascertain:

- Duration of symptoms
- Occupation
- Home environment, pets
- Connective tissue disease



Silicosis, asbestosis, and idiopathic pulmonary fibrosis increase lung cancer risk

FIGURE 2

ADAPTED FROM REFERENCES 3, 34, AND 39.

pathic pulmonary fibrosis have an increased risk of lung cancer compared to the general population. A study from Finland⁴⁷ using a cancer registry found the standardized incidence ratio for all cancers to be 1.5 (95% CI, 1.0–2.1) for silicosis patients and 3.7 (95% CI, 2.8–5.0) for asbestosis patients.

Similarly, a recent population-based cohort study of 890 subjects with idiopathic pulmonary fibrosis in the United Kingdom⁴⁸ confirmed a markedly increased incidence of lung cancer (RR 7.31, 95% CI, 4.47–11.93) which persisted despite adjustment for smoking history. These patients could potentially benefit from cancer screening, although data addressing this issue are sparse.

TREATMENT OPTIONS IN INTERSTITIAL LUNG DISEASE

The treatment of ILD is evolving and controversial. For disorders such as sarcoidosis and hypersensitivity pneumonitis, steroid therapy is useful.^{3,49,50} Unfortunately, in some cases patients cannot tolerate steroid therapy and second-line immunosuppressive agents may be considered.

For pulmonary sarcoidosis, the recommended corticosteroid dosage is 20 to 40 mg of an intermediate-acting corticosteroid such as prednisone or an equivalent daily for 1 to 3 months.⁴⁹ The optimal duration of therapy is controversial, but the current recommenda-

TABLE 6

When is transbronchial lung biopsy useful in the workup of interstitial lung disease?

Very useful when the cause of interstitial lung disease is:

Sarcoidosis
Lymphangitic carcinomatosis
Alveolar proteinosis
Bronchoalveolar carcinoma
Eosinophilic pneumonia
Berylliosis

Occasionally useful when the cause of interstitial lung disease is:

Eosinophilic granuloma
Amyloidosis
Wegener granulomatosis
Pulmonary lymphoma
Hypersensitivity pneumonitis
Pulmonary capillaritis/vasculitis
Lymphocytic interstitial pneumonia
Bronchiolitis obliterans organizing pneumonia

Not useful when the cause of interstitial lung disease is:

Unknown (ie, idiopathic pulmonary fibrosis)
Any disease with underlying usual interstitial pneumonitis

ADAPTED FROM SCHWARZ MI. APPROACH TO THE UNDERSTANDING, DIAGNOSIS, AND MANAGEMENT OF INTERSTITIAL LUNG DISEASE. IN: M. SCHWARZ AND T. KING JR, EDITORS. INTERSTITIAL LUNG DISEASE. B. C. DECKER, INC., HAMILTON, ON. 1998: 3–30.

tion is for responders to be tapered to 10 mg every other day, with treatment continued for at least 1 year.

In the minority of patients not responding to or unable to tolerate steroids, alternative agents such as methotrexate, azathioprine, cyclophosphamide, and chlorambucil can be considered.

In hypersensitivity pneumonitis, the first therapy is antigen avoidance. Corticosteroids may be useful, especially for severely ill patients, although their impact on the long-term course of disease is uncertain.⁵¹ The recommended dose of corticosteroids is prednisone 0.5 mg to 1.0 mg per kg of ideal body

weight (maximum 60 mg/day) for 1 to 2 weeks, tapering over the next 2 to 4 weeks.⁵¹

In contrast, idiopathic pulmonary fibrosis responds to high-dose steroid treatment, but only in a minority of cases and with a high incidence of adverse reactions.⁵² Currently, the recommended therapy for idiopathic pulmonary fibrosis is combination therapy with low-dose prednisone plus azathioprine or cyclophosphamide.³¹

Unfortunately, there is little evidence that any of the therapies currently available improves survival or quality of life.^{31,53,54} However, several encouraging advances are on the horizon.^{55,56} Ziesche and colleagues⁵⁵ randomized 18 patients with IPF (definite or probably in 15) who had experienced progressive disease despite prednisone therapy to subcutaneous interferon gamma 200 mg three times per week plus low-dose oral prednisolone compared to placebo plus the same dose of oral steroid. Of importance, they demonstrated that pulmonary function improved in the interferon gamma group although it decreased in the placebo group. A large, multicenter study has recently been initiated to confirm these exciting, preliminary findings.

Authors' recommendation for treatment

Because ILD often responds poorly to therapy, and because of the potential for progressive deterioration over time, we believe all patients with ILD should have at least an initial evaluation by a pulmonologist before starting any form of therapy. The goal of this approach is to facilitate the enrollment of patients into multicenter treatment trials to determine which agents are most beneficial for the various forms of ILD. Such an approach can also facilitate early evaluation and consideration of other therapeutic interventions, such as lung transplantation.

REFERENCES

1. Coultas DB, Zumwalt RE, Black W, Sobonya RE. The epidemiology of interstitial lung diseases. *Am J Respir Crit Care Med* 1994; 150:967–972.
2. Toews GB. *Interstitial lung disease*. In: Goldman L, Bennett J, editors. *Cecil textbook of medicine*. Philadelphia: W. B. Saunders Company, 2000:409–419.
3. British Thoracic Society. The diagnosis, assessment, and treatment of diffuse parenchymal lung disease in adults: British Thoracic Society recommendations. *Thorax* 1999; 54:S1–S30.
4. Kilburn KH, Lillis R, Anderson HA, et al. Asbestos disease in family contacts of shipyard workers. *Am J Public Health* 1985; 75:615–617.
5. Knishkowsky B, Baker EL. Transmission of occupational disease to family contacts. *Am J Ind Med* 1986; 9:543–550.
6. Grandjean P, Bach E. Indirect exposures: the significance of bystanders at work and at home. *Am Ind Hyg Assoc J* 1986; 47:819–824.
7. O'Donnell AE, Mappin FG, Sebo TJ, Tazelaar H. Interstitial pneumonitis associated with "crack" cocaine abuse. *Chest* 1991; 100:1155–1157.
8. Bailey ME, Fraire AE, Greenberg SD, Barnard J, Cagle PT. Pulmonary histopathology in cocaine abusers. *Hum Pathol* 1994; 25:203–207.



9. Aubry MC, Wright JL, Myers JL. The pathology of smoking-related lung diseases. *Clin Chest Med* 2000; 21:11-135.
10. Carrington CB, Gaensler EA, Coutu RE, Fitzgerald MX, Gupta RG. Natural history and treated course of usual and desquamative interstitial pneumonia. *N Engl J Med* 1978; 298:801-809.
11. Hubbard R, Lewis S, Richards K, Johnston I, Britton J. Occupational exposure to metal or wood dust and etiology of cryptogenic fibrosing alveolitis. *Lancet* 1996; 374:284-289.
12. Iwai K, Mori T, Yamada N, Yamaguchi M, Hosoda Y. Idiopathic pulmonary fibrosis. Epidemiologic approaches to occupational exposure. *Am J Respir Crit Care Med* 1994; 150:670-675.
13. Baumgartner KB, Samet JM, Stidley CA, Colby TV, Waldron JA. Cigarette smoking: A risk factor for idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 1997; 155:242-248.
14. Murin S, Bilello KS, Matthay R. Other smoking-affected pulmonary diseases. *Clin Chest Med* 2000; 1:121-137.
15. Bitterman PB, Rennard SI, Keogh BA, Wewers MD, Adelberg S, Crystal RG. Familial idiopathic pulmonary fibrosis. Evidence of lung inflammation in unaffected family members. *N Engl J Med* 1986; 314:1343-1347.
16. Stinson JC, Tomkin GH. Familial cryptogenic fibrosing alveolitis: a case report. *Ir J Med Sci* 1992; 161:42-43.
17. O'Donnell DE. Physiology of interstitial lung disease. In Schwarz M, King T Jr, editors. *Interstitial lung disease*. Hamilton, Ontario: B. C. Decker Inc, 1998:51-70.
18. Cherniack RM, Colby TV, Flint A, et al. Correlation of structure and function in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 1995; 151:1180-1188.
19. Doherty MJ, Pearson MG, O'Grady EA, Pellegrini V, Calverley PMA. Cryptogenic fibrosing alveolitis with preserved lung volumes. *Thorax* 1997; 52:998-1002.
20. Dunn TL, Watters LC, Hendrix C, Cherniack RM, Schwarz MI, King TE. Gas exchange at a given degree of volume restriction is different in sarcoidosis and idiopathic pulmonary fibrosis. *Am J Med* 1988; 85:221-224.
21. Keogh BA, Lakatos E, Price D, Crystal RG. Importance of the lower respiratory tract in oxygen transfer. Exercise testing in patients with interstitial and destructive lung diseases. *Am Rev Respir Dis* 1984; 129 (Suppl):S76-S80.
22. Robertson HF. Clinical application of pulmonary function and exercise tests in the management of patients with interstitial lung disease. *Sem Respir Crit Care Med* 1994; 15:1-9.
23. Gaensler EA, Carrington CB, Coutu RE, Fitzgerald MX. Radiographic-physiologic-pathologic correlations in interstitial pneumonias. *Prog Respir Res* 1975; 8:223-241.
24. Austrian R, McClement JH, Renzetti AD, et al. Clinical and physiologic features of some types of pulmonary diseases with impairment of alveolar-capillary diffusion. *Am J Med* 1951; 11:667-685.
25. Fulmer JD, Roberts WC, von Gael ER, Crystal RG. Morphologic-physiologic correlates of the severity of fibrosis and degree of cellularity in idiopathic pulmonary fibrosis. *J Clin Invest* 1979; 63:65-76.
26. Epler GR, McCloud TC, Gaensler EA, Mikus JP, Carrington CB. Normal chest roentgenograms in chronic diffuse infiltrative lung disease. *N Engl J Med* 1978; 298:934-939.
27. Orens JB, Kazerooni EA, Martinez FJ, et al. The sensitivity of high-resolution CT in detecting idiopathic pulmonary fibrosis proved by open lung biopsy: A prospective study. *Chest* 1995; 108:109-115.
28. Crystal RG, Fulmer JD, Roberts WC, Moss ML, Line BR, Reynolds HY. Idiopathic pulmonary fibrosis. Clinical, histologic, radiographic, physiologic, scintigraphic, cytologic, and biochemical aspects. *Ann Int Med* 1976; 85:769-788.
29. Jezek V, Fucik J, Michaljanic A, Jiskova L. The prognostic significance of functional tests in cryptogenic fibrosing alveolitis. *Bull Physiolpathol Respir* 1980; 16:711-720.
30. Risk C, Epler GR, Gaensler EA. Exercise alveolar-arterial oxygen pressure difference in interstitial lung disease. *Chest* 1984; 85:69-74.
31. American Thoracic Society. Idiopathic pulmonary fibrosis: Diagnosis and treatment. International Consensus Statement. *Am J Respir Crit Care Med* 2000; 161:646-654.
32. Hansell DM. High-resolution computed tomography in the evaluation of fibrosing alveolitis. *Clin Chest Med* 1999; 20:739-760.
33. Wells A. Clinical usefulness of high-resolution computed tomography in cryptogenic fibrosing alveolitis. *Thorax* 1998; 53:1080-1087.
34. Raghu G, Mageto YN, Lockhart D, Schmidt RA, Wood DE, Godwin JD. The accuracy of the clinical diagnosis of new-onset idiopathic pulmonary fibrosis and other interstitial lung disease: A prospective study. *Chest* 1999; 116:1168-1174.
35. Drent M, von Nierop MAMF, Gerritsen FA, et al. A computer program using BALF analysis results as a diagnostic tool in interstitial lung diseases. *Am J Respir Crit Care Med* 1996; 153:736-741.
36. Schwarz MI. Approach to the understanding, diagnosis, and management of interstitial lung disease. In: Schwarz M, King T Jr, editors. *Interstitial lung disease*. Hamilton, Ontario: B. C. Decker Inc, 1998: 3-30.
37. Leonard C, Tormey VJ, O'Keane C, Burke CM. Bronchoscopic diagnosis of sarcoidosis. *Eur Respir J* 1997; 10:2722-2724.
38. Lynch III JP, Martinez FJ, Travis WD. Idiopathic pulmonary fibrosis: Is lung biopsy essential? *J Respir Dis* 2000; 21:197-214.
39. Raghu G. Interstitial lung disease: A diagnostic approach. Are CT scan and lung biopsy indicated for every patient? *Am J Respir Crit Care Med* 1995; 151:909-914.
40. Panos RJ, Mortenson RL, Niccoli SA, King TE. Clinical deterioration in patients with idiopathic pulmonary fibrosis: Causes and assessment. *Am J Med* 1990; 88:396-404.
41. Katzenstein A, Myers JL. Idiopathic pulmonary fibrosis. Clinical relevance of pathologic classification. *Am J Respir Crit Care Med* 1998; 157:1301-1315.
42. Travis WD, Matsui K, Moss J, Ferrans VJ. Idiopathic nonspecific interstitial pneumonia: Prognostic significance of cellular and fibrosing patterns. *Am J Surg Pathol* 2000; 24:19-33.
43. Nagai S, Kitaichi M, Itoh H, Nishimura K, Colby TV. Idiopathic nonspecific interstitial pneumonia/fibrosis: comparison with idiopathic pulmonary fibrosis and BOOP. *Eur Respir J* 1998; 12:1010-1019.
44. Bjoraker JA, Ryu JH, Edwin MK, et al. Prognostic significance of histopathologic subsets in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 1998; 157:199-203.
45. Baughman R, Winget D, Bowen EH, Lower EE. Predicting respiratory failure in sarcoidosis patients. *Sarcoidosis Vasc Diffuse Lung Dis* 1997; 14:154-158.
46. Kokkarinen J, Tukiainen H, Terho EO. Mortality due to farmer's lung in Finland. *Chest* 1994; 106:509-512.
47. Oksa P, Pukkala E, Karjalainen A, Ojajarvi A, Huuskonen MS. Cancer incidence and mortality among Finnish asbestos sprayers and in asbestosis and silicosis patients. *Am J Ind Med* 1997; 31:693-698.
48. Hubbard R, Venn A, Lewis S, Britton J. Lung cancer and cryptogenic fibrosing alveolitis. A population-based cohort study. *Am J Respir Crit Care Med* 2000; 161:5-8.
49. American Thoracic Society. Statement on sarcoidosis. *Am J Respir Crit Care Med* 1999; 160:736-755.
50. Paramothayan NS, Jones PW. Corticosteroids for pulmonary sarcoidosis (Cochrane Review). In: *The Cochrane Library*, Issue 2 2000. Oxford: Update Software.
51. Weill D, Rose C. Treatment and prognosis of hypersensitivity pneumonitis (extrinsic allergic alveolitis). Wellesley, MA.
52. Flaherty KR, Toews GB, Lynch III JP, et al. Steroids in idiopathic pulmonary fibrosis: Prospective assessment of adverse reactions, response to therapy, and survival. *Am J Med* (in press).
53. Zisman DA, Lynch III JP, Toews GB, Kazerooni EA, Flint A, Martinez FJ. Cyclophosphamide in the treatment of idiopathic pulmonary fibrosis. A prospective study in patients who failed to respond to corticosteroids. *Chest* 2000; 117:1619-1626.
54. Douglas WW, Ryu JH, Schroeder DR. Idiopathic pulmonary fibrosis. Impact of oxygen and colchicine, prednisone, or no therapy on survival. *Am J Respir Crit Care Med* 2000; 161:1172-1178.
55. Ziesche R, Hofbauer E, Wittmann K, Petkov V, Block L. A preliminary study of long-term treatment with interferon gamma-1b and low-dose prednisone in patients with idiopathic pulmonary fibrosis. *N Engl J Med* 1999; 341:1264-1269.
56. Hunninghake GW, Kalica AR. Approaches to the treatment of pulmonary fibrosis. *Am J Respir Crit Care Med* 1995; 151:915-918.

ADDRESS: Fernando J. Martinez MD, 1500 E. Medical Center Drive, Ann Arbor, MI 48109-0360, e-mail fmartine@umich.edu.