
KANDICE KOTTKE-MARCHANT, MD, PhD

Chair, Pathology and Laboratory Medicine Institute,
and Section Head, Hemostasis and Thrombosis,
Department of Clinical Pathology,
Cleveland Clinic, Cleveland, OH

Importance of platelets and platelet response in acute coronary syndromes

■ ABSTRACT

Platelet activation is one of the essential steps in the genesis and propagation of atherothrombosis. Accumulating clinical evidence suggests that an elevated platelet count, platelet activation, and platelet hyperreactivity (defined as residual platelet activity despite antiplatelet drug therapy) may be associated with adverse cardiovascular events in patients with acute coronary syndromes. Platelet function can be analyzed using various assays and measures of platelet activation. The best assays for measuring residual platelet activity in the setting of antiplatelet therapy are still being defined, as are their predictive values. Platelet aggregation remains the gold standard, but other testing methods offer advantages for specific applications, such as detecting overall platelet hyperreactivity in the presence of antiplatelet therapy or detecting inhibition of the adenosine diphosphate receptor P2Y₁₂. Standard testing protocols for platelet aggregation are needed to achieve consistency among studies.

■ KEY POINTS

Platelet function assays are inherently variable because they measure cell function rather than a single analyte.

Screening tests, or global tests for platelet function, do not identify specific causes of platelet dysfunction but combine measurement of different aspects of platelet function.

There appears to be a subgroup of patients with stable cardiovascular disease who have an increased risk of major cardiac events associated with platelet hyperreactivity.

For predicting cardiac events, receiver operating characteristic (ROC) curve analysis should be used to objectively define cutoff values for platelet hyperreactivity as opposed to reliance on arbitrary cutoff values.

Platelets play a substantial role in atherothrombosis, and platelet activation is implicated in the genesis of acute coronary syndromes (ACS) (Figure 1). This review describes platelet function and the mechanisms behind platelet activation, the utility of laboratory tests of platelet function for assessing cardiovascular risk, and the role of platelets in various phases of atherosclerosis. Against this backdrop, the article concludes by reviewing current evidence on the association between platelet hyperreactivity—defined as residual platelet activity despite antiplatelet drug therapy—and ACS. Here and throughout this supplement, ACS is understood to comprise unstable angina and myocardial infarction (MI) with or without ST-segment elevation.

■ PLATELET FUNCTION

Platelets are non-nucleated cells produced by megakaryocytes, which are very large cells (50 to 100 μ m in diameter) found in bone marrow. The megakaryocyte surface membrane forms protoplatelet extensions from which platelets “bud off” and are emitted into the circulation, where they number approximately 200,000 to 400,000 per microliter of blood.

Platelet activation

Platelets play a crucial role in the vascular response to injury, and activation of platelets has long been recognized as an important step. Platelets release dense granules that contain the nucleotide adenosine diphosphate (ADP), which activates other platelets. They also possess alpha granules, which contain proteins and protein mediators (eg, platelet-derived growth factor, platelet factor 4) that are involved in inflammatory processes. The platelet surface is coated with hundreds of thousands of receptors for other cells, including activated vascular wall cells and extracellular matrix proteins. Platelets possess an affinity for adherence, especially to injured vessel walls, where they release their granule contents and then aggregate. These properties promote platelets’ involvement in many vascular processes, including ACS, as will be explored below.

Platelets exist in a nonactivated state and are

See end of article for author disclosures. doi:10.3949/ccjm.76.s1.01

drawn passively into areas of vascular injury. Initially, they adhere to proteins such as von Willebrand factor, which is a large extracellular matrix protein produced by endothelial cells. The platelet glycoprotein Ib/IX/V binds to von Willebrand factor, forming a loose association that results in platelets rolling on the surface of the vessel wall. As a multimer, von Willebrand factor exists in one subunit that is dimerized and then polymerized, making it an ideal substrate for platelets because of the multiple substrates to which platelets can adhere.

Platelets are then engaged through receptors for collagen (ie, glycoprotein Ia/IIa or integrin $\alpha_2\beta_1$) and glycoprotein VI, leading to intracellular signaling and activation of the platelets. Platelet activation is followed by firm adhesion through engagement of another integrin, the $\alpha_{IIb}\beta_3$ (glycoprotein IIb/IIIa) receptor, on platelet surfaces for fibrinogen. The glycoprotein IIb/IIIa receptor is involved in a homotypic platelet–platelet interaction with an $\alpha_{IIb}\beta_3$ receptor on another platelet, which attracts further platelets and results in platelet–platelet adhesion, called platelet aggregation. This platelet cascade is depicted in Figure 2.

Platelet fibrinogen receptor

The platelet fibrinogen receptor (glycoprotein IIb/IIIa receptor) is an $\alpha_{IIb}\beta_3$ integrin that binds to arginine-glycine-aspartic acid (RGD) epitopes of proteins, such as fibrinogen. Fibrinogen has a two-dimensional symmetry, with RGD groups on both ends of the molecule, which makes it an ideal molecule for linking platelet to platelet.

von Willebrand factor has RGD groups, as do both fibronectin and glycoprotein IIb/IIIa vitronectin, and can therefore bind to many plasma and extracellular matrix proteins. The glycoprotein IIb/IIIa receptor is inactive in resting platelets. It becomes activated during the platelet activation process and binds to fibrinogen, which bridges to other platelets, causing aggregation.

ADP receptors

Various receptors on platelet surfaces are responsible for platelet activation. One is a family of receptors for ADP. As ADP is released from platelets, it can then activate other platelets by binding to the receptors. The ADP receptor P2Y₁₂ signals through G protein pathways and is coupled to adenylate cyclase, an enzyme that catalyzes the conversion of adenosine triphosphate to cyclic adenosine monophosphate (cAMP). High levels of cAMP inhibit platelet function; ADP binding to P2Y₁₂ shuts down adenylate cyclase, which leads to phosphoinositide 3-kinase activation and accelerated aggregation and platelet release.

A final notable factor in the mediation of platelet activation and aggregation is phospholipase A₂, which liberates arachidonic acid from the platelet membrane,

Platelet activation and hyperreactivity




FIGURE 1. Resting platelets stimulated by an agonist become activated; the activated platelets can trigger intravascular thrombosis and inflammatory processes and have been implicated in atherogenesis and development of acute coronary syndromes. Antiplatelet drug therapy is designed to result in a therapeutic response due to inhibited or diminished platelet activation in response to agonists. There is increasing evidence, however, that preserved platelet activity despite the presence of antiplatelet drugs, or platelet hyperreactivity, may be associated with adverse cardiovascular events.

metabolizing it through cyclooxygenase and thromboxane synthase to generate thromboxane A₂, which leads to release of platelet granule contents and aggregation of other platelets.

PLATELET FUNCTION TESTS

Platelet function assays are inherently variable because they measure cell function rather than a single analyte. Several new platelet testing devices have come to market with the goal of ease of use; many can now be used at the bedside to measure platelet function.

Platelet count

In my view, the platelet count remains one of the best tests for assessing bleeding risk, as a low platelet count is one of the most common causes of bleeding. However, the platelet count is not a functional assay because it does not evaluate other platelet functions.

Screening tests

Screening tests, or global tests for platelet function, do not identify specific causes of platelet dysfunction but combine measurement of many different aspects of platelet function, such as adhesion, aggregation and granule release.

Bleeding time. The bleeding time is an archaic test because of the poor correlation between bleeding time and

Sequence of platelet activation and aggregation




FIGURE 2. In the process of platelet activation, unactivated circulating platelets initially encounter an area of vascular injury and adhere rapidly to exposed von Willebrand factor (vWF) through the glycoprotein (GP) Ib/IX/V membrane receptor. Collagen in the extracellular matrix is also engaged through receptors $\alpha_2\beta_1$ (GPIa/IIa) and GPIIb/IIIa, leading to platelet shape change and activation. Cell signaling results in conformational change in the fibrinogen receptor GPIIb/IIIa ($\alpha_{IIb}\beta_3$), with binding to fibrinogen and fibrin leading to platelet aggregation and thrombus formation.

Reprinted, with permission, from Kottke-Marchant K, ed. *An Algorithmic Approach to Hemostasis Testing*. Northfield, IL: College of American Pathologists; 2008. Copyright © 2008 College of American Pathologists.

bleeding disorders or thrombotic disorders. Its utility in measuring platelet function is therefore highly limited.

PFA-100. The PFA-100 Platelet Function Analyzer system (PFA-100) is one example of a global platelet function assay that measures multiple platelet functions, including platelet adhesion and aggregation. The instrument, which is about the size of a bread box, uses a citrate-anticoagulated whole blood specimen to measure platelet reaction in a high-shear environment. Blood travels at high shear rates through membranes coated with either collagen and ADP or collagen and epinephrine (epinephrine receptors exist on platelet surfaces). Platelets adhere to the membranes and then activate, aggregate, and occlude a small aperture in the center of each membrane, yielding a measurable closure time.

Since the PFA-100 was developed before the availability of the thienopyridine antiplatelet drugs, its utility lies not in monitoring the effects of those agents but in its ability to detect aspirin-induced platelet dysfunction or intrinsic platelet function disorders. An abnormal epinephrine cartridge closure time in the presence of a normal ADP cartridge closure time indicates aspirin-induced platelet dysfunction. An abnormal closure time on both measures is indicative of von Willebrand disease or a platelet defect such as Glanzmann thrombasthenia or Bernard-Soulier syndrome.

Specific functional tests

Platelet aggregation. One of the most common methods of measuring platelet aggregation is called optical platelet aggregation. This technique, which is a high-complexity laboratory test, involves adding an aggregating agent (eg, ADP, epinephrine, thrombin, arachidonic acid) to platelet-rich plasma, a turbid platelet-rich suspension derived from whole blood. The effect of the aggregating agent on

the suspension's light transmittance is then measured to assess platelet aggregation progress (Figure 3).

Whole blood platelet aggregation is typically a high-complexity laboratory test. Recently, self-contained assay platforms that can measure whole blood aggregation have been developed. These are applicable for smaller hospitals and near-patient settings. One such rapid platelet function analyzer, known commercially as VerifyNow, offers point-of-care assessment of platelet function. The instrument, which is the size of a telephone answering machine, operates by a principle similar to that of optical platelet aggregation: platelet function is measured by the rate and extent of change in light transmittance in response to the introduction of agonists specific to various antiplatelet medications. Low light transmittance indicates a blood sample with inhibited platelet function; high light transmittance indicates normal platelet function.

Measurement of VASP phosphorylation. Vasodilator-stimulated phosphoprotein (VASP) is an intracellular platelet protein that is nonphosphorylated in basal state. The phosphorylation of VASP depends on the level of activation of the P2Y₁₂ receptor, a target of thienopyridine drugs. Thus, measuring VASP phosphorylation by flow cytometry using citrated whole blood can be a highly specific indicator of the action and efficacy of clopidogrel and other thienopyridine drugs.

A flow cytometry assay that measures VASP phosphorylation requires a whole blood sample that is incubated with ADP to measure what is called the platelet reactivity index. Adding ADP to whole blood stimulates adenylate cyclase, lowering cAMP and shutting off protein kinase, which results in low levels of VASP phosphorylation. Thus, if VASP is phosphorylated, the platelets are inhibited; if VASP is not phosphorylated, the platelets are acti-

vated. A satisfactory therapeutic response to clopidogrel or another thienopyridine drug produces a low platelet reactivity index, reflecting platelet inhibition.

■ ROLE OF PLATELETS IN ATHEROSCLEROSIS

Platelets serve major functions in three key aspects of atherosclerosis: atherogenesis, inflammation, and atherothrombosis.

Atherogenesis

Platelets play a pivotal role in atherogenesis.¹ They release matrix metalloproteinases that are involved in degrading the matrix in atherosclerotic plaques. Moreover, they contain and release chemokines and growth factors, including:

- RANTES, a chemokine that stimulates monocytes and T cells to increase the production of monocyte inflammatory mediators
- Platelet-derived growth factor, which stimulates the migration and proliferation of smooth muscle cells
- Transforming growth factor- β , which also stimulates proliferation of smooth muscle cells.

Inflammation

Activated platelets release inflammatory mediators and thereby change the adhesive and chemotactic properties of endothelial cells. Likewise, mediators derived from inflammatory cells (neutrophils) can affect platelet function.

Platelet-derived mediators include the following:

- Pro-interleukin (IL)- β , which triggers the synthesis of E-selectin that enables endothelial cells to interact with leukocytes
- Thromboxane A_2 , which increases neutrophil adhesion to facilitate platelet aggregation
- Platelet-derived growth factor and platelet factor 4, which increase neutrophil chemotaxis (the ability of neutrophils to infiltrate atherosclerotic plaque)
- CD40 ligand, a protein expressed on platelets that induces inflammatory responses in the endothelium
- P-selectin, a cell adhesion molecule expressed on activated platelets that enhances the adhesion of monocytes on activated endothelial cells.

Among the neutrophil-derived mediators, some—such as superoxide and leukotrienes—enhance platelet activation, whereas elastases inhibit platelet activation.

Overall, once inflammation begins in an atherosclerotic plaque, much reciprocal platelet activation can occur, so that the inflammatory process can become a feed-forward loop to eventually promote atherothrombosis.

Atherothrombosis

In the last stage of the atherosclerotic process, platelet enzymes that degrade the matrix may make plaques vulnerable to rupture by creating fissures in the fibrous




FIGURE 3. Optical platelet aggregation involves introduction of an aggregating agent to platelet-rich plasma to assess its effect on light transmission. The greater the increase in light transmission, the greater the platelet aggregation.

plaque cap. This exposes the lipid-rich core, which contains a significant amount of thromboplastin. Exposure to the extracellular matrix can lead to further platelet adhesion, activation, and aggregation. The development of a platelet thrombus is usually one of the ultimate steps in atherothrombosis leading to ACS, including MI.

■ ROLE OF PLATELETS IN ACUTE CORONARY SYNDROMES: WHAT IS THE EVIDENCE?

How predictive is an elevated platelet count?

Evidence suggests that a simple elevated platelet count may predict adverse outcomes following ACS. Among 10,793 patients with ST-segment-elevation MI in the Thrombolysis in Myocardial Infarction (TIMI) trials database, higher platelet counts on presentation were associated with higher rates of death, reinfarction, and development of congestive heart failure at 30 days (**Figure 4**).² In subsequent follow-up, a greater decrease in platelet counts after MI was associated with an increased risk of reinfarction.²

However, another study conducted in a slightly different population—1,616 patients with non-ST-segment-elevation MI/unstable angina—found no correlation between platelet count (by quintiles) and death at 60 months.³ The lowest mortality was observed in patients with a platelet count in the second-lowest quintile, although the highest mortality was indeed observed in the quintile of patients with the lowest platelet counts.³

The differing results in the above two studies suggest that additional platelet factors, beyond platelet count, contribute to the risk of adverse outcomes following ACS.

Platelet hyperreactivity and outcomes in ACS

Platelet hyperreactivity—ie, residual platelet activity despite antiplatelet therapy—appears to be involved

Higher platelet count associated with adverse outcomes in ST-elevation MI




FIGURE 4. Higher platelet counts were associated with adverse outcomes at 30 days among 10,793 patients with ST-segment-elevation myocardial infarction (MI) in the Thrombolysis In Myocardial Infarction (TIMI) trials database.²

Reprinted from *American Journal of Cardiology* (Ly HQ, et al. Association of platelet counts on presentation and clinical outcomes in ST-elevation myocardial infarction. *Am J Cardiol* 2006; 98:1-5). Copyright © 2006, with permission from Elsevier. www.sciencedirect.com/science/journal/00029149

in the spectrum of ACS. A recent study evaluated the association between hyperreactivity of platelets to ADP and outcomes in 600 patients with stable cardiovascular disease who were on aspirin therapy.⁴ Hyperreactivity was defined as a collagen/ADP closure time of less than 90 seconds on the PFA-100 system (short collagen/ADP closure time). On receiver operating characteristic (ROC) curve analysis, a short collagen/ADP closure time served as a significant predictor of recurrent events (relative risk [RR] = 3.65; 95% CI, 1.76-7.57) and death (RR = 6.56; 95% CI, 1.93-22.35) compared with a closure time of 90 seconds or greater. The authors concluded that there appears to be a subgroup of patients with stable cardiovascular disease who have an increased risk of major adverse events associated with platelet hyperreactivity.⁴

An earlier study by Harrison et al assessed platelet function using the PFA-100 in 78 patients presenting with acute chest pain classified as MI, unstable angina, or nonspecific chest pain.⁵ Using the PFA-100, they found shorter collagen/ADP closure times and higher levels of von Willebrand factor in subjects with MI compared with those who had unstable angina or nonspecific chest pain.⁵ Fuchs et al reported a similar association between

von Willebrand factor and outcomes in 208 patients with ACS,⁶ raising the possibility that von Willebrand factor, through its association with increased platelet adhesion and activation, may be a major contributor to risk in ACS.

Similarly, an association between platelet hyperreactivity and cardiovascular events has been suggested in patients with type 2 diabetes. In a 2007 study of 173 patients with type 2 diabetes and coronary artery disease receiving dual antiplatelet therapy (aspirin plus clopidogrel), the 2-year risk of major cardiovascular events was significantly higher in those in the highest quartile of platelet aggregation compared with those in the lower three quartiles (hazard ratio = 3.35; 95% CI, 1.68-6.66).⁷ In a separate study, Serebruany et al measured platelet activity by five different testing methods in 822 patients with coronary artery disease and found significantly higher platelet hyperreactivity by all methods in those patients who had diabetes (n = 257) than in those who did not (n = 565).⁸

Marcucci et al recently examined the relationship between clinical characteristics and residual platelet activity in 386 patients with ACS on dual antiplatelet therapy (aspirin plus clopidogrel).⁹ The presence of residual platelet activity (determined by platelet aggregation in response to the agonists arachidonic acid and ADP, as well as by the PFA-100) was associated with significantly higher inflammatory status, as determined by leukocyte count and erythrocyte sedimentation rate. The same association was observed among a subset of patients in this study undergoing percutaneous coronary intervention (PCI) who were receiving dual antiplatelet therapy; additionally, residual platelet activity was associated with a significantly higher incidence of diabetes and a significantly lower ejection fraction in this subset.⁹

Platelet hyperreactivity while on dual antiplatelet therapy (aspirin plus clopidogrel) was also found to be predictive of clinical outcome in a study of 195 patients with non-ST-elevation MI undergoing PCI.¹⁰ Hyporesponse to antiplatelet therapy, as measured by a high VASP platelet reactivity index (PRI), predicted an increased risk of recurrent ischemic events within 30 days of PCI. Using ROC curve analysis, the investigators found that a VASP PRI cutoff value of 53% (ie, a high PRI [$> 53\%$] indicates residual platelet activity despite clopidogrel) had a sensitivity of 93%, a specificity of 50%, a positive predictive value of 12%, and a negative predictive value of 99% for ischemic events.¹⁰ Similarly, among 144 patients undergoing PCI assessed for decreased platelet reactivity to a loading dose of clopidogrel, Bonello et al also found that a VASP PRI greater than 50% was optimal for predicting major adverse cardiovascular events: all 21 events in the study

occurred among patients whose VASP PRI was in the highest four quintiles.¹¹

CONCLUSIONS AND GENERAL ASSESSMENT OF PLATELET FUNCTION TESTS

Platelets clearly are involved in the pathogenesis of atherothrombosis. Accumulating evidence suggests that both an elevated platelet count and platelet hyperreactivity (residual platelet activity despite dual antiplatelet therapy) may be associated with adverse cardiovascular events in patients with ACS.

Platelet function can be measured using several different assays and measures of platelet activation. The best assays for measuring residual platelet activity in the setting of antiplatelet therapy are still being defined, as are their predictive values. Platelet aggregation remains the gold standard. The PFA-100 may detect overall platelet hyperreactivity despite the use of antiplatelet therapy, and is attracting increasing use for this purpose. VASP phosphorylation may be a good assay for detecting P2Y₁₂ inhibition but is limited to thienopyridines in terms of detecting platelet hyperreactivity. For predicting adverse cardiac events, ROC curve analysis should be used to objectively define cutoff values for platelet hyperreactivity as opposed to reliance on arbitrarily defined cutoff values.

Moving forward, standard testing protocols for platelet aggregation clearly are needed to achieve consistency among studies.

DISCLOSURES

Dr. Kottke-Marchant reported that she has no financial interests or relationships that pose a potential conflict of interest with this article.

This article was developed from an audio transcript of Dr. Kottke-Marchant's lecture at the CME course that formed the basis of this supplement. The transcript was edited and formatted by the *Cleveland Clinic Journal of Medicine* staff for clarity and conciseness, and was then reviewed, revised, and approved by Dr. Kottke-Marchant.

Dr. Kottke-Marchant received honoraria for contributing to this supplement and

the CME course on which it was based. The honoraria were paid by the Cleveland Clinic from the educational grant from Daiichi Sankyo, Inc., and Eli Lilly and Co. that supported the course and this supplement. These grantors had no input on the content of the course or this supplement.

REFERENCES

1. Davi G, Patrono C. Platelet activation and atherothrombosis. *N Engl J Med* 2007; 357:2482–2494.
2. Ly HQ, Kirtane AJ, Murphy SA, et al. Association of platelet counts on presentation and clinical outcomes in ST-elevation myocardial infarction (from the TIMI Trials). *Am J Cardiol* 2006; 98:1–5.
3. Mueller C, Neumann FJ, Hochholzer W, et al. The impact of platelet count on mortality in unstable angina/non-ST-segment elevation myocardial infarction. *Am Heart J* 2006; 151:1214.e1–7.
4. Christie DJ, Kottke-Marchant K, Gorman RT. Hypersensitivity of platelets to adenosine diphosphate in patients with stable cardiovascular disease predicts major adverse events despite antiplatelet therapy. *Platelets* 2008; 19:104–110.
5. Harrison P, Mackie I, Mathur A, et al. Platelet hyperfunction in acute coronary syndromes. *Blood Coagul Fibrinolysis* 2005; 16:557–562.
6. Fuchs I, Frossard M, Spiel A, Riedmüller E, Laggner AN, Jilma B. Platelet function in patients with acute coronary syndrome (ACS) predicts recurrent ACS. *J Thromb Haemost* 2006; 4:2547–2552.
7. Angiolillo DJ, Bernardo E, Sabaté M, et al. Impact of platelet reactivity on cardiovascular outcomes in patients with type 2 diabetes mellitus and coronary artery disease. *J Am Coll Cardiol* 2007; 50:1541–1547.
8. Serebruany V, Pokov I, Kuliczowski W, Chesebro J, Badimon J. Baseline platelet activity and response after clopidogrel in 257 diabetics among 822 patients with coronary artery disease. *Thromb Haemost* 2008; 100:76–82.
9. Marcucci R, Gori AM, Panicia R, et al. Residual platelet reactivity is associated with clinical and laboratory characteristics in patients with ischemic heart disease undergoing PCI on dual antiplatelet therapy. *Atherosclerosis* 2007; 195:e217–e223.
10. Frere C, Cuisset T, Quilici J, et al. ADP-induced platelet aggregation and platelet reactivity index VASP are good predictive markers for clinical outcomes in non-ST elevation acute coronary syndrome. *Thromb Haemost* 2007; 98:838–843.
11. Bonello L, Paganelli F, Arpin-Bornet M, et al. Vasodilator-stimulated phosphoprotein phosphorylation analysis prior to percutaneous coronary intervention for exclusion of postprocedural major adverse cardiovascular events. *J Thromb Haemost* 2007; 5:1630–1636.

Correspondence: Kandice Kottke-Marchant, MD, PhD, Chair, Pathology and Laboratory Medicine Institute, Cleveland Clinic, 9500 Euclid Avenue, L21, Cleveland, OH 44195; marchak@ccf.org