

THE CLINICAL PICTURE

JOSHUA M. BERLIN, MD

Dermatology Associates, PA,
of the Palm Beaches, Boynton Beach, FL

SANDRA M. GOLDBERG, MMS, PA-C

Dermatology Associates, PA, of the Palm Beaches,
Boynton Beach, FL

The Clinical Picture

A rash on the legs and palms



FIGURE 1




FIGURE 2

A 39-YEAR-OLD woman has had a rash on her legs and palms (**FIGURES 1 AND 2**) for 8 weeks. She has been treated with a mid-potency topical steroid, the high-potency topical corticosteroid fluocinonide (Vanos), an oral prednisone taper over 12 days, and loratadine (Claritin), with no improvement. Intermittently, the rash is moderately itchy. Biopsy reveals a psoriasiform hyperplasia with alternating orthokeratosis and parakeratosis.

Q: Which is the most likely diagnosis?

- Psoriasis
- Pityriasis rubra pilaris
- Dyshidrotic eczema
- Keratoderma
- Contact dermatitis

A: The diagnosis is pityriasis rubra pilaris, a rare condition with a prevalence of 1 in 5,000 to 50,000 new dermatology patient visits.¹ It is a papulosquamous disorder that presents with areas of hyperkeratosis on an erythematous base. Large red plaques often coalesce, leaving areas of uninvolved skin (“islands of sparing”).

doi:10.3949/ccjm.77a.09082

The palms and soles often reveal a distinctive orange-red waxy keratoderma.¹

Clinically, pityriasis rubra pilaris can be difficult to differentiate from psoriasis, and it can progress to disabling palmoplantar keratoderma and erythroderma.

■ BASIS OF THE DIAGNOSIS

The diagnosis is based on characteristic findings supported by classic features on skin biopsy. Microscopic study shows a psoriasiform dermatitis with alternating vertical and horizontal orthokeratosis and parakeratosis (the “checkerboard pattern”).

Although an underlying dysfunction in vitamin A metabolism has been suggested, the exact cause and pathogenesis of pityriasis rubra pilaris are not known.

■ TREATMENT

Treatment of pityriasis rubra pilaris can be difficult, as no one single treatment works for all patients. Systemic retinoids, methotrexate, phototherapy, and cyclosporine are commonly used. Recent reports have shown the effectiveness of infliximab (Remicade), a chimeric monoclonal antibody that binds to soluble and membrane-bound forms of tumor necrosis factor alpha.^{2,3}

In our patient, after a 2-month course of acitretin (Soriatane) failed, three treatments with infliximab—5 mg/kg at baseline, at 2 weeks, and at 6 weeks—led to complete resolution of the condition. ■

■ REFERENCES

1. Selvaag E, Haedersdal M, Thomsen K. Pityriasis rubra pilaris: a retrospective study of 12 patients. *J Eur Acad Dermatol Venereol* 2000; 14:514–515.
2. Liao WC, Mutasim DF. Infliximab for the treatment of adult-onset pityriasis rubra pilaris. *Arch Dermatol* 2005; 141:423–425.
3. Müller H, Gattringer C, Zelger B, Höpfl R, Eisendle K. Infliximab monotherapy as first-line treatment for adult-onset pityriasis rubra pilaris: case report and review of the literature on biologic therapy. *J Am Acad Dermatol* 2008; 59(suppl 5):S65–S70.

ADDRESS: Joshua M. Berlin, MD, 10301 Hagen Ranch Road, Suite 930, Boynton Beach, FL 33437; e-mail Joshberlin@hotmail.com.