

Vitamin B₁₂ deficiency

(JULY 2015)

TO THE EDITOR: In the article “An unusual cause of vitamin B₁₂ and iron deficiency,”¹ the diagnosis of vitamin B₁₂ deficiency was made only by a vitamin B₁₂ level of 108 pg/mL.

According to *Harrison's Principles of Internal Medicine*, 18th edition, page 870, the diagnosis of vitamin B₁₂ deficiency requires measurement of methylmalonic acid. Either this test was not performed on the 76-year-old woman described in the article, or the result was not entered. Without a methylmalonic acid level, the title of this article seems incorrect, or the article itself is incomplete by not including this level. The correct diagnosis of anemia due to an intestinal tapeworm was made by capsule endoscopy. She received appropriate therapy and her anemia cleared quickly.

If there is an updated concept for diagnosing vitamin B₁₂ deficiency, I'm open to learning about it.

KENNETH R. PHILLIPS, MS, MD, FCCP
Apponaug Diagnostics and Therapeutics
Warwick, RI

■ REFERENCE

1. Maithel S, Duong AK, Zhang J, Nguyen DL. An unusual cause of vitamin B₁₂ and iron deficiency. *Cleve Clin J Med* 2015; 82:406–408.

doi:10.3949/ccjm.83c.01001

IN REPLY: We thank Dr. Phillips for his inquiry.

In general, serum vitamin B₁₂ concentrations vary greatly, and we acknowledge that serum vitamin B₁₂ may be normal in up to 5% of patients with documented B₁₂ deficiency.¹ In a prospective study of 1,599 patients, Matchar et al² demonstrated that a single vitamin B₁₂ level less than 200 pg/mL

had a specificity greater than 95% at predicting vitamin B₁₂ deficiency.² We acknowledge that additional metabolite testing is necessary in equivocal cases in which the vitamin B₁₂ level is between 200 and 300 pg/mL, which is often considered to be the normal range, but the patient has symptoms of vitamin B₁₂ deficiency such as dementia and unexplained macrocytosis, and neurologic symptoms.³

Based on the patient's symptoms of neuropathy and fatigue in conjunction with a vitamin B₁₂ level well below 200 pg/mL, we believe that the diagnosis can be made.^{2,3} Nonetheless, although we did not mention it in our article, we did indeed send for a methylmalonic acid measurement at the time of the initial evaluation, and the level was elevated at 396 nmol/L (normal 87–318 nmol/L), further confirming her vitamin B₁₂ deficiency.

DOUGLAS L. NGUYEN, MD
Assistant Clinical Professor of Medicine
University of California, Irvine

SHELLY MAITHEL
University of California, Irvine

ALEX K. DUONG
University of California, Irvine

JONATHAN ZHANG, MD
University of California, Irvine


■ REFERENCES

1. Naurath HJ, Joosten E, Riezler R, Stabler SP, Allen RH, Lindenbaum J. Effects of vitamin B₁₂, folate, and vitamin B₆ supplements in elderly people with normal serum vitamin concentrations. *Lancet* 1995; 346:85–89.
2. Matchar DB, McCrory DC, Millington DS, Feussner JR. Performance of the serum cobalamin assay for diagnosis of cobalamin deficiency. *Am J Med Sci* 1994; 308:276–283.
3. Stabler SP. Clinical practice. Vitamin B₁₂ deficiency. *N Engl J Med* 2013; 368:149–160.

doi:10.3949/ccjm.83c.01002

AVAILABLE AT FINE BOOK STORES

Innovation the Cleveland Clinic Way



Innovation is a key asset in business and healthcare. Learn how you and your organization can put ideas to work.

This book tells you how to ignite mission-driven innovation, boost the culture and infrastructure of innovation, recruit and retain brilliant innovators and much more.

By Thomas J. Graham, MD

For details, visit
clevelandclinic.org/ClevelandClinicWay.



LETTERS TO THE EDITOR

Preoperative testing

(OCTOBER 2015)

TO THE EDITOR: I read with great interest your *1-Minute Consult* and the accompanying editorial on preoperative testing. I have long requested from my local hospitals the rationale for the long list of tests that used to be mandated for any surgery. I could not even get the courtesy of a reply from the department of anesthesia. For a while, in addition to the complete blood cell count and chemistry panel, one hospital demanded a urinalysis for cataract surgery.

Finally, without any explanation, the testing is now no longer mandated for cataract surgery but is still required for surgery such as the meniscus repair that was referenced.

These are not tests I want to order, but I am forced to order them or the surgery won't be done. Certainly, in a diabetic patient or a patient treated with a complex regimen for hypertension, tests may be needed.

Thank you for the opportunity to comment.

CYNTHIA POINT, MD
San Francisco, CA

doi:10.3949/ccjm.83c.01003