

THE CLINICAL PICTURE

NATSU FUKUI, MD
Inova Fairfax Hospital,
Falls Church, VA

MARK DELMAN, MD
Inova Fairfax Hospital,
Falls Church, VA

TUYETHOA N. VINH, MD
Inova Fairfax Hospital,
Falls Church, VA

SVETOLIK DJURKOVIC, MD, FACP
Inova Fairfax Hospital,
Falls Church, VA

Necrotizing fasciitis after a watercraft accident

Despite IV antibiotics and surgical debridement, his right leg had to be amputated



Figure 1. Left leg: bullous lesions from *Vibrio vulnificus* infection.

A 57-YEAR-OLD MAN was transferred to our hospital with leg pain and confusion. His family reported that he had injured his leg while launching a motorized personal watercraft at the North Carolina seashore 2 days before. He had a history of cirrhosis secondary to hepatitis C and alcohol abuse.

Physical examination revealed rashes on his legs with hemorrhagic bullous lesions and ecchymosis (**Figure 1**). He was hypotensive and had lactic acidosis, with blood lactate levels as high as 9.4 mmol/L (reference range 0.5–2.2 mmol/L). Despite aggressive hydration and broad-spectrum antibiotics provided at a previous hospital, he needed increasing vasopressor treatment.

Given his septicemia and recent marine exposure, *Vibrio vulnificus* infection was suspected, and antibiotics were switched to doxycycline and ceftazidime. He underwent urgent surgical debridement, ultimately requiring above-the-knee amputation of his right leg. He also required additional surgeries on his left leg.

Blood and wound cultures eventually grew *V vulnificus*, and surgical pathology confirmed the diagnosis of necrotizing fasciitis (**Figure 2**).

RISE IN *V VULNIFICUS* INFECTIONS IS ATTRIBUTED TO GLOBAL WARMING

V vulnificus infection occurs most commonly from consuming raw shellfish, especially oysters, but it also occurs after exposure of an open wound to contaminated salt water. The pathogen is a gram-negative bacterium that resides in coastal waters worldwide, but in the United States it is usually seen on the Pacific and Gulf coasts¹ during the summer.²

doi:10.3949/ccjm.85a.17096

Although only 58 cases of *V vulnificus* infection were reported to the US Centers for Disease Control and Prevention in 1997, the number more than doubled to 124 in 2014.¹ This rise is suspected to be due in part to warmer coastal waters associated with global warming.²

Various marine pathogens can cause wound infections, but *V vulnificus* is most commonly implicated in deaths and hospitalizations.² Immunocompromised patients and those with liver disease are at particularly high risk of rapid progression to septic shock.

First-line antibiotics are doxycycline plus a third-generation cephalosporin.³ Studies have shown a direct correlation between delay of antibiotics and death,⁴ and early surgery is critical.⁵

Given the rising incidence of *V vulnificus* infection, it is increasingly important for providers across the United States to be aware of this infection.




Figure 2. Necrosis of deep dermis, subcutaneous fat, and fascial tissue with polymorphonuclear cell infiltration (hematoxylin and eosin, $\times 200$).

REFERENCES

1. **Centers for Disease Control and Prevention.** National enteric disease surveillance: COVIS annual summary, 2014. US Department of Health and Human Services, Atlanta, GA. 2014. www.cdc.gov/national-surveillance/pdfs/covis-annual-summary-2014-508c.pdf. Accessed May 8, 2018.
2. **Newton A, Kendall M, Vugia DJ, Henao OL, Mahon BE.** Increasing rates of vibriosis in the United States, 1996–2010: review of surveillance data from 2 systems. *Clin Infect Dis* 2012; 54(suppl 5):S391–S395. doi:10.1093/cid/cis243
3. **Stevens DL, Bisno AL, Chambers HF, et al.** Practice guidelines for the diagnosis and management of skin and soft tissue infections: 2014 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2014; 59(2):147–159. doi:10.1093/cid/ciu444
4. **Klontz KC, Lieb S, Schreiber M, Janowski HT, Baldy LM, Gunn RA.** Syndromes of *Vibrio vulnificus* infections. Clinical and epidemiologic features in Florida cases, 1981–1987. *Ann Intern Med* 1988; 109:318–323. PMID:3260760
5. **Chao WN, Tsai CF, Chang HR, et al.** Impact of timing of surgery on outcome of *Vibrio vulnificus*-related necrotizing fasciitis. *Am J Surg* 2013; 206(1):32–39. doi:10.1016/j.amjsurg.2012.08.008

ADDRESS: Natsu Fukui, MD, Inova Fairfax Hospital, Department of Medicine, 3300 Gallows Road, Falls Church, VA 22042; fukui.natsua@gmail.com