

BRIAN AGGANIS, MD
Department of Internal Medicine,
Brown University, Providence, RI

DAVID LEE, MD
Department of Internal Medicine,
Brown University, Providence, RI

THOMAS SEPE, MD
Department of Gastroenterology,
Brown University, Providence, RI

Liver enzymes: No trivial elevations, even if asymptomatic

ABSTRACT

Primary care physicians are at the forefront in screening for abnormal levels of liver enzymes and investigating the likely causes by obtaining a detailed history and physical examination, followed by appropriate laboratory and diagnostic workup. This review outlines common causes for the two main mechanisms of liver injury—cholestasis and hepatocellular insult—and explores the associated risk factors, methods of diagnosis, and management, with a focus on nonalcoholic fatty liver disease, one of the most often encountered causes of abnormal liver enzyme levels.

KEY POINTS

Disorders of hepatocellular injury tend to elevate levels of aminotransferases, whereas cholestatic disorders cause elevations of alkaline phosphatase and bilirubin.

The three most common causes of liver enzyme elevation are alcohol toxicity, medication overdose, and fatty liver disease.

Other disorders of liver dysfunction include hereditary hemochromatosis, viral hepatitis, autoimmune hepatitis, primary biliary cholangitis, primary sclerosing cholangitis, and alpha-1 antitrypsin disease.

Nonhepatic causes of elevated “liver enzymes” also need to be considered. For instance, rhabdomyolysis causes elevations in aminotransferase levels.

ELEVATED LEVELS OF CIRCULATING enzymes that are frequently of hepatic origin (aminotransferases and alkaline phosphatase) and bilirubin in the absence of symptoms are common in clinical practice. A dogmatic but true statement holds that there are no trivial elevations in these substances. All persistent elevations of liver enzymes need a methodical evaluation and an appropriate working diagnosis.¹

Here, we outline a framework for the workup and treatment of common causes of liver enzyme elevations.

■ PATTERN OF ELEVATION: CHOLESTATIC OR HEPATOCELLULAR

Based on the pattern of elevation, causes of elevated liver enzymes can be sorted into disorders of cholestasis and disorders of hepatocellular injury (Table 1).¹

Cholestatic disorders tend to cause elevations in alkaline phosphatase, bilirubin, and gamma-glutamyl transferase (GGT).

Hepatocellular injury raises levels of alanine aminotransferase (ALT) and aspartate aminotransferase (AST).

■ HOW SHOULD ABNORMAL RESULTS BE EVALUATED?

When approaching liver enzyme elevations, the clinician should develop a working differential diagnosis based on the medical and social history and physical examination.

Think about alcohol, drugs, and fat

The most common causes of liver enzyme elevation are alcohol toxicity, medication over-

dose, and fatty liver disease.

Alcohol intake should be ascertained. “Significant” consumption is defined as more than 21 drinks per week in men or more than 14 drinks per week in women, over a period of at least 2 years.²

The exact pathogenesis of alcoholic hepatitis is incompletely understood, but alcohol is primarily metabolized by the liver, and damage likely occurs during metabolism of the ingested alcohol. AST elevations tend to be higher than ALT elevations; the reason is ascribed to hepatic deficiency of pyridoxal 5'-phosphate, a cofactor of the enzymatic activity of ALT, which leads to a lesser increase in ALT than in AST.

Alcoholic liver disease can be difficult to diagnose, as many people are initially reluctant to fully disclose how much they drink, but it should be suspected when the ratio of AST to ALT is 2 or greater.

In a classic study, a ratio greater than 2 was found in 70% of patients with alcoholic hepatitis and cirrhosis, compared with 26% of patients with postnecrotic cirrhosis, 8% with chronic hepatitis, 4% with viral hepatitis, and none with obstructive jaundice.³ Importantly, the disorder is often correctable if the patient is able to remain abstinent from alcohol over time.

A detailed medication history is important and should focus especially on recently added medications, dosage changes, medication overuse, and use of nonprescription drugs and herbal supplements. Common medications that affect liver enzyme levels include statins, which cause hepatic dysfunction primarily during the first 3 months of therapy, nonsteroidal anti-inflammatory drugs, antiepileptic drugs, antibiotics, anabolic steroids, and acetaminophen (Table 2).¹ Use of illicit drugs and herbal remedies should be discussed, as they may cause toxin-mediated hepatitis.

Although inflammation from drug toxicity will resolve if the offending agent is discontinued, complete recovery may take weeks to months.⁴

A pertinent social history includes exposure to environmental hepatotoxins such as amatoxin (contained in some wild mushrooms) and occupational hazards (eg, vinyl chloride). Risk factors for viral hepatitis should be evaluated, including intravenous drug use,

TABLE 1

Liver disease and associated liver enzyme elevations

Hepatocellular disease

(aminotransferase elevations predominate)

Common

Alcoholic liver disease
Autoimmune hepatitis
Chronic viral hepatitis
Genetic hemochromatosis
(northern European ethnicity)
Medication toxicity (see Table 2)
Nonalcoholic fatty liver disease

Less common

Alpha-1 antitrypsin deficiency
Wilson disease

Cholestatic disease

(alkaline phosphatase, bilirubin, and gamma-glutamyl transferase elevations predominate)

Common

Biliary obstruction (eg, gallstones)
Drug hepatotoxicity
Neoplasm
Primary biliary cirrhosis
Primary sclerosing cholangitis

Less common

Autoimmune cholangiopathy
Sarcoidosis

From Aragon G, Zounossi ZM. When and how to evaluate mildly elevated liver enzymes in apparently healthy patients. *Cleve Clin J Med* 2010; 77(3):195–204. doi:10.3949/ccjm.77a.09064.

blood transfusions, unprotected sexual contact, organ transplant, perinatal transmission, and a history of work in healthcare facilities or travel to regions in which hepatitis A or E is endemic.

The medical and family history should include details of associated conditions, such as:

- Right heart failure (a cause of congestive hepatopathy)
- Metabolic syndrome (associated with fatty liver disease)
- Inflammatory bowel disease and primary sclerosing cholangitis
- Early-onset emphysema and alpha-1 antitrypsin deficiency.

The physical examination should be thor-

‘Significant’ alcohol consumption is > 21 drinks per week in men, and > 14 drinks per week in women, for ≥ 2 years

TABLE 2

Hepatotoxicity of selected drugs**Hepatocellular abnormalities**

Acetaminophen—acute hepatitis
 Allopurinol—granuloma
 Azathioprine—veno-occlusive disease, nodular regenerative hyperplasia
 Chaparral—portal inflammation with bile duct proliferation, lobar necrosis, and collapse
 Diclofenac, other nonsteroidal anti-inflammatory drugs
 Hydralazine—granuloma
 Isoniazid
 Methotrexate—fibrosis
 Methyldopa
 Mistletoe—hepatocellular injury
 Nitrofurantoin—autoimmune-like disease
 Quinidine—granuloma
 Statins
 Toxic alkaloid—veno-occlusive disease

Cholestatic abnormalities

Amoxicillin-clavulanate, other penicillin derivatives
 Anabolic steroids—cholestasis, peliosis hepatis, neoplasm
 Captopril
 Carbamazepine
 Chlorpromazine
 Erythromycin estolate
 Estrogens
 Kava—hepatic necrosis, cholestasis, lobular hepatitis
 Oral contraceptives
 Phenytoin—mononucleosis-like syndrome
 Sulfa drugs

Drug-induced fatty liver

(with or without hepatocellular abnormalities)
 Amiodarone—phospholipidosis
 Anabolic steroids—cholestasis, peliosis hepatis, neoplasm
 Cocaine—microvesicular steatosis
 Corticosteroids
 Jin Bu Huan—focal hepatic necrosis, steatosis, bridging fibrosis
 Tetracycline
 Valproic acid

Adapted from Aragon G, Zounossi ZM. When and how to evaluate mildly elevated liver enzymes in apparently healthy patients. *Cleve Clin J Med* 2010; 77(3):195–204. doi:10.3949/ccjm.77a.09064.

ough, with emphasis on the abdomen, and search for stigmata of advanced liver disease such as hepatomegaly, splenomegaly, ascites, edema, spider angiomas, jaundice, and asterix. Any patient with evidence of chronic

liver disease should be referred to a subspecialist for further evaluation.

Further diagnostic workup

Abnormal liver enzyme findings or physical examination findings should direct the subsequent diagnostic workup with laboratory testing and imaging.⁵

For cholestasis. If laboratory data are consistent with cholestasis or abnormal bile flow, it should be further characterized as extrahepatic or intrahepatic. Common causes of extrahepatic cholestasis include biliary tree obstruction due to stones or malignancy, often visualized as intraductal biliary dilation on ultrasonography of the right upper quadrant. Common causes of intrahepatic cholestasis include viral and alcoholic hepatitis, nonalcoholic steatohepatitis, certain drugs and toxins such as alkylated steroids and herbal medications, infiltrative diseases such as amyloid, sarcoid, lymphoma, and tuberculosis, and primary biliary cholangitis.

Abnormal findings on ultrasonography should be further pursued with advanced imaging, ie, computed tomography or magnetic resonance cholangiopancreatography (MRCP). The confirmation of a lesion on imaging is often followed by endoscopic retrograde cholangiopancreatography (ERCP) in an attempt to obtain biopsy samples, remove obstructions, and place therapeutic stents. In instances when endoscopic attempts fail to relieve the obstruction, surgical referral may be appropriate.

For nonhepatobiliary problems. Depending on clinical presentation, it may also be important to consider nonhepatobiliary causes of elevated liver enzymes.

Alkaline phosphatase is found in many other tissue types, including bone, kidney, and the placenta, and can be elevated during pregnancy, adolescence, and even after fatty meals due to intestinal release.⁶ After screening for the aforementioned physiologic conditions, isolated elevated alkaline phosphatase should be further evaluated by obtaining GGT or 5-nucleotidase levels, which are more specifically of hepatic origin. If these levels are within normal limits, further evaluation for conditions of bone growth and cellular turnover such as Paget disease, hyperparathyroidism,

Up to 1/3
of the US
population—
100 million
people—
may have
NAFLD,
and 4–6 million
may have NASH

and malignancy should be considered. Specifically, Stauffer syndrome should be considered when there is a paraneoplastic rise in the alkaline phosphatase level in the setting of renal cell carcinoma without liver metastases.

AST and ALT levels may also be elevated in clinical situations and syndromes unrelated to liver disease. Rhabdomyolysis, for instance, may be associated with elevations of AST in more than 90% of cases, and ALT in more than 75%.⁷ Markers of muscle injury including serum creatine kinase should be obtained in the setting of heat stroke, muscle weakness, strenuous activity, or seizures, as related elevations in AST and ALT may not always be clinically indicative of liver injury.

Given the many conditions that may cause elevated liver enzymes, evaluation and treatment should focus on identifying and removing offending agents and targeting the underlying process with appropriate medical therapy.

■ FATTY LIVER

With rates of obesity and type 2 diabetes on the rise in the general population, identifying and treating nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH) require increased awareness and close coordination between primary care providers and subspecialists.

According to current estimates, up to one-third of the US population (100 million people) may have NAFLD, and 1% to 3% of the population (4–6 million people) likely have NASH, defined as steatosis with inflammation. Development of NASH places patients at a significantly higher risk of fibrosis, hepatocellular injury, and cancer.⁸

NAFLD is more common in men than in women. It is present in around 80% to 90% of obese adults, two-thirds of adults with type 2 diabetes, and many people with hyperlipidemia. It is also becoming more common in children, with 40% to 70% of obese children likely having some element of NAFLD.

Diagnosis of fatty liver

Although liver enzymes are more likely to be abnormal in individuals with NAFLD, many individuals with underlying NAFLD may have normal laboratory evaluations. ALT may be elevated in only up to 20% of cases and

does not likely correlate with the level of underlying liver damage, although increasing GGT may serve as a marker of fibrosis over time.^{9–11} In contrast to alcohol injury, however, the AST-ALT ratio is usually less than 1.0.

Noninvasive tools for diagnosing NAFLD include the NAFLD fibrosis score, which incorporates age, hyperglycemia, body mass index, platelet count, albumin level, and AST-ALT ratio. This and related scoring algorithms may be useful in differentiating patients with minimal fibrosis from those with advanced fibrosis.^{12,13}

Ultrasonography is a first-line diagnostic test for steatosis, although it may demonstrate fatty infiltration only around 60% of the time. Computed tomography and magnetic resonance imaging are more sensitive, but costlier. Transient elastography (FibroScan; Echosens, Paris, France) has become more popular and has been shown to correlate with findings on liver biopsy in diagnosing or excluding advanced liver fibrosis.^{14,15}

The gold standard for diagnosing NAFLD and NASH is identifying fat-laden hepatocytes or portal inflammation on biopsy; however, biopsy is generally reserved for cases in which the diagnosis remains uncertain.

Behavioral treatment

The primary treatment for NAFLD consists of behavioral modification including weight loss, exercise, and adherence to a low-fat diet, in addition to tight glycemic control and treatment of any underlying lipid abnormalities. Studies have shown that a reduction of 7% to 10% of body weight is associated with a decrease in the inflammation of NAFLD, though no strict guidelines have been established.¹⁶

Given the prevalence of NAFLD and the need for longitudinal treatment, primary care physicians will play a significant role in long-term monitoring and management of patients with fatty liver disease.

■ OTHER DISORDERS OF LIVER FUNCTION

Hereditary hemochromatosis

Hereditary hemochromatosis is the most common inherited liver disorder in adults of European descent,¹⁷ and can be effectively treated if discovered early. But its clinical diagnosis can be challenging, as many patients have no

Treatment for NAFLD: weight loss, exercise, low-fat diet, tight glycemic control, lipid control

Screen for HCV in those with HCV exposure, HIV infection, IV drug use, incarceration, or in those born 1945 to 1965 (one-time testing)

symptoms at presentation despite abnormal liver enzyme levels. Early symptoms may include severe fatigue, arthralgias, and, in men, impotence, before the appearance of the classic triad of “bronze diabetes” with cirrhosis, diabetes, and darkening of the skin.¹⁸

If hemochromatosis is suspected, laboratory tests should include a calculation of percent transferrin saturation, with saturation greater than 45% warranting serum ferritin measurement to evaluate for iron overload (ferritin > 200–300 ng/mL in men, > 150–200 ng/mL in women).¹⁹ If iron overload is confirmed, referral to a gastroenterologist is recommended.

Genetic evaluation is often pursued, but patients may ultimately require liver biopsy regardless of the findings, as some patients homozygous for the *HFE* mutation C282Y may not have clinical hemochromatosis, whereas others with hereditary hemochromatosis may not have the *HFE* mutation.

Therapeutic phlebotomy is the treatment of choice, and most patients tolerate it well.

Chronic hepatitis B virus and hepatitis C virus infections

Chronic hepatitis B virus (HBV) and hepatitis C virus (HCV) infections are common in the United States, with HBV affecting more than 1 million people and HCV affecting an estimated 3.5 million.

Chronic HCV infection. Direct-acting antiviral drugs have revolutionized HCV treatment and have led to a sustained viral response and presumed cure at 12 weeks in more than 95% of cases across all HCV genotypes.²⁰ Given the recent development of effective and well-tolerated treatments, primary care physicians have assumed a pivotal role in screening for HCV.

The American Association for the Study of Liver Diseases and the Infectious Diseases Society of America²¹ recommend screening for HCV in people who have risk factors for it, ie:

- HCV exposure
- HIV infection
- Behavioral or environmental risks for contracting the virus such as intravenous drug use or incarceration
- Birth between 1945 and 1965 (one-time testing).

If HCV antibody screening is positive,

HCV RNA should be obtained to quantify the viral load and confirm active infection, and genotype testing should be performed to guide treatment. Among the 6 most common HCV genotypes, genotype 1 is the most common in North America, accounting for over 70% of cases in the United States.

Although recommendations and therapies are constantly evolving, the selection of a treatment regimen and the duration of therapy are determined by viral genotype, history of prior treatment, stage of liver fibrosis, potential drug interactions, and frequently, medication cost and insurance coverage.

HBV infection. The treatment for acute HBV infection is generally supportive, though viral suppression with tenofovir or entecavir may be required for those who develop coagulopathy, bilirubinemia, or liver failure. Treatment of chronic HBV infection may not be required and is generally considered for those with elevated ALT, high viral load, or evidence of liver fibrosis on noninvasive measurements such as transient elastography.

Autoimmune hepatitis

Autoimmune causes of liver enzyme elevations should also be considered during initial screening. Positive antinuclear antibody and positive antismooth muscle antibody tests are common in cases of autoimmune hepatitis.²² Autoimmune hepatitis affects women more often than men, with a ratio of 4:1. The peaks of incidence occur during adolescence and between ages 30 and 45.²³

Primary biliary cholangitis

Additionally, an elevated alkaline phosphatase level should raise concern for underlying primary biliary cholangitis (formerly called primary biliary cirrhosis), an autoimmune disorder that affects the small and medium intrahepatic bile ducts. Diagnosis of primary biliary cholangitis can be assisted by a positive test for antimitochondrial antibody, present in almost 90% of patients.²⁴

Primary sclerosing cholangitis

Elevated alkaline phosphatase is also the hallmark of primary sclerosing cholangitis, which is associated with inflammatory bowel disease.²⁵ Primary sclerosing cholangitis is

characterized by inflammation and fibrosis of the intrahepatic and extrahepatic bile ducts, which are visualized on MRCP and confirmed by biopsy if needed.

REFERRAL

Subspecialty referral should be considered if the cause remains ambiguous or unknown, if

there is concern for a rare hepatic disorder such as an autoimmune condition, Wilson disease, or alpha-1 antitrypsin deficiency, or if there is evidence of advanced or chronic liver disease.

Primary care physicians are at the forefront of detecting and diagnosing liver disease, and close coordination with subspecialists will remain crucial in delivering patient care.

REFERENCES

1. Aragon G, Younossi ZM. When and how to evaluate mildly elevated liver enzymes in apparently healthy patients. *Cleve Clin J Med* 2010; 77(3):195–204. doi:10.3949/ccjm.77a.09064
2. Chalasani N, Younossi Z, Lavine JE, et al; American Gastroenterological Association; American Association for the Study of Liver Diseases; American College of Gastroenterology. The diagnosis and management of non-alcoholic fatty liver disease: practice guideline by the American Gastroenterological Association, American Association for the Study of Liver Diseases, and American College of Gastroenterology. *Gastroenterology* 2012; 142(7):1592–1609. doi:10.1053/j.gastro.2012.04.001
3. Cohen JA, Kaplan MM. The SGOT/SGPT ratio—an indicator of alcoholic liver disease. *Dig Dis Sci* 1979; 24(11):835–838. PMID:520102
4. Kaplan MM. Alanine aminotransferase levels: what's normal? *Ann Intern Med* 2002; 137(1):49–51. PMID:12093245
5. Pratt DS, Kaplan MM. Evaluation of abnormal liver enzyme results in asymptomatic patients. *N Engl J Med* 2000; 342(17):1266–1271. doi:10.1056/NEJM200004273421707
6. Sharma U, Pal D, Prasad R. Alkaline phosphatase: an overview. *Indian J Clin Biochem* 2014; 29(3):269–278. doi:10.1007/s12291-013-0408-y
7. Weibrecht K, Dayno M, Darling C, Bird SB. Liver aminotransferases are elevated with rhabdomyolysis in the absence of significant liver injury. *J Med Toxicol* 2010; 6(3):294–300. doi:10.1007/s13181-010-0075-9
8. Bellentani S, Scaglioni F, Marino M, Bedogni G. Epidemiology of non-alcoholic fatty liver disease. *Dig Dis* 2010; 28(1):155–161. doi:10.1159/000282080
9. Adams LA, Feldstein AE. Non-invasive diagnosis of nonalcoholic fatty liver and nonalcoholic steatohepatitis. *J Dig Dis* 2011; 12(1):10–16. doi:10.1111/j.1751-2980.2010.00471.x
10. Fracanzani AL, Valenti L, Bugianesi E, et al. Risk of severe liver disease in nonalcoholic fatty liver disease with normal aminotransferase levels: a role for insulin resistance and diabetes. *Hepatology* 2008; 48(3):792–798. doi:10.1002/hep.22429
11. Tahan V, Canbakan B, Balci H, et al. Serum gamma-glutamyltranspeptidase distinguishes non-alcoholic fatty liver disease at high risk. *Hepatogastroenterology* 2008; 55(85):1433–1438. PMID:18795706
12. McPherson S, Stewart S, Henderson E, Burt AD, Day CP. Simple non-invasive fibrosis scoring systems can reliably exclude advanced fibrosis in patients with non-alcoholic fatty liver disease. *Gut* 2010; 59(9):1265–1269. doi:10.1136/gut.2010.216077
13. Angulo P, Hui JM, Marchesini G, et al. The NAFLD fibrosis score: a noninvasive system that identifies liver fibrosis in patients with NAFLD. *Hepatology* 2007; 45(4):846–854. doi:10.1002/hep.21496
14. Petta S, Vanni E, Bugianesi E, et al. The combination of liver stiffness measurement and NAFLD fibrosis score improves the noninvasive diagnostic accuracy for severe liver fibrosis in patients with nonalcoholic fatty liver disease. *Liver Int* 2015; 35(5):1566–1573. doi:10.1111/liv.12584
15. Hashemi SA, Alavian SM, Gholami-Fesharaki M. Assessment of transient elastography (FibroScan) for diagnosis of fibrosis in non-alcoholic fatty liver disease: a systematic review and meta-analysis. *Caspian J Intern Med* 2016; 7(4):242–252. PMID:27999641
16. Promrat K, Kleiner DE, Niemeier HM, et al. Randomized controlled trial testing the effects of weight loss on nonalcoholic steatohepatitis. *Hepatology* 2010; 51(1):121–129. doi:10.1002/hep.23276
17. Adams PH, Reboussin DM, Barton JC, et al. Hemochromatosis and iron-overload screening in a racially diverse population. *N Engl J Med* 2005; 352(17):1769–1778. doi: 10.1056/NEJMoa041534
18. Brissot P, de Bels F. Current approaches to the management of hemochromatosis. *Hematology Am Soc Hematol Educ Program* 2006; 2006(1):36–41. doi:10.1182/asheducation-2006.1.36
19. Bacon BR, Adams PC, Kowdley KV, Powell LW, Tavill AS; American Association for the Study of Liver Diseases. Diagnosis and management of hemochromatosis: 2011 practice guideline by the American Association for the Study of Liver Diseases. *Hepatology* 2011; 54(1):328–343. doi:10.1002/hep.24330
20. Weiler N, Zeuzem S, Welker MW. Concise review: interferon-free treatment of hepatitis C virus-associated cirrhosis and liver graft infection. *World J Gastroenterol* 2016; 22(41):9044–9056. doi:10.3748/wjg.v22.i41.9044
21. American Association for the Study of Liver Disease, Infectious Diseases Society of America. HCV guidance: recommendations for testing, managing, and treating hepatitis C. www.hcvguidelines.org. Accessed July 16, 2018.
22. Manns MP, Czaja AJ, Gorham JD, et al; American Association for the Study of Liver Diseases. Diagnosis and management of autoimmune hepatitis. *Hepatology* 2010; 51(6):2193–2213. doi:10.1002/hep.23584
23. Liberal R, Krawitt EL, Vierling JM, Manns MP, Mieli-Vergani G, Vergani D. Cutting edge issues in autoimmune hepatitis. *J Autoimmun* 2016; 75:6–19. doi:10.1016/j.jaut.2016.07.005
24. Mousa HS, Carbone M, Malinverno F, Ronca V, Gershwin ME, Invernizzi P. Novel therapeutics for primary biliary cholangitis: Toward a disease-stage-based approach. *Autoimmun Rev* 2016; 15(9):870–876. doi:10.1016/j.autrev.2016.07.003
25. de Vries AB, Janse M, Blokzijl H, Weersma RK. Distinctive inflammatory bowel disease phenotype in primary sclerosing cholangitis. *World J Gastroenterol* 2015; 21(6):1956–1971. doi:10.3748/wjg.v21.i6.1956

ADDRESS: Brian Agganis, MD, Department of Internal Medicine, Brown University, 593 Eddy Street, Jane Brown Ground, Suite 0100, Providence, RI 02903; brian.agganis@lifespan.org

Liver enzymes

AUGUST 2018

In the article by Agganis B, Lee D, Sepe T (Liver enzymes: No trivial elevations, even if asymptomatic. *Cleve Clin J Med* 2018; 85(8):612–617, doi:10.3949/ccjm.85a.17103), an error occurred on page 613, in the second paragraph in the section about alcohol intake. The words ALT and AST were reversed. The paragraph should read as follows:

The exact pathogenesis of alcoholic hepatitis is

incompletely understood, but alcohol is primarily metabolized by the liver, and damage likely occurs during metabolism of the ingested alcohol. AST elevations tend to be higher than ALT elevations; the reason is ascribed to hepatic deficiency of pyridoxal 5'-phosphate, a cofactor of the enzymatic activity of ALT, which leads to a lesser increase in ALT than in AST.

We thank Avinash Alexander, MD, Texas Tech University Health Sciences Center, for calling this to our attention. The correction has been made online.