

Diabetic retinopathy: Screening, prevention, and treatment

In Reply: We thank Ashley Lim for the comments on our review article.¹ We are pleased that readers are interested in our review and appreciate the opportunity to discuss any concerns raised in the letter.

We agree that new therapies such as glucagon-like peptide-1 (GLP-1) receptor agonists and sodium-glucose cotransporter 2 inhibitors have significant value in achieving glycemic control to prevent diabetic retinopathy, and that GLP-1 receptor agonists have complex and unclear effects on the risk for diabetic retinopathy. Yet, several studies reported findings showing that sodium-glucose cotransporter 2 inhibitors are either not associated with a significantly increased risk of diabetic retinopathy or may even be linked to a significantly reduced risk, findings that seem to support their safety.²⁻⁶

However, as we discuss in the “Glucagon-like peptide-1 receptor agonists, rapid HbA1c reduction, and retinopathy” section of our review,¹ there

is mixed evidence on the effect of GLP-1 receptor agonists on the risk of diabetic retinopathy, with some meta-analyses and trials reporting an increased risk⁷⁻¹³ and others reporting no significant association between GLP-1 receptor agonists and risk for diabetic retinopathy or diabetic retinopathy progression.¹⁴⁻¹⁸ Consistent with the findings of Wai et al,¹⁹ the studies cited in our review that reported an increased risk of diabetic retinopathy observed an early rise in risk from 3 months to 3 years after patients started a GLP-1 receptor agonist.¹¹

Because of the mixed results in the literature, further investigations on treatment methods that can drastically reduce hemoglobin A1c, such as specific types of GLP-1 receptor agonists, other pharmacologic therapies, and bariatric surgery, are necessary to elucidate the mechanism of potential early worsening of diabetic retinopathy and safe management of this complication of diabetes.

In clinical practice, it is important for clinicians to follow the American Diabetes Association screening recommendations²⁰ for diabetic retinopathy and to consider referring patients to an ophthalmologist to assess retinopathy status when prescribing GLP-1 receptor agonists.

David D. Chong, BA
Case Western Reserve University School of
Medicine, Cleveland, OH

Nikhil Das, MD
Center for Ophthalmic Bioinformatics,
Cole Eye Institute, Cleveland Clinic, Cleveland, OH

Rishi P. Singh, MD
Vice President, Chief Medical Officer,
Cleveland Clinic Martin Hospitals,
Cleveland Clinic Florida, Stuart, FL; Professor,
Cleveland Clinic Lerner College of Medicine of
Case Western Reserve University, Cleveland, OH

REFERENCES

1. Chong DD, Das N, Singh RP. Diabetic retinopathy: screening, prevention, and treatment. *Clev Clin J Med* 2024; 91(8):503–510. doi:10.3949/ccjm.91a.24028
2. Tang H, Li G, Zhao Y, et al. Comparisons of diabetic retinopathy events associated with glucose-lowering drugs in patients with type 2 diabetes mellitus: a network meta-analysis. *Diabetes Obes Metab* 2018; 20(5):1262–1279. doi:10.1111/dom.13232
3. Ott C, Jumar A, Striepe K, et al. A randomised study of the impact of the SGLT2 inhibitor dapagliflozin on microvascular and macrovascular circulation. *Cardiovasc Diabetol* 2017; 16(1):26. doi:10.1186/s12933-017-0510-1
4. Yen FS, Wei JC, Yu TS, Hung YT, Hsu CC, Hwu CM. Sodium-glucose cotransporter 2 inhibitors and risk of retinopathy in patients with type 2 diabetes. *JAMA Netw Open* 2023; 6(12):e2348431. doi:10.1001/jamanetworkopen.2023.48431
5. Inzucchi SE, Wanner C, Hehnke U, et al. Retinopathy outcomes with empagliflozin versus placebo in the EMPA-REG OUTCOME trial. *Diabetes Care* 2019; 42(4):e53–e55. doi:10.2337/dc18-1355
6. Li C, Zhou Z, Neuen BL, et al. Sodium-glucose co-transporter-2 inhibition and ocular outcomes in patients with type 2 diabetes: a systematic review and meta-analysis. *Diabetes Obes Metab* 2021; 23(1):252–257. doi:10.1111/dom.14197
7. Marso SP, Bain SC, Consoli A, et al. Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2016; 375(19):1834–1844. doi:10.1056/NEJMoa1607141
8. Wang F, Mao Y, Wang H, Liu Y, Huang P. Semaglutide and diabetic retinopathy risk in patients with type 2 diabetes mellitus: a meta-analysis of randomized controlled trials. *Clin Drug Investig* 2022; 42(1):17–28. doi:10.1007/s40261-021-01110-w
9. Albert SG, Wood EM, Ahir V. Glucagon-like peptide 1-receptor agonists and A1c: good for the heart but less so for the eyes? *Diabetes Metab Syndr* 2023; 17(1):102696. doi:10.1016/j.dsx.2022.102696
10. Tsapas A, Avgerinos I, Karagiannis T, et al. Comparative effectiveness of glucose-lowering drugs for type 2 diabetes: a systematic review and network meta-analysis. *Ann Intern Med* 2020; 173(4):278–286. doi:10.7326/M20-0864
11. Bethel MA, Diaz R, Castellana N, Bhattacharya I, Gerstein HC, Lakshmanan MC. HbA1c change and diabetic retinopathy during GLP-1 receptor agonist cardiovascular outcome trials: a meta-analysis and meta-regression. *Diabetes Care* 2021; 44(1):290–296. doi:10.2337/dc20-1815
12. Husain M, Birkenfeld AL, Donsmark M, et al. Oral semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2019; 381(9):841–851. doi:10.1056/NEJMoa1901118
13. Yoshida Y, Joshi P, Barri S, et al. Progression of retinopathy with glucagon-like peptide-1 receptor agonists with cardiovascular benefits in type 2 diabetes—a systematic review and meta-analysis. *J Diabetes Complications* 2022; 36(8):108255. doi:10.1016/j.jdiacomp.2022.108255
14. Marso SP, Daniels GH, Brown-Frandsen K, et al. Liraglutide and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2016; 375(4):311–322. doi:10.1056/NEJMoa1603827
15. Bethel MA, Mentz RJ, Merrill P, et al. Microvascular and cardiovascular outcomes according to renal function in patients treated with once-weekly exenatide: insights from the EXSCEL Trial. *Diabetes Care* 2020; 43(2):446–452. doi:10.2337/dc19-1065
16. Wei J, Yang B, Wang R, et al. Risk of stroke and retinopathy during GLP-1 receptor agonist cardiovascular outcome trials: an eight RCTs meta-analysis. *Front Endocrinol (Lausanne)* 2022; 13:1007980. doi:10.3389/fendo.2022.1007980
17. Zhang X, Shao F, Zhu L, Ze Y, Zhu D, Bi Y. Cardiovascular and microvascular outcomes of glucagon-like peptide-1 receptor agonists in type 2 diabetes: a meta-analysis of randomized controlled cardiovascular outcome trials with trial sequential analysis. *BMC Pharmacol Toxicol* 2018; 19(1):58. doi:10.1186/s40360-018-0246-x
18. Gerstein HC, Colhoun HM, Dagenais GR, et al. Dulaglutide and cardiovascular outcomes in type 2 diabetes (REWIND): a double-blind, randomised placebo-controlled trial. *Lancet* 2019; 394(10193):121–130. doi:10.1016/S0140-6736(19)31149-3
19. Wai KM, Mishra K, Koo E, et al. Impact of GLP-1 agonists and SGLT-2 inhibitors on diabetic retinopathy progression: an aggregated electronic health record data study. *Am J Ophthalmol* 2024; 265:39–47. doi:10.1016/j.ajo.2024.04.010
20. American Diabetes Association Professional Practice Committee. 12. Retinopathy, neuropathy, and foot care: standards of care in diabetes—2024. *Diabetes Care* 2024; 47(suppl 1):S231–S243. doi:10.2337/dc24-S012

doi:10.3949/ccjm.91c.12002