

Noura Nachawi, MD

Department of Metabolism, Endocrinology & Diabetes, University of Michigan, Ann Arbor, MI; Assistant Professor, University of Michigan, Ann Arbor, MI

Dingfeng Li, MD

Department of Endocrinology, Diabetes, and Metabolism, Cleveland Clinic, Cleveland, OH; Assistant Professor, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University, Cleveland, OH

M. Cecilia Lansang, MD, MPH

Department of Endocrinology, Diabetes, and Metabolism, Cleveland Clinic, Cleveland, OH; Professor, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University, Cleveland, OH

Glucocorticoid-induced adrenal insufficiency and glucocorticoid withdrawal syndrome: Two sides of the same coin

ABSTRACT

Diseases of the adrenal glands can lead to primary adrenal insufficiency, and suppression of the hypothalamic-pituitary-adrenal axis can cause secondary adrenal insufficiency (adrenal suppression). The most common cause of adrenal suppression is exogenous steroids, a condition recently termed *glucocorticoid-induced adrenal insufficiency* (GIAI). Similarly, weaning from high doses of glucocorticoids or giving insufficient glucocorticoid replacement after curative surgery for endogenous hypercortisolism (Cushing syndrome) can lead to glucocorticoid withdrawal syndrome, which overlaps with GIAI.

KEY POINTS

GIAI is common in patients treated with glucocorticoids.

GIAI may go unrecognized when caused by non-oral formulations of glucocorticoids: intra-articular, epidural, inhaled, and even topical.

When tapering high doses of glucocorticoids, patients can develop symptoms of glucocorticoid withdrawal similar to those of GIAI.

Patients with GIAI are a vulnerable population with a poor baseline quality of life. Lack of awareness of GIAI among patients and physicians often leads to worse clinical outcomes and quality of life.

GLUCCORTICOID-INDUCED ADRENAL insufficiency (GIAI) is a well-known side effect of glucocorticoid therapy, and clinicians usually expect it in patients who receive systemic (oral, intravenous, and intramuscular) glucocorticoids in doses equivalent to more than 5 mg of prednisone for at least 3 weeks.¹ However, glucocorticoids given through other routes can also suppress the adrenal glands.

Unfamiliarity with GIAI, especially when caused by nonsystemic formulations of glucocorticoids, can lead to delay in diagnosis or misdiagnosis and lack of proper patient education. This lack of awareness often leads to failure to implement an adrenal action plan and underuse of injectable glucocorticoids at home or, in cases of adrenal crisis, in the emergency room.² Ultimately, gaps in care in managing adrenal suppression often worsen clinical outcomes and quality of life in this vulnerable patient population, who tend to have a poor quality of life at baseline.³

This review highlights the differences between primary adrenal insufficiency, secondary adrenal insufficiency (including GIAI), and glucocorticoid withdrawal syndrome.

■ DEFINITION AND TYPES OF ADRENAL INSUFFICIENCY

The adrenal cortex produces 3 main types of hormones⁴:

TABLE 1

Common causes of primary adrenal insufficiency

Autoimmune

Isolated

Autoimmune polyglandular syndrome type 1

Autoimmune polyglandular syndrome type 2

Adrenal infection

Tuberculosis

Human immunodeficiency virus

Cytomegalovirus

Fungal infections: candidiasis, histoplasmosis, paracoccidioidomycosis

Syphilis

African trypanosomiasis

Adrenal metastases

Breast, lung, colon, stomach cancers or lymphoma

Adrenal hemorrhage

Trauma

Anticoagulation

Antiphospholipid syndrome

Congenital adrenal hyperplasia

21-hydroxylase deficiency

11-hydroxylase deficiency

3B-hydroxysteroid dehydrogenase II deficiency

Drug-induced primary adrenal insufficiency

Adrenal enzyme inhibitors: mitotane, ketoconazole, metyrapone, etomidate

Drugs that accelerate cortisol metabolism: fluconazole, phenytoin, rifampin, barbiturates

Immune checkpoint inhibitors

Anti-PD-1 (programmed cell death protein 1) monoclonal antibodies: pembrolizumab, nivolumab

CTLA-4 (cytotoxic T-lymphocyte antigen 4) inhibitor: ipilimumab

Others

Adrenoleukodystrophy and adrenomyeloneuropathy

Familial glucocorticoid deficiency

Familial glucocorticoid resistance

- Glucocorticoids (primarily cortisol) from the zona fasciculata
- Mineralocorticoids (aldosterone, deoxycorticosterone) from the zona glomerulosa
- Androgens and their precursors (androstenedione, dihydroepiandrostenedione, dihydroepiandrosterone sulfate, testosterone, and 11-oxygenated 19-carbon androgens) from the zona reticularis.

Adrenal insufficiency is the inability of the adrenal cortex to synthesize and produce glucocorticoids, mineralocorticoids, or both.

Primary adrenal insufficiency

Diseases of the adrenal cortex can lead to primary adrenal insufficiency, with insufficient production of glucocorticoids, mineralocorticoids, or both. The prevalence of primary adrenal insufficiency in the United States is not well documented. However, it is rising in Europe, where it has been reported to be as high as 22.1 per 100,000 population.^{5,6}

Autoimmune adrenalitis (also known as Addison disease, for Thomas Addison,⁷ who first described it in 1855) is the most common cause of primary adrenal insufficiency in developed countries.⁸ Other causes include tuberculosis, human immunodeficiency virus infection, trauma, and use of immune checkpoint inhibitors (Table 1).

Secondary adrenal insufficiency

Secondary adrenal insufficiency occurs when the hypothalamus does not produce enough corticotropin-releasing hormone or the anterior pituitary gland does not produce enough adrenocorticotrophic hormone, so that the adrenal cortex is not stimulated and does not produce enough glucocorticoids. Mineralocorticoid secretion, however, is usually preserved, as the renin-angiotensin-aldosterone system, involving the cardiovascular and renal systems, is not affected.⁹ Therefore, patients with secondary adrenal insufficiency are less likely to have hypotension than those with primary adrenal insufficiency.

Adrenal insufficiency caused by suppressed corticotropin-releasing hormone is sometimes called *tertiary adrenal insufficiency*. However, this term remains controversial. Here, we will use *secondary adrenal insufficiency* for both pituitary and hypothalamic causes of adrenal insufficiency.

Secondary adrenal insufficiency is more common than primary, with an estimated prevalence of up to 28 per 100,000 people.¹⁰ Common causes include pituitary tumors, other tumors metastasizing to the pituitary gland, and head trauma (Table 2).

Other important causes are the many drugs that can affect the hypothalamic-pituitary-adrenal axis (HPAA) at different levels (Figure 1). The drugs that primary care clinicians most often encounter are immune checkpoint inhibitors, opioids, and glucocorticoids. Secondary adrenal insufficiency caused by emerging immunotherapies such as monoclonal antibody targeting programmed cell death protein 1 (PD-1; nivolumab and pembrolizumab) and monoclonal antibody targeting cytotoxic T-lymphocyte antigen 4 (CTLA-4; ipilimumab) is also common, more so when these agents are used in combination or sequence.¹¹ Of note, these

medicines can also cause primary adrenal insufficiency, though infrequently.¹²

Opioids are believed to suppress the adrenal glands by binding to receptors in the hypothalamus and pituitary, exerting tonic inhibition on the HPA. Opioid-induced adrenal insufficiency is estimated to affect approximately 15% of patients treated with opioids for at least 3 to 6 months.¹⁴ Li et al¹⁵ reported that 9 (9%) of 102 patients who were receiving more than 20 morphine milligram equivalents per day developed adrenal insufficiency. All were treated with glucocorticoid replacement while weaning off opioids until their HPA recovered, which occurred within 1 to 14 months of stopping the opioid. Glucocorticoid replacement improved pain, quality of life, and physical function.¹⁵

THE DEEP SLEEP OF ADRENAL GLANDS: ADRENAL SUPPRESSION AND GIAI

Glucocorticoids are powerful anti-inflammatory agents used to treat autoimmune and other conditions. However, long-term use in supraphysiologic doses can suppress the HPA and consequently cause GIAI.

GIAI is a fairly new term and has been used by some authors interchangeably with *adrenal suppression*.¹⁶ Other authors use the term more specifically to describe symptoms in patients with HPA suppression who receive inadequate treatment with glucocorticoids during stressful situations.¹⁷ The rest of the discussion will focus on GIAI, given that exogenous glucocorticoid use is the most common cause of adrenal suppression.

In a systematic review and meta-analysis of 73 studies, the median prevalence of GIAI was 37% in patients receiving any form of glucocorticoids.¹⁸ In another meta-analysis, the median prevalence was 48.7% in those receiving oral glucocorticoids and 52.2% in those receiving intra-articular injections.¹⁹

Excessive glucocorticoids, whether endogenous due to adrenal lesions secreting excessive cortisol or from an exogenous source, bind to receptors in the hypothalamus and pituitary, triggering negative feedback on adrenocorticotrophic hormone release. Chronic suppression of adrenocorticotrophic hormone eventually leads to atrophy of the zona fasciculata but not the zona glomerulosa, resulting in impaired cortisol secretion but intact mineralocorticoid secretion.²⁰

Risk factors for GIAI

Although high-quality evidence is lacking, available data suggest that many factors affect the risk of GIAI, including glucocorticoid dose, duration, formulation, frequency and timing of administration, pharmaco-

TABLE 2

Common causes of secondary adrenal insufficiency

Pituitary tumors

Pituitary tumors replacing normal corticotrophic cells
Adrenocorticotrophic hormone deficiency after tumor resection or radiation treatment

Nonpituitary tumors

Meningioma
Craniopharyngioma
Sellar or suprasellar metastases (lung, colon, and breast cancer)

Pituitary infiltration

Granulomatosis with polyangiitis
Sarcoidosis
Amyloidosis
Hemochromatosis
Lymphoma

Autoimmune

Lymphocytic hypophysitis
Isolated (usually with pregnancy)
Associated with other autoimmune disease (thyroid, vitiligo, type 1 diabetes, pernicious anemia)

Sheehan syndrome

Infarction in the pituitary gland due to excessive postpartum hemorrhage

Pituitary apoplexy

Acute hemorrhage in the pituitary adenoma

Head trauma

Severe head trauma leading to fracture of the skull base and injury in the pituitary gland

Drug-induced central adrenal insufficiency

See Figure 1

Rare congenital causes

Mutations of *TBX19* (T-box transcription factor 19) and *PCSK1* kexin (proprotein convertase subtilisin) genes
Mutations of *POMC* (proopiomelanocortin) gene

kinetics, interaction with other medications, and cushingoid features.¹⁶

Glucocorticoid dose and duration. In studies in patients with asthma,¹⁹ GIAI occurred in 2.4% of those treated with low doses of systemic glucocorticoids, 8.5% of those receiving medium doses, and 21.5% of those receiving high doses. Short-term use (< 1 month) resulted in GIAI in 1.4%, medium-term use (1 month to 1 year) resulted in GIAI in 11.9%, and long-term use (> 1 year) resulted in GIAI in 27.4%. The patterns were similar in patients treated only with inhaled glucocorticoids. However, other studies have found no correlations between glucocorticoid dose or duration and risk of GIAI.¹⁸




Figure 1. How various drugs can cause secondary adrenal insufficiency.

CTLA-4 = cytotoxic T-lymphocyte antigen 4; PD-1 = programmed cell death protein 1

TABLE 3
Risk factors for glucocorticoid-induced adrenal insufficiency

Route of administration	Reported risk	Factors that increase the risk	Factors that decrease the risk
Inhaled	Dose- and duration-dependent ⁴¹ 20.3% in patients treated for > 1 year ¹⁹	High doses (any glucocorticoid > 0.8 mg/day or fluticasone propionate > 0.75 mg/day) ⁴² Concurrent use of intranasal or oral glucocorticoids ^{43,44} Use of spacer device to deliver more medication to the lower airways ⁴⁵ Higher lung volumes ⁴⁵	Beclomethasone dipropionate, budesonide, and triamcinolone acetonide are less likely to suppress the HPA axis compared with fluticasone propionate ⁴² Ciclesonide has the lowest risk of HPA axis suppression ⁴⁶ Lower lung volumes ⁴⁵
Intranasal	Low ($\leq 4.2\%$) ^{19,46,47}	Long-term use (> 12 months) ⁴⁶	Short-term use ⁴⁶
Intra-articular injections	52.2% ¹⁹ GIAI usually occurs 1 to 8 weeks after injection ⁴⁸ After single and repeated injections ⁴⁹	Higher doses ⁴⁹ Patients with inflammatory disease ⁴⁹ Administration in bilateral joints simultaneously ⁴⁸	Patients with degenerative disease ⁴⁹
Epidural injections	52.2% ¹⁹	Higher doses ⁵⁰ Longer-acting glucocorticoids (eg, methylprednisolone, triamcinolone) ⁵¹	Lower doses ⁵⁰ Shorter-acting glucocorticoids ⁵¹
Topical	4.7% ^{19,52} Shampoo formulations are not linked to GIAI ⁵³	Disruption of skin barrier ^{52,54} Long-term use (> 12 months) ^{52,55} Higher-potency topical glucocorticoids (eg, betamethasone dipropionate, clobetasol propionate) ^{52,55,56} Higher doses Application to larger body surface ^{52,54} Use of occlusive bandage ^{52,54} Application on the eyelids, scrotum, and mucosal surfaces ^{52,54}	Lower-potency topical glucocorticoids (eg, dexamethasone cream 0.1%, hydrocortisone 0.5%, hydrocortisone 1%, hydrocortisone 2.5%, methylprednisolone 1%) ⁵⁵

GIAI = glucocorticoid-induced adrenal insufficiency; HPA axis = hypothalamic-pituitary-adrenal axis

Based on information in references 19 and 41–56.

Formulation. Dexamethasone is 25 times more potent than hydrocortisone, and prednisone is 4 times more potent. Duration of effect is more than 36 hours for dexamethasone, 18 to 36 hours for prednisone, and 8 to 12 hours for hydrocortisone.^{21,22} At equivalent doses (0.75 mg of dexamethasone is equivalent to 5 mg of prednisone or 20 mg of hydrocortisone),^{21,22} dexamethasone has stronger suppressive effects on the HPA axis compared with hydrocortisone. However, studies have not shown any difference in HPA axis suppression in patients treated with equivalent doses of prednisone compared with dexamethasone.^{23,24}

Frequency and timing of administration. Pulse therapy with high-dose glucocorticoids (eg, intravenous methylprednisolone 250–500 mg weekly for 6–12 weeks)^{25,26} and alternate single-day dosing are less likely to cause GIAI^{27,28} compared with bedtime dosing and frequent dosing (more than once daily).^{29–31}

GIAI after short bursts of glucocorticoids (7–14 days) has been infrequently reported,^{28,32,33} particularly in patients with chronic obstructive pulmonary disease who receive frequent short bursts of glucocorticoids³⁴ and patients with malignancies who receive bursts of dexamethasone to mitigate chemotherapy-related nausea.³⁵

Interactions with other medications. Concomitant use of glucocorticoids and hepatic cytochrome P450 3A4 inhibitors (eg, protease inhibitors, azoles, clarithromycin, erythromycin) increases the levels of active metabolites of glucocorticoids, and consequently, the risk of GIAI.^{36,37} This happens with all glucocorticoid formulations metabolized by cytochrome P450 3A4 regardless of route of administration: oral, injectable, intra-articular, and even inhaled and intranasal formulations.^{38–40} Primary care clinicians should be aware of these interactions when

they suspect GIAI, especially in patients with chronic obstructive pulmonary disease or human immunodeficiency virus infection.

Cushingoid features. A cushingoid appearance usually indicates that the glucocorticoid dose is excessive. Some authors have indicated that patients with cushingoid features while on glucocorticoids are at a very high risk for GIAI.¹⁶

Unrecognized sources of exogenous glucocorticoids

Inhaled glucocorticoids bind to receptors in the lungs, mouth, and oropharynx, leading to systemic exposure and possibly HPA suppression.²¹ Table 3 summarizes the reported risk of GIAI after exposure to the different formulations and factors that can increase or decrease the risk.^{19,41–56}

Intra-articular and epidural injections. Systemic absorption of intra-articular glucocorticoids has been widely described.^{57,58} Similarly, HPA suppression after epidural corticosteroid injections has been reported with multiple formulations, doses, and frequencies (after both single and recurrent doses).⁵⁹

Some patients do not know that these injections contain steroids and therefore may not report this exposure if they present with GIAI symptoms.⁶⁰ Serum and urine testing for synthetic steroids are important tools when GIAI is suspected.⁶¹ Urine screening for synthetic glucocorticoids (liquid chromatography-tandem mass spectrometry with stable isotope dilution analysis) is reported to detect prednisone and prednisolone for up to 40 days after epidural injections and for up to 62 days after triamcinolone epidural injections.⁶⁰

Topical formulations. Several studies reported GIAI induced by topical cutaneous glucocorticoids.^{19,52}

Eye drops. GIAI due to ophthalmic glucocorticoids has been reported in adult, pediatric, and animal studies.^{62,63}

Locally active enteral formulations. Rectal glucocorticoids and oral budesonide are used to treat inflammatory bowel disease. The risk of GIAI is dose- and duration-dependent in patients taking oral budesonide, being higher when patients take more than 6 mg daily for at least 8 weeks.⁶⁴ GIAI has been reported in patients using prednisolone enemas,⁶⁵ whereas beclomethasone dipropionate enemas seem to be safer.⁶⁶

Other medications with glucocorticoid activity

Megestrol acetate is a synthetic progestin with glucocorticoid-like activity commonly used as an appetite stimulant in patients with malignancy and anorexia. Several reports have highlighted the incidence of adrenal insufficiency, Cushing syndrome, or both in patients

treated with megestrol acetate,^{67,68} specifically, when megestrol acetate is combined with dexamethasone.³⁵

Medroxyprogesterone acetate, another progestin that binds glucocorticoid receptors,^{69,70} is used to treat endometrial cancer, endometriosis, and abnormal uterine bleeding, and as a contraceptive, and is reported to cause HPA suppression.⁷¹

GLUCOCORTICOID WITHDRAWAL SYNDROME

Excessive endogenous hormone secretion or exogenous administration often leads to tolerance (decreased response to the elevated hormones and the need for even higher levels to achieve the same effect) followed by physiologic and psychologic dependence.¹ In this situation, gradually tapering or abruptly stopping the glucocorticoids can induce glucocorticoid withdrawal syndrome,¹ even while patients are still receiving supraphysiologic doses of glucocorticoids.

Glucocorticoid withdrawal syndrome manifests as a spectrum of nonspecific symptoms and is mediated by multiple mechanisms. Chronic suppression of corticotropin-releasing hormone after stopping or tapering from glucocorticoids leads to adrenal insufficiency, adrenal crisis, depressive mood changes,⁷² hypersomnia, and lethargy.^{73,74} Prolonged suppression of proopiomelanocortin-related peptides causes myalgia, arthralgia, fever, and headache.¹ Depressed central noradrenergic and dopaminergic systems cause nonspecific withdrawal symptoms along with anorexia, nausea, vomiting, and weight loss.¹ Loss of glucocorticoid's suppressive effect on calcium absorption results in hypercalcemia and hyperphosphatemia.¹

These symptoms can develop at any time—during glucocorticoid taper (while the patient is still on supraphysiologic doses), after completely stopping glucocorticoids, and even after there is biochemical evidence of HPA recovery.¹

Long-term treatment with supraphysiologic doses of glucocorticoids often leads to HPA suppression and adrenal insufficiency. At the same time, tolerance to and dependence on high doses of glucocorticoids causes glucocorticoid withdrawal syndrome when attempting to taper or discontinue these drugs. Therefore, adrenal insufficiency and glucocorticoid withdrawal syndrome share similar clinical features (Table 4); however, they are completely different clinical entities that often overlap until the HPA recovers. Results of biochemical testing including early morning cortisol levels and the corticotropin stimulation test can be normal or suboptimal, and hence, not helpful in making this diagnosis.⁷⁵

TABLE 4

Adrenal insufficiency, glucocorticoid-induced adrenal insufficiency, and glucocorticoid withdrawal syndrome

	Adrenal insufficiency	Glucocorticoid-induced adrenal insufficiency	Glucocorticoid withdrawal syndrome
Diagnosis	Clinical symptoms and biochemical testing: Low 8 AM cortisol (< 4.8 µg/dL) ^a Abnormal response to corticotropin stimulation test (cortisol peak < 12.6 µg/dL at 30 minutes and 60 minutes) ^a Variable adrenocorticotrophic hormone (for primary adrenal insufficiency > 63.3 pg/mL, for secondary adrenal insufficiency < 7.2 pg/mL) ^b	After abrupt discontinuation or quick taper of exogenous glucocorticoid or Cushing syndrome: Low 8 AM cortisol (< 4.8 µg/dL) ^a Low adrenocorticotrophic hormone (< 7.2 pg/mL) ^b Low dehydroepiandrosterone sulfate ^c Abnormal response to corticotropin stimulation test (cortisol peak < 12.6 µg/dL at 30 and 60 minutes) ^a	Clinical symptoms of adrenal insufficiency with or without cushingoid features while gradually tapering or after abrupt discontinuation of glucocorticoid No laboratory test to diagnose
Mechanism	Lack of glucocorticoid secretion from adrenal cortex due to either adrenal etiology (primary adrenal insufficiency) or pituitary or hypothalamic etiology (secondary adrenal insufficiency)	HPAA suppression due to excessive endogenous or exogenous glucocorticoid, leading to atrophy of adrenal cortex	Tolerance of and dependence on supraphysiologic doses of glucocorticoid
Prevention	Replace with physiologic doses of glucocorticoid	Gradually taper glucocorticoid until completely stopped	Use the lowest effective supraphysiologic glucocorticoid dose when indicated
Treatment	Replace with physiologic doses of glucocorticoid	Gradually taper glucocorticoid until completely stopped Consider stress-dose glucocorticoid under stressors	No effective treatment: empirically increase glucocorticoid to prolong HPAA suppression

^aCortisol values per the Elecsys Cortisol II assay.^bAdrenocorticotrophic hormone values per the Electro Chemiluminescence Immunoassay.^cDehydroepiandrosterone sulfate normal values (µg/dL) per the Electro Chemiluminescence Immunoassay for females, by age:

15–19 years 65.1–368.0; 20–24 years 148–407; 25–34 years 98.8–340; 35–44 years 60.9–337; 45–54 years 35.4–256; 55–64 years 18.9–205; 65–74 years < 247; 75–99 years 12–154.

For males, by age:

15–19 years 70.2–492; 20–24 years 211–492; 25–34 years 160–449; 35–44 years 88.9–427; 45–54 years 44.3–331; 55–64 years 51.7–295; 65–74 years 33.6–249; 75–99 years 16.2–123.

HPAA = hypothalamic-pituitary-adrenal axis

Glucocorticoid withdrawal syndrome after successful treatment of Cushing syndrome

Evidence on glucocorticoid withdrawal syndrome in patients with GIAI caused by exogenous glucocorticoid use is lacking. However, several studies have looked into glucocorticoid withdrawal syndrome in patients with GIAI caused by adrenal lesions secreting excessive endogenous cortisol (adrenocorticotrophic hormone-independent Cushing syndrome). Up to 99% of patients with Cushing syndrome have HPAA suppression.⁷⁶ Patients with Cushing syndrome can develop tolerance to and dependence on excessive endogenous cortisol, and hence, suffer from glucocorticoid withdrawal

syndrome postoperatively.⁷⁶ After resection, glucocorticoid taper is indicated until the HPAA recovers.

Postoperative glucocorticoid withdrawal syndrome is usually characterized by biochemical evidence of HPAA suppression, with many signs and symptoms consistent with cortisol deficiency despite the use of supraphysiologic doses of glucocorticoids. Common symptoms include myalgias, arthralgias, fatigue, weakness, sleep disturbance, and mood changes. In a recent prospective observational study, myalgias, arthralgias, and weakness got progressively worse 5 to 12 weeks after surgery.⁷⁷ Glucocorticoid withdrawal syndrome can be difficult to differentiate from adrenal

TABLE 5

Approach to glucocorticoid taper in patients with glucocorticoid-induced adrenal insufficiency and after surgery for Cushing syndrome

Average daily prednisone dose

> 40 mg/day: decrease by 10 mg weekly until 40 mg daily
 20–40 mg/day: decrease by 5 mg weekly until 20 mg daily
 10–20 mg/day: decrease by 1–2.5 mg weekly until 10 mg daily
 5–10 mg/day: decrease by 1 mg weekly until < 5 mg daily
 < 5 mg/day: switch to equivalent dose of hydrocortisone (eg, 10 mg in the morning and 5 mg in the early afternoon); hold hydrocortisone for 24 hours and retest HPA

Testing for HPA recovery

If patient has been on prednisone 5 mg/day, switch to equivalent dose of hydrocortisone, wait for 2–4 weeks, and hold hydrocortisone for 24 hours before testing

Check 8 AM serum cortisol:

If < 10 µg/dL,^a continue current dose of hydrocortisone and retest in 4–8 weeks

If ≥ 10 µg/dL, perform 250-µg corticotropin stimulation test:

- If suboptimal (cortisol peak < 12.6 µg/dL at 30 minutes and 60 minutes), consider stopping daily glucocorticoid replacement if patient has no withdrawal symptoms, but continue the sick-day rule (using stress-dose glucocorticoid) until repeating corticotropin stimulation test
 - If optimal (peak cortisol ≥ 12.6 µg/dL), stop glucocorticoid if patient is comfortable
- If 8 AM serum cortisol ≥ 12.6 µg/dL, consider stopping glucocorticoid if patient is ready in terms of withdrawal symptoms, or performing 250-µg corticotropin stimulation test or tapering glucocorticoid dose

Frequency of testing:

- If the results of tests are abnormal, recheck every 2–3 months
- If no recovery within 1 year, reassess every 3–6 months

Things to consider

- If glucocorticoid withdrawal syndrome develops at any point, increase the glucocorticoid dose to the most recent dose on which the patient did not have glucocorticoid withdrawal syndrome; consider decrements every other week rather than weekly
- If patient is on dexamethasone, consider switching to prednisone
- If patient is on twice-daily prednisone dosing, consider switching to equivalent dose of prednisone in the morning once daily

^aValues per the Elecsys Cortisol II assay.

HPAA = hypothalamic-pituitary-adrenal axis

Based in part on information in reference 81.

insufficiency, which complicates glucocorticoid dosing and tapering regimens.

In a retrospective study of the postoperative course of 81 patients with adrenocorticotrophic hormone-independent Cushing syndrome,⁷⁸ glucocorticoid withdrawal syndrome was most common when the 8 AM serum cortisol level 24 hours after the last glucocorticoid dose was less than 5 µg/dL, whereas no withdrawal symptoms were reported when it was higher than 10 µg/dL.

ASSESSING AND EXPEDITING HPA RECOVERY IN GIAI

Currently, there is no consensus on the best approach to assessing HPA recovery in patients with GIAI and those who have undergone surgery for Cushing syndrome. However, several factors related to the patient's characteristics, glucocorticoid course of therapy, and biochemical testing could be used to estimate the

recovery of the HPA and help clinicians with their approach to patients with GIAI.

Studies have looked at recovery of the HPA after successful surgery for endogenous adrenocorticotrophic hormone-independent Cushing syndrome, and we could extrapolate some of their conclusions to GIAI.⁷⁸ Slower HPA recovery is expected in patients treated with higher doses of glucocorticoids, women, patients with lower body mass index, and patients with cushingoid features. Faster recovery (in weeks to months) is reported in patients treated with high doses of oral glucocorticoids for less than 1 month.¹⁹ HPA recovery could take up to 6 to 12 months in patients treated with glucocorticoids for more than 12 months.^{19,79} Future studies are needed to prove the hypothesis.

An observational study by Pofi et al⁷⁹ involving 776 patients suggested a cutoff of 3.6 µg/dL (using the Roche Modular System) between baseline cortisol and

30-minute cortisol levels after a 250- μ g corticotropin stimulation test to predict recovery of the HPA. If the change in cortisol level is less than 3.6 μ g/dL and the random cortisol level is less than 7.2 μ g/dL after 1 year, patients are less likely to recover HPA function.⁷⁹

Switching from a longer-acting glucocorticoid (eg, dexamethasone, prednisone) to a shorter-acting one (eg, hydrocortisone) has been hypothesized to expedite HPA recovery, but evidence remains inadequate to recommend one glucocorticoid vs others for HPA recovery.^{16,80}

Corticosteroid taper in patients with GIAI and after surgery for Cushing syndrome

Clinicians should work in multidisciplinary teams and closely monitor conditions that could possibly worsen or relapse due to lowering glucocorticoid doses. Glucocorticoids should be tapered when appropriate to safely induce HPA recovery while at the same time avoiding glucocorticoid withdrawal syndrome or adrenal crisis.

Based on available literature and expert opinion,⁸¹ we suggest the approach to tapering glucocorticoids in patients with GIAI outlined in Table 5.

REFERENCES

- Hochberg Z, Pacak K, Chrousos GP. Endocrine withdrawal syndromes. *Endocr Rev* 2003; 24(4):523–538. doi:10.1210/er.2001-0014
- Li D, Genere N, Behnken E, et al. Determinants of self-reported health outcomes in adrenal insufficiency: a multisite survey study. *J Clin Endocrinol Metab* 2021; 106(3):e1408–e1419. doi:10.1210/clinem/dgaa668
- Li D, Brand S, Hamidi O, et al. Quality of life and its determinants in patients with adrenal insufficiency: a survey study at 3 centers in the United States. *J Clin Endocrinol Metab* 2022; 107(7):e2851–e2861. doi:10.1210/clinem/dgac175
- Melmed S, Polonsky KS, Larsen PR, Kronenberg HM, eds. *Williams Textbook of Endocrinology*. 13th ed. Bucharest, Romania: Elsevier; 2016.
- Olafsson AS, Sigurjonsdottir HA. Increasing prevalence of Addison disease: results from a nationwide study. *Endocr Pract* 2016; 22(1):30–35. doi:10.4158/EP15754.0R
- Løvås K, Husebye ES. High prevalence and increasing incidence of Addison's disease in western Norway. *Clin Endocrinol (Oxf)* 2002; 56(6):787–791. doi:10.1046/j.1365-2265.2002.t01-1-01552.x
- Addison T. On the constitutional and local effects of disease of the supra-renal capsules / by Thomas Addison. London, UK: Wellcome Collection; 1855. <https://wellcomecollection.org/works/xsmzqpdw>. Accessed March 14, 2024.
- Erichsen MM, Løvås K, Skiningsrud B, et al. Clinical, immunological, and genetic features of autoimmune primary adrenal insufficiency: observations from a Norwegian registry. *J Clin Endocrinol Metab* 2009; 94(12):4882–4890. doi:10.1210/jc.2009-1368
- Bancos I, Hahner S, Tomlinson J, Arlt W. Diagnosis and management of adrenal insufficiency. *Lancet Diabetes Endocrinol* 2015; 3(3):216–226. doi:10.1016/S2213-8587(14)70142-1
- Arlt W, Allolio B. Adrenal insufficiency. *Lancet* 2003; 361(9372):1881–1893. doi:10.1016/S0140-6736(03)13492-7
- Iglesias P, Sánchez JC, Díez JJ. Isolated ACTH deficiency induced by cancer immunotherapy: a systematic review. *Pituitary* 2021; 24(4):630–643. doi:10.1007/s11102-021-01141-8
- Grouthier V, Lebrun-Vignes B, Moey M, et al. Immune checkpoint inhibitor-associated primary adrenal insufficiency: WHO Vigibase Report analysis. *Oncologist* 2020; 25(8):696–701. doi:10.1634/theoncologist.2019-0555

TAKE-HOME POINTS

Primary care clinicians should be aware of the high incidence of GIAI in patients who are treated with formulations of glucocorticoids other than oral forms. Glucocorticoid withdrawal syndrome develops due to dependence on supraphysiologic doses. Its symptoms closely resemble those of GIAI.

Primary care clinicians are encouraged to taper glucocorticoids when possible and test for HPA recovery. If patients develop symptoms of glucocorticoid withdrawal syndrome while tapering, clinicians could consider increasing the glucocorticoid dose slightly and reattempting a slower taper.

DISCLOSURES

Dr. Li has disclosed conducting research for BridgeBio Pharma. Dr. Lansang has disclosed receiving research funding support from Abbott, serving as a research principal investigator for Abbott and Dexcom, and conducting research for Xeris. Dr. Nachawi reports no relevant financial relationships which, in the context of their contributions, could be perceived as a potential conflict of interest.

- Donegan D, Bancos I. Opioid-induced adrenal insufficiency. *Mayo Clin Proc* 2018; 93(7):937–944. doi:10.1016/j.mayocp.2018.04.010
- de Vries F, Bruin M, Lobatto DJ, et al. Opioids and their endocrine effects: a systematic review and meta-analysis. *J Clin Endocrinol Metab* 2020; 105(3):1020–1029. doi:10.1210/clinem/dgcz022
- Li T, Cunningham JL, Gilliam WP, Loukianova L, Donegan DM, Bancos I. Prevalence of opioid-induced adrenal insufficiency in patients taking chronic opioids. *J Clin Endocrinol Metab* 2020; 105(10):e3766–e3775. doi:10.1210/clinem/dgaa499
- Prete A, Bancos I. Glucocorticoid induced adrenal insufficiency [published correction appears in *BMJ* 2021; 374:n1936]. *BMJ* 2021; 374:n1380. doi:10.1136/bmj.n1380
- Borresen SW, Klose M, Glinthorp D, Watt T, Andersen MS, Feldt-Rasmussen U. Approach to the patient with glucocorticoid-induced adrenal insufficiency. *J Clin Endocrinol Metab* 2022; 107(7):2065–2076. doi:10.1210/clinem/dgac151
- Joseph RM, Hunter AL, Ray DW, Dixon WG. Systemic glucocorticoid therapy and adrenal insufficiency in adults: a systematic review. *Semin Arthritis Rheum* 2016; 46(1):133–141. doi:10.1016/j.semarthrit.2016.03.001
- Broersen LH, Pereira AM, Jørgensen JO, Dekkers OM. Adrenal insufficiency in corticosteroids use: systematic review and meta-analysis. *J Clin Endocrinol Metab* 2015; 100(6):2171–2180. doi:10.1210/jc.2015-1218
- Crowley RK, Argese N, Tomlinson JW, Stewart PM. Central hypoadrenalism. *J Clin Endocrinol Metab* 2014; 99(11):4027–4036. doi:10.1210/jc.2014-2476
- Williams DM. Clinical pharmacology of corticosteroids. *Respir Care* 2018; 63(6):655–670. doi:10.4187/respcare.06314
- Meikle AW, Tyler FH. Potency and duration of action of glucocorticoids. Effects of hydrocortisone, prednisone and dexamethasone on human pituitary-adrenal function. *Am J Med* 1977; 63(2):200–207. doi:10.1016/0002-9343(77)90233-9
- Kuperman H, Odone Filho V, Cristofani LM, Assis de Almeida MT, Setian N, Damiani D. Evaluation of adrenal reserve in children with acute lymphocytic leukemia treated with prednisone or dexamethasone. *Horm Res Paediatr* 2012; 78(2):73–80. doi:10.1159/000339830

24. Grossmann C, Scholz T, Rochel M, et al. Transactivation via the human glucocorticoid and mineralocorticoid receptor by therapeutically used steroids in CV-1 cells: a comparison of their glucocorticoid and mineralocorticoid properties. *Eur J Endocrinol* 2004; 151(3):397–406. doi:10.1530/eje.0.1510397
25. Jespersen S, Nygaard B, Kristensen LØ. Methylprednisolone pulse treatment of Graves' ophthalmopathy is not associated with secondary adrenocortical insufficiency. *Eur Thyroid J* 2015; 4(4): 222–225. doi:10.1159/000440834
26. Giotaki Z, Fountas A, Tsirouki T, Bargiota A, Tigas S, Tzatsoulis A. Adrenal reserve following treatment of Graves' orbitopathy with intravenous glucocorticoids. *Thyroid* 2015; 25(4):462–463. doi:10.1089/thy.2014.0533
27. Ackerman GL, Nolsn CM. Adrenocortical responsiveness after alternate-day corticosteroid therapy. *N Engl J Med* 1968; 278(8):405–409. doi:10.1056/NEJM196802222780801
28. Carter ME, James VH. Effect of alternate-day, single-dose, corticosteroid therapy on pituitary-adrenal function. *Ann Rheum Dis* 1972; 31(5):379–383. doi:10.1136/ard.31.5.379
29. Jasani MK, Boyle JA, Greig WR, et al. Corticosteroid-induced suppression of the hypothalamo-pituitary-adrenal axis: observations on patients given oral corticosteroids for rheumatoid arthritis. *Q J Med* 1967; 36(143):261–276. PMID:6049762
30. Nichols T, Nugent CA, Tyler FH. Diurnal variation in suppression of adrenal function by glucocorticoids. *J Clin Endocrinol Metab* 1965; 25:343–349. doi:10.1210/jcem-25-3-343
31. Grant SD, Forsham PH, DiRaimondo VC. Suppression of 17-hydroxycorticosteroids in plasma and urine by single and divided doses of triamcinolone. *N Engl J Med* 1965; 273(21):1115–1118. doi:10.1056/NEJM196511182732101
32. Brigell DF, Fang VS, Rosenfield RL. Recovery of responses to ovine corticotropin-releasing hormone after withdrawal of a short course of glucocorticoid. *J Clin Endocrinol Metab* 1992; 74(5):1036–1039. doi:10.1210/jcem.74.5.1314844
33. Lević Z, Micić D, Nikolić J, et al. Short-term high dose steroid therapy does not affect the hypothalamic-pituitary-adrenal axis in relapsing multiple sclerosis patients. Clinical assessment by the insulin tolerance test. *J Endocrinol Invest* 1996; 19(1):30–34. doi:10.1007/BF03347855
34. Fleishaker DL, Mukherjee A, Whaley FS, Daniel S, Zeiher BG. Safety and pharmacodynamic dose response of short-term prednisone in healthy adult subjects: a dose ranging, randomized, placebo-controlled, crossover study. *BMC Musculoskelet Disord* 2016; 17:293. doi:10.1186/s12891-016-1135-3
35. Han HS, Park JC, Park SY, et al. A prospective multicenter study evaluating secondary adrenal suppression after antiemetic dexamethasone therapy in cancer patients receiving chemotherapy: a Korean South West Oncology Group study. *Oncologist* 2015; 20(12):1432–1439. doi:10.1634/theoncologist.2015-0211
36. Lebrun-Vignes B, Archer VC, Diquet B, et al. Effect of itraconazole on the pharmacokinetics of prednisolone and methylprednisolone and cortisol secretion in healthy subjects. *Br J Clin Pharmacol* 2001; 51(5):443–450. doi:10.1046/j.1365-2125.2001.01372.x
37. Busse KH, Formentini E, Alfaro RM, Kovacs JA, Penzak SR. Influence of antiretroviral drugs on the pharmacokinetics of prednisolone in HIV-infected individuals. *J Acquir Immune Defic Syndr* 2008; 48(5):561–566. doi:10.1097/QAI.0b013e31817bebeb
38. Yombi JC, Maiter D, Belkhir L, Nzeusseu A, Vandercam B. Iatrogenic Cushing's syndrome and secondary adrenal insufficiency after a single intra-articular administration of triamcinolone acetonide in HIV-infected patients treated with ritonavir. *Clin Rheumatol* 2008; 27(suppl 2):S79–S82. doi:10.1007/s10067-008-1022-x
39. Foisy MM, Yakiwchuk EM, Chiu I, Singh AE. Adrenal suppression and Cushing's syndrome secondary to an interaction between ritonavir and fluticasone: a review of the literature. *HIV Med* 2008; 9(6):389–396. doi:10.1111/j.1468-1293.2008.00579.x
40. Saberi P, Phengrasamy T, Nguyen DP. Inhaled corticosteroid use in HIV-positive individuals taking protease inhibitors: a review of pharmacokinetics, case reports and clinical management. *HIV Med* 2013; 14(9):519–529. doi:10.1111/hiv.12039
41. Lasky-Su J. Inhaled corticosteroid use for asthma is linked to adrenal suppression. *Nat Med* 2022; 28(4):645–646. doi:10.1038/s41591-022-01732-3
42. Lipworth BJ. Systemic adverse effects of inhaled corticosteroid therapy: a systematic review and meta-analysis. *Arch Intern Med* 1999; 159(9):941–955. doi:10.1001/archinte.159.9.941
43. Zöllner EW, Lombard C, Galal U, Hough S, Irusen E, Weinberg E. Hypothalamic-pituitary-adrenal axis suppression in asthmatic children on inhaled and nasal corticosteroids—more common than expected? *J Pediatr Endocrinol Metab* 2011; 24(7–8):529–534. doi:10.1515/jpem.2011.198
44. Zöllner EW, Lombard CJ, Galal U, Hough FS, Irusen EM, Weinberg E. Hypothalamic-pituitary-adrenal axis suppression in asthmatic children. *Pediatrics* 2012; 130(6):e1512–1519. doi:10.1542/peds.2012-1147
45. Lipworth B, Kuo C, Jabbal S. Adrenal suppression with inhaled corticosteroids: the seed and the soil. *Lancet Respir Med* 2018; 6(6):e19. doi:10.1016/S2213-2600(18)30148-6
46. Derendorf H, Nave R, Drollmann A, Cerasoli F, Wurst W. Relevance of pharmacokinetics and pharmacodynamics of inhaled corticosteroids to asthma. *Eur Respir J* 2006; 28(5):1042–1050. doi:10.1183/09031936.00074905
47. Sampieri G, Namavarian A, Lee JJW, Hamour AF, Lee JM. Hypothalamic-pituitary-adrenal axis suppression and intranasal corticosteroid use: a systematic review and meta-analysis. *Int Forum Allergy Rhinol* 2022; 12(1):11–27. doi:10.1002/alr.22863
48. Habib G, Khazin F, Jabbour A, et al. Simultaneous bilateral knee injection of methylprednisolone acetate and the hypothalamic-pituitary adrenal axis: a single-blind case-control study. *J Invest Med* 2014; 62(3):621–626. doi:10.2310/JIM.0000000000000048
49. Mader R, Lavi I, Luboshitzky R. Evaluation of the pituitary-adrenal axis function following single intra-articular injection of methylprednisolone. *Arthritis Rheum* 2005; 52(3):924–928. doi:10.1002/art.20884
50. Sim SE, Hong HJ, Roh K, Seo J, Moon HS. Relationship between epidural steroid dose and suppression of the hypothalamus-pituitary-adrenal axis. *Pain Physician* 2020; 23(4S):S283–S294. PMID:32942788
51. Friedly JL, Comstock BA, Heagerty PJ, et al. Systemic effects of epidural steroid injections for spinal stenosis. *Pain* 2018; 159(5): 876–883. doi:10.1097/j.pain.0000000000001158
52. Böckle BC, Jara D, Nindl W, Aberer W, Sepp NT. Adrenal insufficiency as a result of long-term misuse of topical corticosteroids. *Dermatology* 2014; 228(4):289–293. doi:10.1159/000358427
53. Andres P, Poncet M, Farzaneh S, Soto P. Short-term safety assessment of clobetasol propionate 0.05% shampoo: hypothalamic-pituitary-adrenal axis suppression, atrophogenicity, and ocular safety in subjects with scalp psoriasis. *J Drugs Dermatol* 2006; 5(4):328–332. PMID:16673799
54. Hengge UR, Ruzicka T, Schwartz RA, Cork MJ. Adverse effects of topical glucocorticosteroids. *J Am Acad Dermatol* 2006; 54(1):1–15. doi:10.1016/j.jaad.2005.01.010
55. Weston WL, Fennessey PV, Morelli J, et al. Comparison of hypothalamic-pituitary-adrenal axis suppression from superpotent topical steroids by standard endocrine function testing and gas chromatographic mass spectrometry. *J Invest Dermatol* 1988; 90(4):532–535. doi:10.1111/1523-1747.ep12461062
56. Walsh P, Aeling JL, Huff L, Weston WL. Hypothalamus-pituitary-adrenal axis suppression by superpotent topical steroids. *J Am Acad Dermatol* 1993; 29(3):501–503. doi:10.1016/s0190-9622(08)82011-7
57. Armstrong RD, English J, Gibson T, Chakraborty J, Marks V. Serum methylprednisolone levels following intra-articular injection of methylprednisolone acetate. *Ann Rheum Dis* 1981; 40(6):571–574. doi:10.1136/ard.40.6.571
58. Derendorf H, Möllmann H, Grüner A, Haack D, Gyselby G. Pharmacokinetics and pharmacodynamics of glucocorticoid suspensions after intra-articular administration. *Clin Pharmacol Ther* 1986; 39(3):313–317. doi:10.1038/clpt.1986.45
59. Lee MS, Moon HS. Safety of epidural steroids: a review. *Anesth Pain Med (Seoul)* 2021; 16(1):16–27. doi:10.17085/apm.21002
60. Lansang MC, Farmer T, Kennedy L. Diagnosing the unrecognized systemic absorption of intra-articular and epidural steroid injections. *Endocr Pract* 2009; 15(3):225–228. doi:10.4158/EP.15.3.225

61. Cizza G, Nieman LK, Doppman JL, et al. Factitious Cushing syndrome. *J Clin Endocrinol Metab* 1996; 81(10):3573–3577. doi:10.1210/jcem.81.10.8855803
62. Kröger L, Kotaniemi K, Jääskeläinen J. Topical treatment of uveitis resulting in adrenal insufficiency. *Acta Paediatr* 2009; 98(3): 584–585. doi:10.1111/j.1651-2227.2008.01091.x
63. Roberts SM, Lavach JD, Macy DW, Severin GA. Effect of ophthalmic prednisolone acetate on the canine adrenal gland and hepatic function. *Am J Vet Res* 1984; 45(9):1711–1714. PMID:6497127
64. Löfberg R, Rutgeerts P, Malchow H, et al. Budesonide prolongs time to relapse in ileal and ileocaecal Crohn's disease. A placebo controlled one year study. *Gut* 1996; 39(1):82–86. doi:10.1136/gut.39.1.82
65. Luman W, Gray RS, Pendek R, Palmer KR. Prednisolone metasulphobenzoate foam retention enemas suppress the hypothalamo-pituitary-adrenal axis. *Aliment Pharmacol Ther* 1994; 8(2):255–258. doi:10.1111/j.1365-2036.1994.tb00284.x
66. Kumana CR, Seaton T, Meghji M, Castelli M, Benson R, Sivakumaran T. Beclomethasone dipropionate enemas for treating inflammatory bowel disease without producing Cushing's syndrome or hypothalamic pituitary adrenal suppression. *Lancet* 1982; 1(8272):579–583. doi:10.1016/s0140-6736(82)91747-0
67. González Villarroel P, Fernández Pérez I, Páramo C, et al. Mestrol acetate-induced adrenal insufficiency. *Clin Transl Oncol* 2008; 10(4):235–237. doi:10.1007/s12094-008-0188-7
68. Mehta K, Weiss I, Goldberg MD. Megace mystery: a case of central adrenal insufficiency. *Case Rep Endocrinol* 2015; 2015:147265. doi:10.1155/2015/147265
69. Thomas CP, Liu KZ, Vats HS. Medroxyprogesterone acetate binds the glucocorticoid receptor to stimulate alpha-ENaC and sgk1 expression in renal collecting duct epithelia. *Am J Physiol Renal Physiol* 2006; 290(2):F306–F312. doi:10.1152/ajprenal.00062.2005
70. Kontula K, Paavonen T, Luukkainen T, Andersson LC. Binding of progestins to the glucocorticoid receptor. Correlation to their glucocorticoid-like effects on in vitro functions of human mononuclear leukocytes. *Biochem Pharmacol* 1983; 32(9):1511–1518. doi:10.1016/0006-2952(83)90474-4
71. Malik KJ, Wakelin K, Dean S, Cove DH, Wood PJ. Cushing's syndrome and hypothalamic-pituitary-adrenal axis suppression induced by medroxyprogesterone acetate. *Ann Clin Biochem* 1996; 33(pt 3):187–189. doi:10.1177/000456329603300302
72. Kling MA, Roy A, Doran AR, et al. Cerebrospinal fluid immunoreactive corticotropin-releasing hormone and adrenocorticotropin secretion in Cushing's disease and major depression: potential clinical implications. *J Clin Endocrinol Metab* 1991; 72(2):260–271. doi:10.1210/jcem-72-2-260
73. Gold PW, Chrousos GP. The endocrinology of melancholic and atypical depression: relation to neurocircuitry and somatic consequences. *Proc Assoc Am Physicians* 1999; 111(1):22–34. doi:10.1046/j.1525-1381.1999.09423.x
74. Opp M, Obál F Jr, Krueger JM. Corticotropin-releasing factor attenuates interleukin 1-induced sleep and fever in rabbits. *Am J Physiol* 1989; 257(3 pt 2):R528–R535. doi:10.1152/ajpregu.1989.257.3.R528
75. Dixon RB, Christy NP. On the various forms of corticosteroid withdrawal syndrome. *Am J Med* 1980; 68(2):224–230. doi:10.1016/0002-9343(80)90358-7
76. Di Dalmazi G, Berr CM, Fassnacht M, Beuschlein F, Reincke M. Adrenal function after adrenalectomy for subclinical hypercortisolism and Cushing's syndrome: a systematic review of the literature. *J Clin Endocrinol Metab* 2014; 99(8):2637–2645. doi:10.1210/jc.2014-1401
77. Zhang CD, Li D, Singh S, et al. Glucocorticoid withdrawal syndrome following surgical remission of endogenous hypercortisolism: a longitudinal observational study. *Eur J Endocrinol* 2023; 188(7): 592–602. doi:10.1093/ejendo/lvad073
78. Hurtado MD, Cortes T, Natt N, Young WF Jr, Bancos I. Extensive clinical experience: hypothalamic-pituitary-adrenal axis recovery after adrenalectomy for corticotropin-independent cortisol excess. *Clin Endocrinol (Oxf)* 2018; 89(6):721–733. doi:10.1111/cen.13803
79. Pofi R, Feliciano C, Sbardella E, et al. The short synacthen (corticotropin) test can be used to predict recovery of hypothalamic-pituitary-adrenal axis function. *J Clin Endocrinol Metab* 2018; 103(8):3050–3059. doi:10.1210/jc.2018-00529
80. Sagar R, Mackie S, Morgan AW, Stewart P, Abbas A. Evaluating tertiary adrenal insufficiency in rheumatology patients on long-term systemic glucocorticoid treatment. *Clin Endocrinol (Oxf)* 2021; 94(3):361–370. doi:10.1111/cen.14405
81. He X, Findling JW, Auchus RJ. Glucocorticoid withdrawal syndrome following treatment of endogenous Cushing syndrome. *Pituitary* 2022; 25(3):393–403. doi:10.1007/s11102-022-01218-y

Address: Noura Nachawi, MD, 24 Frank Lloyd Wright Drive, Ste 1300, Lobby C, Ann Arbor, MI 48105; nachawin@med.umich.edu