

EXPANDED TABLE 4
Summary of patients with acquired hemophilia A and thrombosis

Study number	Authors	Age/sex	Risk factors for thrombosis	Bleeding site	Site of thrombosis	At diagnosis		Inhibitor, Bethesda units/mL	Hematologic treatment	Immunosuppression regimen
						Activated partial thromboplastin time, seconds	Factor VIII			
1	Poli et al ¹⁸	36/F	Pregnancy	Ecchymosis	Distal right lower extremity deep vein thrombosis	57 at deep vein thrombosis diagnosis and on anticoagulation	1.9 U/dL	NA	Vitamin K	Prednisone 1 g/kg/day (4 weeks)
2.	Deitcher et al ¹⁹	60/F	Idiopathic	Forearm hematoma	Right peroneal deep vein thrombosis	54.3 (24–33)	5.3	57	Factor VIII concentrate and desmopressin	NA
		80/M	Lymphoma	Left flank and thigh hematoma	Proximal left limb deep vein thrombosis	61.4 (22–33)	2	47	Factor VIII concentrate and desmopressin	NA
		76/F	Postoperative	Ecchymosis	Proximal left lower limb deep vein thrombosis	54.8 (22–33)	2	5	Factor VIII concentrate and desmopressin	NA
3.	Spencer et al ²⁰	37/F	Pregnancy	Right calf hematoma and upper extremity bruising	Proximal right lower limb deep vein thrombosis	3.01	2 U/dL	35	Recombinant activated factor VII	Prednisolone 1 mg/kg/day (6 weeks) and cyclophosphamide 100 mg/day (6 months)
4.	Paudel et al ²¹	21/M	Trauma/surgery	Abdominal hematoma	Proximal right lower limb deep vein thrombosis, bilateral cephalic vein thrombi	> 100	< 1	12	Vitamin K, Fresh frozen plasma, recombinant activated factor VII	Methylprednisolone 120 mg/day, then rituximab 375 mg/m ² /weekly for 4 doses
5.	Wool et al ²²	73/M	Idiopathic	Thalamic hemorrhage	Pulmonary embolism	76.9 (24–34)	4	27	Factor VIII concentrate (2 days), then recombinant activated factor VII (30 µg/kg/dose), then therapeutic plasma exchange	Dexamethasone 8 mg/12 hours; therapeutic plasma exchange days 5 and 6, then rituximab 375 mg/m ² /weekly (2 doses)
6.	Maral et al ²³	46/F	Malignancy	Ecchymosis	Inferior vena cava thrombosis	107 (22–34)	3	350	NA	Prednisolone 1 mg/kg/day and cyclophosphamide 500 mg/week
7.	Chhabra et al ²⁴	72/M	Idiopathic	Retroperitoneal hematoma	Left leg distal deep vein thrombosis	78.6	< 1	82	Recombinant activated factor VII	Prednisolone (1 mg/kg/day) and cyclophosphamide (100 mg/day)
8.	Aslam et al ²⁵	64/F	Addison disease	Left thigh	Pulmonary embolism	81.8	< 6	NA	Factor VIII inhibitor bypassing activity, vitamin K	Intravenous methylprednisolone (30 mg/12 hours)

NA = not available